

DOCKETED	
Docket Number:	21-AFC-02
Project Title:	Willow Rock Energy Storage Center
TN #:	264487
Document Title:	COMMENTS OF CALIFORNIA UNIONS FOR RELIABLE ENERGY ON THE PRELIMINARY STAFF ASSESSMENT ATTACHMENT D (4 OF 7)
Description:	N/A
Filer:	Alisha Pember
Organization:	California Unions for Reliable Energy
Submitter Role:	Intervenor
Submission Date:	6/27/2025 1:51:50 PM
Docketed Date:	6/27/2025

STATE OF CALIFORNIA
STATE ENERGY RESOURCES CONSERVATION
AND DEVELOPMENT COMMISSION

IN THE MATTER OF:

WILLOW ROCK ENERGY STORAGE
CENTER

Docket No. 21-AFC-02

COMMENTS OF CALIFORNIA UNIONS FOR RELIABLE ENERGY
ON THE PRELIMINARY STAFF ASSESSMENT

ATTACHMENT D (4 OF 7)

June 16, 2025

Richard M. Franco
Adams Broadwell Joseph & Cardozo
601 Gateway Blvd., Suite 1000
South San Francisco, CA 94080
(650) 589-1660
rfranco@adamsbroadwell.com

Attorneys for California Unions for Reliable
Energy




PHOTO ON PREVIOUS PAGE: Sonoran Desert toad, Cochise County, Arizona. Courtesy of Rob Schell Photography.

and have a conspicuous black keratinized spade on the ventral surface of the rear feet.

Taxonomic Relationships

The validity of this taxon has never been questioned, although confusion about the type specimen and locality has been discussed (Fouquette 1968). Osteological and genetic data, as well as call characteristics, suggest that it is related to Central American bufonids (Tihen 1962, Martin 1972, Sullivan and Malmos 1994, Pauly et al. 2004, Frost et al. 2006a).

Frost et al. (2006a) suggested a taxonomic revision that placed this species in the genus *Cranopsis*. Based on subsequent discoveries of older available names, the genus name for this taxon was later revised to *Ollotis* (Frost et al. 2006b) and then *Incilius* (Frost et al. 2009b, Pauly et al. 2009). We retain the older taxonomy both for taxonomic stability and because the analyses supporting the original rearrangement are controversial (Crother 2009, Frost et al. 2009a, Pauly et al. 2009).

Life History

The life history of this species in California is unknown, and we base the following discussion on observations from other areas. *Bufo alvarius* spends much of the year underground, presumably in rodent burrows (Degenhardt et al. 1996). *Bufo alvarius* is primarily nocturnal and becomes active before summer rains. It is more strongly aquatic than most North American toads (Stebbins 1951). Breeding behavior appears to be generally associated with summer rains (Sullivan and Malmos 1994), although amplexus has been reported in stock ponds before rains have occurred (Degenhardt et al. 1996). Several years may pass between breeding events depending on the presence of sufficient rainfall (Sullivan and Fernandez 1999). The species sometimes congregates in large numbers for breeding, with nearly all reproduction of a local breeding population occurring in a single night (Degenhardt et al. 1996). The time required for hatching and metamorphosis is unknown but may be less

than a month (notes of Thornber, reported in Ruthven 1907 and Storer 1925). This species appears to be a dietary generalist, feeding on any live arthropod or small vertebrate prey that it can successfully capture (Stebbins 1951, Cole 1962). Poison secreted by the skin and paratoid glands is particularly toxic and has caused death and paralysis in dogs and is a potent hallucinogen in humans (Musgrave 1930, Stebbins 1951, Stebbins 2003).

Habitat Requirements

The habitat requirements for *Bufo alvarius* in California are unknown. In arid habitats of Arizona and New Mexico, the species can be found in and around a variety of water sources used for breeding, including springs, stock ponds, washes, river bottoms, and irrigation ditches (Stebbins 1951, Stebbins 1972), though it is occasionally found at great distances (>1 mi) from water (Slevin 1928). Upland habitat surrounding known aquatic breeding localities elsewhere in the range include mesquite-creosote desert lowland, arid grassland, rocky riparian zones, oak-sycamore-walnut assemblages in mountain canyons, and montane pine-oak juniper plant communities (Stebbins 2003, Fouquette et al. 2005).

Distribution (Past and Present)

There are no known extant populations in California. Historically, the species ranged in California along the Lower Colorado River and into the Imperial Valley (Grinnell and Camp 1917, Stebbins 1951, Jennings and Hayes 1994a), likely ranging as far north as the southern tip of Nevada (Cooper 1869). It is not known if records in the Imperial Valley are a natural part of the historic range or whether they represent recent range expansion following the development of irrigation (Stebbins 1951).

The last verified record (LACM 87044) from California dates to 31 July 1955, 7 km north of Winterhaven. More recent surveys have failed to detect the species (King and Robbins 1991, Jennings and Hayes 1994b). Sporadic records continue to be reported on the Arizona side of

the Colorado River, however. Several individuals were found near the Cibola National Wildlife Refuge in 1980 (Anderson and Ohmart 1982; B. Anderson, pers. comm.), and a single individual was found at the refuge itself in 1986 (J. Rorabaugh, pers. comm.). On 1 July 2004, a large individual was found "by the golf course on the Parker Strip," La Paz County, Arizona (J. Rorabaugh, pers. comm.). On 29 July 2009, an amplexing pair was found along the Bill Williams River at Planet Ranch, Mohave County, Arizona, and the species is reportedly "fairly common" 24–32 km above the confluence of the Gila and Colorado Rivers, Yuma County, Arizona (J. Rorabaugh, pers. comm.). A single, unverified record of a calling *Bufo alvarius* was reported near Bard, California, in the spring of 2007 or 2008, though the time of year was unexpected and the observer was inexperienced with the species (J. Rorabaugh, pers. comm.).

Outside of California, *B. alvarius* ranges across southern Arizona to the southwestern corner of New Mexico and south into Sonora and the northern edge of Sinaloa, Mexico. The known elevational range extends from near sea level to 1615 m (Cole 1962).

Trends in Abundance

Though the paucity of records from California makes assessing former abundance difficult, *Bufo alvarius* was apparently common at Yuma, Arizona, on the California border, along the Lower Colorado River, and in parts of the Imperial Valley (Slevin 1928, Klauber 1934). As no populations are currently known in these areas, declines leading to probable population extirpations or extremely low population sizes have clearly occurred. The species is also known to be declining in New Mexico (Degenhardt et al. 1996). Throughout the rest of the range the species appears to be stable and abundant at many localities (Fouquette et al. 2005, Lazaroff et al. 2006), though some have suggested that declines are occurring throughout the range (B. Brattstrom, R. Ruibal, and C. Schwalbe, pers. comms., reported in Jennings and Hayes 1994a).

Nature and Degree of Threat

The causes of declines, and therefore the threats to this species, are poorly understood. Declines occurred before any studies were carried out in California, though it is likely that landscape modification and pesticide applications that occurred with the growth of agriculture in the Imperial Valley contributed to declines (Ohmart et al. 1988, Jennings and Hayes 1994a). Bufonids are generally very susceptible to amphibian declines (Stuart et al. 2004). In California, toad declines have been linked to habitat loss and pesticide use (Davidson et al. 2002) and pathogenic fungi (Green and Kagarise Sherman 2001).

Status Determination

The declines and possible extirpation of *Bufo alvarius* in California are the primary concerns for this taxon. The species may require permanent aquatic environments making it a moderate ecological specialist, given the arid environments that characterize its range. This also makes the taxon sensitive to the effects of climate change, particularly changes in hydrology and the increasing year-to-year variation in precipitation that have been projected (Cayan et al. 2008b). Finally, because little understanding of the causes of declines in California exists, we are poorly positioned to protect any remaining populations should they be found in future surveys.

Management Recommendations

The development of an effective management strategy for *Bufo alvarius* in California is not possible without further distributional and ecological information. As no populations are currently known, the first management priority should be to undertake comprehensive surveys, as described below, aimed at identifying remaining fragmentary California populations. Habitat protection and enhancement would then become the critical management tools to build these populations to larger and viable sizes. Simultaneous ecological research is also needed on habitat use, home range size, life

history, and population connectivity before more complex management programs focused on reestablishing the species are considered.

Monitoring, Research, and Survey Needs

A critical first step toward developing a comprehensive management plan for this species is to undertake comprehensive surveys of remaining potential habitat in southeastern California. These surveys should take place during the summer rains and should involve biologists who are familiar with *Bufo alvarius*' breeding behavior. If any remaining populations are found, a population-monitoring program should rapidly be established to determine both geographical extent and population size. As little is known about this species in California, this monitoring program should take place in conjunction with a study of the species' life history and habitat use, in California and/or adjacent Arizona. These surveys should specifically target the remaining moist areas of the southwest California deserts that are known to support other water-dependent vertebrate species, such as the desert mule deer (*Odocoileus hemionus crooki*). Using existing survey data from other, and better known, species may help to guide toad survey efforts

toward the wettest areas or most consistent water supplies, thereby increasing odds of detection.

A second critical priority is to work with wildlife managers in Arizona to survey for and study the nearest remaining populations on the Arizona side of the Lower Colorado River. These populations are likely the most ecologically similar to the former California populations and should therefore provide information valuable to the eventual development of management programs in California. Genetic samples from both California and Arizona should be collected to help inform managers about levels of genetic differentiation, and therefore the appropriateness of possible reintroduction of Arizona animals to California.

As any populations that remain in California are likely isolated, study of these populations is unlikely to yield information on the metapopulation dynamics that we presume are key in sustaining this species elsewhere. Reestablishing these dynamics would form an important part of a comprehensive management program in California, and research focused on better understanding these dynamics will also need to take place outside of California, preferably in adjacent Arizona.



ARROYO TOAD

Bufo californicus Camp 1915

Status Summary

Bufo californicus is a Priority 1 Species of Special Concern, receiving a Total Score/Total Possible of 93% (102/110). During the previous evaluation, it was also considered a Species of Special Concern (Jennings and Hayes 1994a), and it has been listed as federally Endangered since 1995.

Identification

Bufo californicus is a small to medium-sized (4.6–8.6 cm SVL), light-gray to tannish-brown toad that often has some greenish or olive and dark-brown mottling on the back and sides (Camp 1915, Stebbins 2003). The underside is buff or dirty white and usually unmarked (Stebbins 2003). A light middorsal stripe is rarely present (Jennings and Hayes 1994a, Stebbins 2003). Weak cranial crests are often present and the paratoids are oval-shaped and widely separated (Stebbins 2003). The advertisement call of this species is a musical trill that lasts 3–10 s. The pitch of the call rises quickly and is held constant for the

remainder of the call, which ends abruptly (Stebbins 2003, Elliott et al. 2009). Like most toads, the tadpoles are small and black early in life. However, several weeks post-hatching they develop a cryptic tan coloration that closely

Arroyo Toad: Risk Factors

Ranking Criteria (Maximum Score)	Score
i. Range size (10)	10
ii. Distribution trend (25)	25
iii. Population concentration/ migration (10)	10
iv. Endemism (10)	7
v. Ecological tolerance (10)	10
vi. Population trend (25)	20
vii. Vulnerability to climate change (10)	10
viii. Projected impacts (10)	10
Total Score	102
Total Possible	110
Total Score/Total Possible	0.93




PHOTO ON PREVIOUS PAGE: Arroyo toad, Baja California, Mexico. Courtesy of Rob Schell Photography.

matches the substrate (Sweet 1992, Hancock 2009).

Metamorphosed individuals of this species may be confused with the western toad (*B. boreas*), which is the only sympatrically occurring toad. *Bufo boreas* has a prominent white or cream dorsal stripe and lacks cranial crests (Stebbins 2003). Young tadpoles that still retain the black coloration are difficult to distinguish from *B. boreas*, but older tadpoles are readily distinguishable.

Taxonomic Relationships

Until recently, *Bufo californicus* was considered a subspecies of the Arizona toad (*B. microscaphus*) (Price and Sullivan 1988), although recommendations to recognize it as a full species date back to Myers (1930). Frost and Hillis (1990) recognized this species as distinctive based on the general observation that few allopatrically distributed polytypic species represent single genetically cohesive units, as is implied by retaining *B. californicus* as a subspecies under *B. microscaphus*. Later analyses of allozyme data confirm that *B. californicus* is a distinct lineage, providing support for its recognition as a full species (Gergus 1998). Additional analyses of advertisement calls indicated a substantial amount of variation within the species complex, although the results were equivocal with respect to species status (Gergus et al. 1997). Lovich (2009a) analyzed data from two mitochondrial genes and found additional evidence that *B. californicus* is a distinct species. This work also identified clades within *B. californicus* that roughly correspond to parts of the range north and south of the Los Angeles Basin, respectively.

Frost et al. (2006a) recommended placing this species and many other North American bufonids in the genus *Anaxyrus*, although this proposal and the analyses that support it are controversial (Crother 2009, Frost et al. 2009a, Pauly et al. 2009). We choose not to follow this recommendation at the present time, pending further analyses, and to maintain taxonomic stability.

Life History

Bufo californicus is primarily nocturnal and feeds predominantly on nocturnally active ant species (Cunningham 1962, Sweet 1992, Sweet 1993, Mahrtdt et al. 2002). Adults typically emerge from retreats approximately 30–40 min after sunset, remaining active down to temperatures of around 13°C on dry nights and 10°C on rainy nights, with nocturnal activity increased during wet periods (Cunningham 1962, Sweet 1992, Sweet 1993).

Males begin calling at varying times of the year depending on local conditions and elevation, although calling activity appears to initiate when water temperatures reach or exceed 11–13°C (Myers 1930, Sweet 1992). Choruses generally begin in late February in coastal populations and late March or April at higher elevation inland sites, and they may continue into July (Sweet 1992, Sweet 1993, Stebbins 2003, Hancock 2009). Eggs are laid near the male's calling site on a substrate of mud, sand, or gravel, away from vegetation and other submerged debris (Sweet 1992). Hatching occurs after 4–6 days at typical water temperatures (12–16°C), although the larvae remain associated with the egg mass for an additional 5–6 days. Metamorphosis can occur in as few as 65 days, although typically 72–80 days are required (Sweet 1992, Hancock 2009). Larger males and females are more sedentary and tend to breed in the same pools throughout the reproductive season and from year to year (Sweet 1993, Hancock 2009, Mitrovich et al. 2011). The seasonal activity period for adults extends roughly from the beginning of the breeding season to late June or July, after which most toads become inactive (Cunningham 1962, Sweet 1993, Hancock 2009). Juveniles may remain active into October or later following rains (Sweet 1993).

Bufo californicus usually attain reproductive condition in their second (males) or third (females) year. The species is relatively short-lived, with few toads living beyond 5 years of age. In the absence of nonnatural disturbances, survivorship of adult toads is high during the active season, but decreases markedly during

the inactive season. Sweet (1993) documented that toads experience ~55% per year mortality mostly during the winter, though other estimates suggest even higher mortality (D. Holland and N. Sisk, unpublished data, reported in Sweet and Sullivan 2005). Eggs and young larvae are apparently unpalatable to most predators, although garter snakes and nonnative fishes prey upon older tadpoles (Sweet 1992). Juvenile toads that have not yet adopted the nocturnal activity pattern characteristic of adults also experience high predation pressure (Hancock 2009). Adult toads experience intense predation from introduced bullfrogs in areas where that species occurs (Miller et al. 2012, R. Fisher pers., comm.). In the absence of bullfrogs, adult toads experience much lower predation intensity (Sweet 1993, Hancock 2009).

Habitat Requirements

Along with its close relative *Bufo microscaphus*, *B. californicus* may have the most specialized habitat requirements of any North American anuran (Stebbins 2003). This species requires shallow, slow-moving stream and riparian habitat. In some areas they may occupy first-order streams, although most populations inhabit second- to sixth-order streams that have extensive braided channels and sediment deposits of sand, gravel, or pebbles that are occasionally reworked by flooding (Sweet and Sullivan 2005). These toads will use either permanent or seasonal streams, although seasonal streams must flow for a minimum 4–5 months for successful reproduction and recruitment (Sweet and Sullivan 2005). At inland sites, radiotelemetry studies indicate that this species rarely moves beyond the immediate upland margin of streams, although in coastal sites arroyo toads appear to occasionally use and disperse across hotter and drier upland sites (Sweet 1992, Sweet 1993, Griffin and Case 2001, Hancock 2009, Mitrovich et al. 2011). Mitrovich et al. (2011) found that radio-tracked toads actively selected channel and terrace stream habitats, and largely avoided surrounding scrub, grassland, and forest. On average, males were found

in closer proximity to flowing sections of stream than females, possibly to maximize reproductive opportunity (Mitrovich et al. 2011). *Bufo californicus* is known to occasionally use and breed in human-made habitats, such as artificial stream terraces and ponds (Price and Sullivan 1988, Mahrtdt et al. 2002). It is unknown whether the species can persist in these habitats.

Distribution (Past and Present)

Bufo californicus historically occurred in coastal drainages from the San Antonio River, Monterey County, California, southward through the Transverse and Peninsular Ranges to the vicinity of Arroyo San Simón in Baja California Norte, Mexico (Price and Sullivan 1988, Cergus et al. 1997, Grismer 2002, Lovich 2009a). Almost all populations occur along the coast or on the coastal slopes of the southern California mountains. Six localities were previously recognized from the desert slopes of Los Angeles, Riverside, San Bernardino, and San Diego Counties, California (Patten and Myers 1992, Jennings and Hayes 1994a). Desert slope populations are known to occur at Little Rock Creek, Los Angeles County, and the Mojave River, San Bernadino County. Populations at Whitewater River, Riverside County, Borrego Springs (listed as San Felipe Creek in Jennings and Hayes 1994a), Vallecito Creek, and Pinto Canyon, San Diego County, are probably in error and are the result of misidentifications (Ervin et al. 2013). The known elevational range extends from near sea level to approximately 1000 m (Stebbins 2003; S. Sweet, pers. comm.).

The present distribution of *B. californicus* is considerably smaller than it once was. Jennings and Hayes (1994a) estimated that this species had disappeared from 76% of its former range in California, although more recent estimates place this loss at 65% (Sweet and Sullivan 2005).

Trends in Abundance

In addition to the extirpations discussed above, extensive declines in abundance have been

documented in most *Bufo californicus* populations that do survive. Extensive collections from the 1930s, largely stemming from the work of L.M. Klauber, suggest that this species was formerly present at much higher densities (S. Sweet, pers. obs., reported in Sweet and Sullivan 2005).

Nature and Degree of Threat

A recent 5-year review of the status of *Bufo californicus* thoroughly discusses the ongoing threats to this taxon (USFWS 2009). We follow the findings of that document and recommend that readers consult it for additional detail.

The greatest threat facing this taxon is loss and degradation of habitat that stems from modifications to hydrology from reservoir construction, roads, flood control, development, recreational activity, and mining (USFWS 2009). In addition, declines are occurring even in areas that are not subject to development and direct habitat degradation from human activities (Hancock 2009). These additional declines stem largely from introduced predators (primarily bullfrogs and green sunfish) and introduced plants, which degrade habitat and/or decrease survivorship of toads (Sweet 1992, Hancock 2009, USFWS 2009, Miller et al. 2012). Off-highway vehicle use has also caused both habitat degradation and direct mortality in this species (Ervin et al. 2006)

Status Determination

Major declines in both distribution and abundance, coupled with several ongoing threats, combine to warrant a Priority 1 Species of Special Concern status for *Bufo californicus*.

Management Recommendations

Management efforts for *Bufo californicus* should mirror those outlined by the USFWS recovery

plan and 5-year review for this taxon (USFWS 1999, USFWS 2009). The recent 5-year review suggests that management efforts to date have been effective, and the outlook for this species has improved somewhat since it was initially listed (USFWS 2009). The most important management strategy is to preserve existing stream habitat that supports this species and to restore additional habitat that can support self-sustaining populations. Restoration efforts should include dam removal to allow streams to meander and rebuild sand and gravel bars, and removal of exotic plants and vertebrate predators.

Monitoring, Research, and Survey Needs

Monitoring, research, and survey needs are covered in depth in the USFWS recovery plan for this taxon and the recent 5-year review. We refer the reader to these documents for additional detail (USFWS 1999, USFWS 2009). Monitoring efforts should focus on recovering populations, particularly those in newly restored habitat. It is particularly important to continue monitoring through drought and El Niño cycles given that this is a short-lived species and several years of consistent drought could be extremely damaging to recovering populations.

In addition, research aimed at characterizing variation in this species' life history in different parts of its range should be undertaken, as these differences might have an impact on future management efforts. For example, the two desert slope populations may differ substantially in several aspects of life history relative to the coastal slope populations. Additional research into the prevalence and potential impacts of *Bd* fungus on this species is also particularly important. Finally, molecular analyses of population size and connectivity might be particularly valuable in this taxon.



YOSEMITE TOAD

Bufo canorus Camp 1916a

Status Summary

Bufo canorus is a Priority 1 Species of Special Concern, receiving a Total Score/Total Possible of 84% (92/110). During the previous evaluation, it was also considered a Species of Special Concern (Jennings and Hayes 1994a). This species is also listed as Threatened under the US Endangered Species Act.

Identification

Bufo canorus is a moderately sized (1.0–7 cm SVL) sexually dichromatic toad (Camp 1916a, Grinnell and Storer 1924, Kagarise Sherman 1980, Stebbins 2003). Females and juveniles have tan or brown dorsal coloration with extensive dark blotches over the dorsal surface and legs that are edged with white or cream. Males are pale green-yellow or olive green above without, or with only minimal, dark blotching or flecking (Camp 1916a, Stebbins 2003). A pale, very thin middorsal line is often present in juveniles and young females, but is usually lost in males and older females (Camp 1916a). The paratoid glands

are large, flat, and circular, and are separated by a space less than or equal to their diameter (Camp 1916a). The advertisement call of this species is a musical trill lasting 3–9 s (Elliott et al. 2009). The specific epithet “*canorus*” (Latin for

Yosemite Toad: Risk Factors

Ranking Criteria (Maximum Score)	Score
i. Range size (10)	10
ii. Distribution trend (25)	15
iii. Population concentration/ migration (10)	10
iv. Endemism (10)	10
v. Ecological tolerance (10)	10
vi. Population trend (25)	20
vii. Vulnerability to climate change (10)	10
viii. Projected impacts (10)	7
Total Score	92
Total Possible	110
Total Score/Total Possible	0.84




PHOTO ON PREVIOUS PAGE: Yosemite toad, Mono County, California. Courtesy of Rob Grasso.

“tuneful”) refers to the melodic quality of the call (Camp 1962a, Karlstrom 1962).

This species may be confused with high-elevation populations of the western toad (*B. boreas*). *Bufo boreas* often has a conspicuous light middorsal stripe and smaller, more widely spaced paratoid glands that are separated by a space approximately twice their diameter (Stebbins 2003). *Bufo boreas* also has more extensive webbing on the hind feet than *B. canorus* (Camp 1917). Populations of *B. boreas* that occur in the Sierra Nevada do not produce advertisement calls; thus, toad breeding choruses are diagnostic for *B. canorus* (Mullally 1956). Juveniles of the two species (<20–30 mm) are very similar to one another, and identifications of this size class should be made with caution (Karlstrom 1962).

Taxonomic Relationships

Bufo canorus was initially described on the basis of coloration and morphology (Camp 1962a). Its status as a distinct species has gone largely unquestioned since this time. Subsequent genetic analyses based on mitochondrial data suggest that this taxon is a close relative of *B. boreas* (Shaffer et al. 2000, Pauly et al. 2004). In addition, mitochondrial DNA data suggest that *B. canorus* may be paraphyletic with respect to the black toad, *B. exsul*, and some lineages of *B. boreas* (Graybeal 1993, Shaffer et al. 2000, Goebel et al. 2009). These relationships have not been corroborated with nuclear sequence data, and thus it is unclear whether cryptic diversity exists within the taxon or if this is a case of mitochondrial introgression. Unpublished genetic data suggest that mitochondrial introgression associated with past or ongoing hybridization may explain these results (G. Pauly, unpublished data). Some morphological variation has been observed in size and degree of melanism in eggs and larvae of this taxon (Karlstrom and Livezey 1955, Karlstrom 1962). Whether this is plasticity in response to elevation and/or the local environment or genetic differentiation has not been investigated further. Mitochon-

drial data also indicate that some intraspecific variation and isolation by distance may exist within the taxon (Shaffer et al. 2000, Pauly et al. 2004, Goebel et al. 2009), although sample sizes in these studies were small. Wang (2012) collected data from 10 microsatellites for toads from 24 populations in Yosemite National Park and found significant variation in the amount of genetic distance between populations. This analysis concluded that environmental factors such as slope and precipitation were associated with genetic structure.

Frost et al. (2006a) recommended placing this species and many other North American bufonids in the genus *Anaxyrus*, although this proposal and the analyses that support it are controversial (Crother 2009, Frost et al. 2009a, Pauly et al. 2009).

Life History

Bufo canorus is primarily a diurnal toad that occasionally exhibits crepuscular or nocturnal activity on warm days (Mullally 1956, Martin 2008). Males emerge from hibernation as soon as snowmelt pools form along the margins of preferred high-elevation meadow habitat and quickly form breeding choruses (Karlstrom 1962). As in many high-elevation amphibians, the timing of emergence is correlated with elevation, and generally occurs in May and June (Karlstrom 1962, Kagarise Sherman 1980). Males are territorial and often maintain interindividual spacing of 7–14 m (Karlstrom 1962, Kagarise Sherman 1980). Fighting occurs between males that encroach on one another's territory (Kagarise Sherman and Morton 1984). Breeding activity and egg-laying commence soon after males begin calling, with females depositing eggs along shallow edges of pools and streams in meadows (Karlstrom 1962, Kagarise Sherman 1980). Hatching occurs in as few as 3–4 days at relatively high water temperatures (20–23°C) or up to 10–14 days at lower temperatures (16–17°C) (Karlstrom 1962, Kagarise Sherman 1980). Metamorphosis occurs approximately 40–60 days after oviposition, again depending on temperature and

elevation (Karlstrom 1962, Kagarise Sherman 1980, Kagarise Sherman and Morton 1984). The seasonal activity period extends into late September and early October, after which toads hibernate in rodent burrows, crevices under rocks, and root tangles (Kagarise Sherman 1980). Adult toads do not begin to breed until they are 3–6 years old, after which females may only breed every few years (Kagarise Sherman 1980, Kagarise Sherman and Morton 1984). Adults grow slowly, averaging only 2.5 mm per year at Tioga Pass, Tuolumne County, California (Kagarise Sherman and Morton 1984). The post-metamorphic diet consists of a variety of small arthropods including ants, bees, flies, wasps, beetles, millipedes, and spiders (Grinnell and Storer 1924, Mullally 1953, Kagarise Sherman and Morton 1984). The slow growth rate and lack of breeding every year is likely attributable to low metabolic rates associated with low caloric intake and relatively cold temperatures (Kagarise Sherman and Morton 1984). In the wild, adults appear to be able to tolerate a relatively wide range of temperatures (from 2°–30°C) (Karlstrom 1962), although they prefer temperatures higher in this range (Cunningham 1963). The estimated critical thermal maximum is 37–40°C for adults and 36–38°C for larvae (Karlstrom 1962).

Bufo canorus is known to occur sympatrically with *B. boreas* in two areas (see the "Distribution" section) and may occasionally hybridize. At the Frog Lakes locality, individuals that are morphologically intermediate in paratoid gland width and the extent of webbing on the hind feet occur and may represent natural hybrids (Morton and Sokolski 1978). No putative hybrids have been described from the Blue Lakes locality (Karlstrom 1962), although some authors suggest that hybridization may also occur there (Stebbins 2003). Artificial crosses in the laboratory readily produce hybrids (Karlstrom 1962).

Habitat Requirements

Bufo canorus prefers relatively open high-elevation meadows vegetated with grasses, sedges, rushes, and/or willow stands (Karlstrom 1962).

This species can be found in the margins of water bodies that form from snowmelt runoff, as well as in moist meadows. During the early part of the active season, individuals are often localized along meadow margins within approximately 30 m of the forest edge. This behavior may allow them to easily retreat to forest cover at night to avoid freezing temperatures (Karlstrom 1962). As the active season progresses and nights become warmer, the toads tend to move toward the center of meadows and become less restricted to the margins (Karlstrom 1962). This species prefers shallow (probably <7.5 cm) snowmelt pools on the margins of meadows or very slow moving runoff streams in which to breed, although they have also been found in deeper (>3 m) permanent pools (G. Fellers, pers. comm.). These need to be deep enough to avoid premature desiccation—a significant cause of mortality for larvae but shallow enough to achieve the temperatures needed for rapid development (Karlstrom 1962). This species may prefer to oviposit in dark bottomed pools, particularly at high elevations, as these may provide warmer water temperatures and more rapid larval development (Karlstrom 1962). The presence of pocket gopher, mouse, and vole burrows may provide additional beneficial cover and protection from predation (Grinnell and Storer 1924, Karlstrom 1962).

Distribution (Past and Present)

Bufo canorus is restricted to a relatively small area approximately 240 km (north–south) by 60 km (east–west) in higher elevation areas of the Sierra Nevada (Karlstrom 1962, Kagarise Sherman and Morton 1993). It ranges from the vicinity of Blue Lakes, Alpine County, California, south past Kaiser Pass to the Evolution Lakes area, Fresno County, California (Grinnell and Storer 1924, Livezey 1955, Karlstrom 1962, Jennings and Hayes 1994a, Stebbins 2003, Davidson and Fellers 2005). The known elevational range extends from 1950 to 3599 m, with most localities between 2590 and 3048 m (Karlstrom 1962).

Between 1915 and 1992, this species exhibited declines throughout some areas of its

range. Drost and Fellers (1996) resurveyed localities from Grinnell and Storer (1924) and found that this species had disappeared from 6 of 13 sites in the Yosemite area. Jennings and Hayes (1994a) also estimated that the species has disappeared from low-elevation areas on the western edge of the range, as well as at the northern edge of the range.

Trends in Abundance

In areas where *Bufo canorus* persists, marked declines in abundance have also been documented. In the Drost and Fellers (1996) resurveys, *B. canorus* was present in lower densities than in 1915 at three sites where it was still present. Between 1976 and 1982, the number of male toads entering breeding pools at Tioga Pass meadow declined from a maximum of 342 individuals to a low of 28, a ninefold decrease from the 1974–1978 mean (Kagarise Sherman and Morton 1993). However, the number of females entering breeding pools showed no obvious changes during this time period (Kagarise Sherman and Morton 1993). The average number of toads encountered in daily surveys also declined in the vicinity of Tioga Pass meadow between the early 1970s and 1990. In addition, these surveys documented declines in female toads, although they were not as severe as those documented in males (Kagarise Sherman and Morton 1993). Similar declines in abundance have also been documented at six additional localities in this region (Kagarise Sherman and Morton 1993).

Nature and Degree of Threat

The causes of decline in *Bufo canorus* require additional study. The declines have occurred in seemingly undisturbed areas and do not appear to be localized, suggesting that they are being driven by general changes to the environment, rather than localized causes such as habitat destruction. Several possible causes have been advanced, and more than one factor may be playing a role. These causes include environmental contamination, disease, drought and/or climate change, habitat modification due to

grazing or other activities, human disturbance of breeding choruses, increased predation pressure from birds and fish, and pesticides. Based on current data it is not possible to understand in detail which, if any, of these factors are most important in *B. canorus* declines.

Snowmelt pools have extremely low acid neutralizing capacity, leading to the hypothesis that acidification of aquatic breeding habitat due to atmospheric deposition may be contributing to declines. Bradford et al. (1992) examined the effect of increasing acidification and the associated increase in dissolved aluminum on embryos and hatchlings of *B. canorus*. Embryos and hatchlings exposed to decreasing pH (and increasing aluminum solute) showed no increase in mortality at levels found in nature. However, these factors did cause earlier hatching and smaller body size at metamorphosis. Bradford et al. (1994) attempted to correlate the distribution of declining Sierran amphibians with these environmental factors and found no relationship, concluding that acid deposition was an unlikely source of amphibian declines in the Sierra Nevada.

Disease has also been considered as a factor in declines, though there is little evidence to date. Green and Kagarise Sherman (2001) examined the cause of death in 12 adult *B. canorus* that were found during a die-off that immediately preceded the population declines documented at Tioga Pass meadow by Kagarise Sherman and Morton (1993). They found that a variety of diseases and parasites were present in the population, and chytridiomycosis and septicemia, alone or in combination, caused the death of at least four individuals. However, no single infectious disease was present in more than 25% of the samples, which is far below the proportion typically observed in other die-offs caused by these diseases (Worthylake and Hovingh 1989, Berger et al. 1998, Vredenburg et al. 2010).

California experienced a relatively severe drought between 1987 and 1992, a time when *B. canorus* population declines were occurring (Roos 1992, Drost and Fellers 1996). Although

it may have played an exacerbating role, drought alone seems unlikely to be responsible for declines. California experiences drought with some regularity, including during the time period of the Grinnell and Storer (1924) survey, which occurred before any major declines in *B. canorus* were observed (Drost and Fellers 1996). Drought does affect year-to-year reproductive success for this species, and prolonged drought may have a cumulative effect on populations (Kagarise Sherman and Morton 1993). Because climate change is expected to impact the amount of snow present in the Sierra Nevada and the speed and timing of snowmelt (Cayan et al. 2008b), drought might play an increasing role in declines of this species in the future.

Habitat modification is a leading cause of decline in many species throughout California and has been suggested as a factor for *B. canorus*. However, *B. canorus* is found largely within the boundaries of Yosemite National Park and other public (mostly National Forest) lands that have experienced varying impacts over the last 100 years. Drost and Fellers (1996) compared photos of habitat from the Grinnell and Storer (1924) surveys with current habitat and saw no apparent differences. Over the course of their 20-year study, Kagarise Sherman and Morton (1993) were also unable to detect significant habitat changes. That said, local impacts from changing habitat remain a potential driver of declines. All-terrain vehicle and snowmobile use in some localized areas may degrade habitat quality (D. Emery, pers. comm.). Some workers have postulated that livestock grazing in alpine meadows of National Forest land causes changes to hydrology, which may affect the suitability of breeding habitat and increase sedimentation in pools. Two recent, relatively short-term studies (5 years) have addressed this hypothesis: one that used experimental fencing treatments to exclude livestock from *B. canorus* breeding meadows and a second that included occupancy surveys across gradients of meadow moisture and livestock use levels (K. Tate and A. Lind, pers. comm.). Both of these studies demonstrated that meadow wetness was more influen-

tial in determining the current distribution and abundance of *B. canorus* than the level of livestock use. Sean Barry (pers. comm.) documented that toads seem to persist and even concentrate in areas that had been disturbed by cattle in the Kaiser Meadow population. It is also possible that the presence of cattle feces increases insect food supply for adult toads, although this remains untested. Martin (2008) suggests that the practice of fencing individual breeding pools to prevent grazing might actually lead to stronger habitat disturbance from cattle grazing in the terrestrial foraging habitats, potentially increasing the overall impact from grazing. These local-scale influences of livestock grazing along with more detailed and longer-term investigations of livestock use in the context of *B. canorus* metapopulation dynamics require further study.

Some researchers have suggested that increasing predation pressure could be causing declines. A variety of avian predators are known to feed on adult and larval toads, and increasing densities of common raven (*Corvus corax*) have been postulated as a possible cause of decline (Kagarise Sherman and Morton 1993). Ravens are known to prey upon other toad species in the *B. boreas* complex and likely also take *B. canorus*. Evidence suggests that declines in other amphibian species have occurred in areas where fish have been introduced (Drost and Fellers 1996). However, fish alone are unlikely to explain the declines in *B. canorus*. Most *B. canorus* reproduction takes place in ephemeral water bodies that do not contain fish (Drost and Fellers 1996). Knapp (2005) found no evidence for an effect of introduced trout on *B. canorus* presence and absence. Further, Grasso (2005) and Grasso et al. (2010) examined the palatability of early life stages of *B. canorus* to introduced brook trout and found that all life stages were highly unpalatable, suggesting that introduced trout may have little direct impact on populations.

Some authors have noted that breeding choruses of *B. canorus* are sensitive to human disturbance. Grinnell and Storer (1924) docu-

mented that choruses would abruptly stop calling when humans entered a meadow. They specifically noted that *B. canorus* seemed to be more sensitive to this disturbance than the sympatric Pacific treefrog (*Pseudacris regilla*). Karlstrom (1962) as well as several biologists presently working on this species disagree that this species' calling behavior is impacted more strongly by human disturbance than other toad species. Karlstrom (1962) did notice wariness at night and that cars moving through the area even at 0.8 km distance would cause choruses to cease calling and that "the almost continual daytime traffic in [Yosemite National Park] might help to explain the paucity of roadside populations" of *B. canorus*. It has also been suggested that the relatively frequent handling and study experienced by some populations could induce stress and immunosuppression, which may also be playing a role in declines (Green and Kagarise Sherman 2001). To our knowledge, this possibility has not been investigated.

Davidson et al. (2002) found that areas where toads had disappeared were downwind from disproportionately large areas of agricultural land (primarily the low-elevation populations on the western side of the Sierra Nevada), suggesting that wind-borne agrochemicals may be a factor in declines. However, this relationship was not statistically significant.

When this evidence is taken together, it is clear that the causes of decline for *B. canorus* are still poorly understood. It is possible that several factors act in combination, perhaps interacting with variation in life history or metapopulation dynamics. Individual populations may be susceptible to localized extirpation due to small population sizes and the species' slow maturation rate. Increasing frequency of localized extirpations could cause a breakdown of broader-scale metapopulation dynamics, leading to additional declines as recolonization ceases to counteract local population extirpations. Landscape genetic data suggest that migration rates between local populations are already low in several areas and this situation would likely be exacerbated by additional

localized population declines and extirpations (Wang 2012). These inferences are still speculative, however, and further research is needed on many aspects of *B. canorus* population biology to better understand ongoing declines.

Status Determination

Declines in both distribution and abundance, coupled with a poor understanding of the factors leading to decline, are the major factors justifying a Priority 1 Species of Special Concern status.

Management Recommendations

An effective management program for this species will depend on identifying and prioritizing the factors leading to observed, ongoing declines. Until this is accomplished, protecting breeding meadows from disturbance of natural hydrologic regimes and water table dynamics and limiting human disturbance to meadows during the breeding season may be helpful in safeguarding populations. In addition, upland wintering habitats adjacent to breeding areas should also be protected from grazing and other disturbances.

Monitoring, Research, and Survey Needs

Ongoing monitoring and study of this species is required with a particular aim of identifying the major factors leading to decline. It is possible that some populations are relatively stable, and comparisons with declining sites could lead to important insights into reasons for declines and potential management solutions. Experimental work, going beyond the primarily correlational studies that have been carried out thus far, could also be helpful in identifying the most important factors. In particular, experimental studies of human disturbance, susceptibility to disease, and the potential role of reduced snowpack on hibernation and breeding biology would all be useful. Populations should also be monitored for disease outbreaks.

Further genetic work also needs to be completed to characterize genetic diversity within the species. Several studies have already been

carried out, although they rely primarily on mitochondrial data alone, which is unable to distinguish true population substructure (or multiple lineages) from introgression from nearby *B. boreas* populations. Wang (2012) adds important information from the nuclear genome, but focuses on Yosemite National Park rather than the species' range as a whole. Future studies should utilize multiple unlinked nuclear markers to clarify the diversity present

in the species, gene flow among meadows, and effective population sizes.

Finally, the majority of survey efforts to date have focused on populations within the boundaries of Yosemite National Park. A committed survey effort is needed to better understand the location of populations, their trends in distribution and abundance, and their disease status and level of infection (or lack of) in areas outside of the park itself.



NORTHERN RED-LEGGED FROG

Rana aurora Baird and Girard 1852

Status Summary

Rana aurora is a Priority 2 Species of Special Concern, receiving a score of 55% (61/110). Previously this species was included as a Species of Special Concern by Jennings and Hayes (1994a).

Identification

Rana aurora is a medium-sized (70–100 mm SVL) brown, red, gray, or tan frog with dorsolateral folds (Dumas 1966, Nussbaum et al. 1983). The dorsum varies from having indistinct, irregular black spots 2–3 mm in diameter with many tiny flecks to an all-over network pattern of black lines (Dunlap 1955, Dumas 1966, Nussbaum et al. 1983). The dorsum can also be largely unmarked, though this is less common. A light lip line from eye to shoulder is usually present, often with a dark mask above (Nussbaum et al. 1983). Ventrally, the chest and abdomen are often marbled gray, with the groin area heavily and darkly mottled (Dunlap 1955). Red coloration on the venter and underneath the

hind legs is typical but varies in intensity and extent (Dunlap 1955). Yellow coloration is common in the groin, as well as red and sometimes green coloration (Dunlap 1955). Larvae are dark brown from above, with scattered small clumps

Northern Red-Legged Frog: Risk Factors

Metric (Maximum Score)	Score
i. Range size (10)	10
ii. Distribution trend (25)	10
iii. Population concentration/ migration (10)	10
iv. Endemism (10)	0
v. Ecological tolerance (10)	7
vi. Population trend (25)	10
vii. Vulnerability to climate change (10)	7
viii. Projected impacts (10)	7
Total Score	61
Total Possible	110
Total Score/Total Possible	0.55




PHOTO ON PREVIOUS PAGE: Northern red-legged frog, Humboldt County, California. Courtesy of William Flaxington.

of metallic flecks and are 50–75 mm in TL (Nussbaum et al. 1983). In southern Mendocino County, this species could be confused with *R. draytonii* (Shaffer et al. 2004). *Rana draytonii* is a larger frog (up to 138 mm SVL; Stebbins 2003), and typically its dark dorsal markings have light centers (Stebbins 2003).

Taxonomic Relationships

Although initially described as a distinct species (Baird and Girard 1852), for much of the twentieth century northern red-legged frogs were regarded as the subspecies *Rana aurora aurora* (Camp 1917). Studies over the last few decades have supported the current taxonomic arrangement, with *R. aurora* and *R. draytonii* both recognized as distinct species. While they look superficially similar, these two species differ in morphology (vocal sacs, body size) and breeding behavior, and are genetically distinct (Hayes and Miyamoto 1984, Green 1985, Green 1986a, Green 1986b, Hayes and Kremples 1986, Shaffer et al. 2004). A narrow contact zone occurs between *R. aurora* and *R. draytonii* in southern Mendocino County (Shaffer et al. 2004).

Life History

Limited information is available on *Rana aurora* life history, with most studies occurring outside of its California range. Adults migrate to wetlands to breed for a few weeks between December and April when temperatures range from 4°C to 18°C (Storm and Pimentel 1954, Storm 1960, Dumas 1966, Licht 1969, Calef 1973). Males call beneath the water (Licht 1969, Brown 1975b). During one breeding season in Humboldt Bay National Wildlife Refuge, two breeding events more than a month apart produced over half of the egg masses (J. Betasso et al., unpublished data). Egg masses are 15–25 cm in diameter and contain approximately 500–600 eggs on average (Calef 1973, Licht 1974, Brown 1975b). Eggs are attached to emergent and floating vegetation, branches, or logs up to 150 cm below the water surface (Brown 1975b, Storm 1960, Calef 1973, Cary 2010). Surveys in Humboldt Bay National Wildlife Refuge found that

most egg masses occurred between 30 and 60 cm elevation in the stream channel, at an average height of 37 cm above the bottom and 8 cm below the water surface (J. Betasso et al., unpublished data). In southwestern British Columbia, eggs were found at least a meter away from the pond edge or river bank (Licht 1969). Water temperatures near developing eggs in a Washington pond were 6.2°C on average (Brown 1975b), and embryos tolerate temperatures from 4°C to 21°C (Licht 1971). Dumas (1966) reared embryos at 11°C, 15°C, and 20°C, and observed the greatest embryo mortality at 20°C.

Embryonic development (from laying to hatching) takes 34–49 days (Storm 1960, Licht 1971, Brown 1975b). In Humboldt Bay Wildlife Refuge, most egg masses (103/232) persisted for 4 weeks before completely hatching out (J. Betasso et al., unpublished data). Larvae hatch at 8–12 mm long (Storer 1925, Storm 1960, Brown 1975b) and grow to up to 80 mm TL (Brown 1975b). Metamorphosis occurs after 3–7 months (Brown 1975b, Storer 1925) and metamorphs are 18–29 mm (Brown 1975h, Storm 1960, Calef 1973). Larger metamorphs are more likely to survive and to emigrate farther (Chelgren et al. 2006). In a Washington population, eggs were laid in February and March, the first larvae hatched in April, and metamorphosis was completed in late July (Brown 1975b).

Rana aurora juveniles disperse from breeding sites within days or weeks after transformation (Licht 1974, Licht 1986a). While daily movements of adults may be on average only a few meters per day, movements of several hundred meters to 4.8 kilometers have been documented over longer periods (Haggard 2000, Hayes et al. 2001, Chan-McLeod and Wheeldon 2004, Hayes et al. 2007).

Larvae are algal grazers (Dickman 1968). Metamorphs and adults are generalist predators of insects, spiders, and mollusks (Licht 1986b).

Habitat Requirements

Rana aurora occurs in mesic forests and riparian areas, which in its northern California range are primarily steep coniferous forests,

coastal terraces, and floodplains (Nussbaum et al. 1983, Stebbins 2003). *Rana aurora* is relatively terrestrial for a ranid frog. Adults can occur hundreds of meters from water, and are often found in dense vegetated or downed log cover (Dunlap 1955, Dumas 1966). Adult frogs radio-tracked from March to July in Humboldt County were detected on land 90% of the time and usually within 5 m of water, though animals were found up to 80 m away from water (Haggard 2000). In habitat choice experiments, juvenile frogs spent most of their time out of the water (Pearl et al. 2004).

Both permanent and temporary breeding habitats are used, such as ponds, freshwater lagoons, lakes, and slow-moving streams (Licht 1969, Cary 2010, Sun 2012). Artificial habitats such as drainage ditches are also used (T. Fuller, J. Garwood, and M. van Hatterm, pers. comm.). Coastal streams may be important dispersal corridors to inland populations. For example, *R. aurora* have been found outside of the breeding season in coastal streams in Humboldt and Del Norte Counties, and egg masses have been found in backwaters and alcoves of the Smith River where surrounding areas have been diked, drained, and converted (J. Garwood, pers. comm.). Both aquatic and terrestrial vegetation are important determinants of breeding habitat quality. In Humboldt County, egg mass presence was positively correlated with low canopy cover (ponds with less than ~40% canopy cover are more likely to have egg masses present; Cary 2010). Egg mass density was higher in smaller ponds (< 2000 m²) and in ponds where the percentage of floating and emergent vegetation cover was at least ~40% (Cary 2010). Surveys in Oregon also found support for the importance of emergent vegetation, as wetlands used for breeding had 27% open water on average compared to 50% open water in unused wetlands (Pearl et al. 2005a). Occupancy models fit to 5 years of survey data in Oregon predicted that local extinction probability decreased as the percentage of trees along the shoreline increased and surface area of emergent vegetation increased (Adams et al. 2011).

Distribution (Past and Present)

Rana aurora occurs from Mendocino County, California, north along the west side of the Cascade Crest up through Vancouver Island and the adjacent mainland coast of British Columbia (Stebbins 2003). Populations also occur on Graham Island, British Columbia (Ovaska et al. 2002), and on Chichagof Island, Alaska (Hodge 2004). The elevational range extends from near sea level to 1160 m in Lane County, Oregon (Dunlap 1955), with populations in California occurring up to approximately 300 m (Jennings and Hayes 1994a). Two localities included on our map possibly extend the eastern edge and elevation range in California, and are in need of further investigation. A specimen collected by Camp in 1913 from eastern Mendocino County is in the UC Berkeley collection (MVZ 5068), photographs of which were reviewed by several experts. It is possible that the specimen is a misidentified *R. draytonii*, or it may be that *R. aurora* was historically more widespread. Despite the presence of potentially suitable habitat, contemporary CDFW biologists working in this region have not observed any *R. aurora* east of Highway 101 or in Mendocino National Forest (T. Fuller, pers. comm.). At another site, two individuals were found recently in eastern Humboldt County at around 800 m elevation (M. van Hatterm, pers. comm.).

In California, surveys have found *R. aurora* to be mostly absent from the river bottom lands of the Eel, Mad, and Smith Rivers. These areas have undergone extensive habitat conversion to beef, dairy, and bulb farming, though populations may persist on inaccessible private lands (M. van Hatterm, unpublished data). Surveys in Oregon's Willamette valley found *R. aurora* at 50% of sites, with highest occupancy probability observed in seasonal sites without fish (Rowe and Garcia 2013).

Trends in Abundance

Population declines have been suspected for *Rana aurora*, particularly in Oregon's Willamette Valley (e.g., Nussbaum et al. 1983; Hayes and

Jennings 1986). However, systematic surveys are lacking. Data on *R. aurora* abundance in California are limited, particularly with regard to documenting trends over time. Mean density of egg masses in breeding ponds in Humboldt County during one breeding season was 0.2/m², with densities up to 0.7/m² observed (Cary 2010). In Del Norte County, 382 egg masses were found in a 40 × 40 m area of a pond near the confluence of East Fork Mill Creek and West Branch Mill Creek (J. Garwood, unpublished data). Surveys in California have found more egg masses in areas where natural vegetation buffers the breeding habitat compared to developed areas (M. van Hattem, unpublished data).

Nature and Degree of Threat

The major threat to *Rana aurora* is development and forest conversion leading to habitat loss and degradation. Other threats include introduced predators, disease, and climate change, though more data are needed on each of these stressors.

Due to issues such as low capture rates, it is unclear whether *R. aurora* abundance varies consistently with stand age in harvested forests (reviewed in Pearl 2005). For example, terrestrial (Welsh et al. 2007) and aquatic (Ashton et al. 2006) amphibian surveys in northwestern California forests have documented only a handful of *R. aurora*. In Washington, breeding sites with high primary forest cover within 2 km had higher egg mass counts, as did breeding sites greater than 0.25 km away from roads (Holcomb 2012). On Vancouver Island, radio-tracked frogs tended to move toward old-growth stands and away from clear-cuts <12 years old, suggesting that recolonization of impacted sites may require several years (Chan-McLeod 2003). In an Oregon study, the highest capture rates of *R. aurora* were in mature, mixed large sawtimber forest (Martin and McComb 2003).

Agricultural and residential development has likely contributed to habitat loss and degradation for *R. aurora*, and is projected to continue to increase in the future. For example, much of the Smith River coastal plain in Del Norte

County has been converted to lily bulb production (J. Garwood, pers. comm.). In addition to habitat loss, such agricultural conversion can further degrade habitat through use of chemicals such as pesticides, herbicides, and fungicides. Similarly, the emerging issue of largely unregulated marijuana cultivation can degrade watersheds through grading and roadbuilding (which both destroy habitat and create runoff into aquatic habitats), application of pesticides and herbicides, and through dewatering of springs, streams, and wetlands used for irrigation (e.g., Thompson et al. 2014). Residential and commercial development is likely to increase in northern California, potentially leading to losses of breeding habitat or loss of access to remaining habitat. For example, the Humboldt County General Plan is currently being updated, with some proposals considering a doubling or tripling of rural development. However, *R. aurora* does use artificial habitat for breeding, and amount of urban cover was not a strong predictor of frog occurrence in surveys in Oregon (Rowe and Garcia 2013), suggesting some tolerance for certain kinds of habitat modification.

Introduced predatory fish and bullfrogs are widespread throughout *R. aurora* habitat in California, including sites near the coast (T. Fuller, J. Garwood, and M. van Hattem, pers. comm.). Negative impacts have been documented in mesocosm experiments, but field observations have yielded both negative and neutral effects of fish and bullfrogs on *R. aurora* distribution and abundance. Field-enclosure experiments in Oregon have shown reduced survivorship, shifts in microhabitat use, slower development, and smaller size at metamorphosis of *R. aurora* in the presence of fish and bullfrogs (Kiesecker and Blaustein 1998). Surveys in Oregon and Washington have found evidence for negative associations between *R. aurora* presence or abundance and the presence of nonnative fish but weak or no evidence for an effect of bullfrogs (Adams 1999, Pearl et al. 2005a, Rowe and Garcia 2013). Other studies in Oregon and Washington have not detected any effects of fish or bullfrogs on *R. aurora*

presence (Richter and Azous 1995, Adams et al. 1998, Adams et al. 2011). Little data are available from California. Freshwater Lagoon and Big Lagoon in Humboldt County both have a long history of fish stocking, and surveys of suitable habitat during the 2010 and 2011 breeding seasons never found more than 1 egg mass in either lagoon (M. van Hattem, unpublished data). While introduced fish and bullfrogs can prey upon *R. aurora*, the population-level impacts of such predation are unknown. Gut content analysis of 5075 bullfrogs collected over 5 years on Vancouver Island found *R. aurora* in only 0.2% of stomachs (Jancowski and Orchard 2013).

Expected climate changes within the California range of *R. aurora* over the next 100 years include increased temperatures, sea-level rise, changes in hydrology, changes in fire regime, and vegetation shifts (reviewed in PRBO 2011). The frequency of extremely hot days is projected to increase, with roughly nine additional days over 32.2°C (Bell et al. 2004), though the effects of increased temperature are difficult to predict. A mesocosm experiment on larval *R. aurora* found that the combined effects of warming and drying can offset each other: warmer conditions result in more algal resources, allowing larvae to develop faster and escape costs of drying (O'Regan et al. 2014). Sea-level rises as high as 72 cm above 1990 levels are predicted under some models for California (reviewed in PRBO 2011), which may cause saltwater intrusion into estuarine habitat used for breeding. Upwelling is expected to intensify, which may increase fog development and contribute to cooler, moister conditions (Snyder et al. 2003, Lebassi et al. 2009), possibly facilitating terrestrial habitat use by this species along the coast. Potential changes in precipitation are less clear, some models predict either modest increases or decreases in rainfall, while others predict sharp reductions of up to 28%. (reviewed in PRBO 2011). Reductions in water availability due to reduced snowpack and possibly reduced precipitation will affect the timing and magnitude of stream flows, which may negatively affect

habitat (Snyder et al. 2004, Stewart et al. 2005, Cayan et al. 2008b). How fire regime will be affected by climate change in northwestern California is not well understood. Some models predict little change in fire regime or even decreases in area burned along the northern coast (Fried et al. 2004, Lenihan et al. 2008), while increases in area burned have been predicted for the southern coast of northwestern California (Lenihan et al. 2008). Westerling et al. (2011) projected a 100% increase in area burned in northwestern California under some scenarios. How *R. aurora* responds to wildfire is unknown. Vegetation communities are expected to shift from moist conifer to drier mixed evergreen forest, with reductions in Douglas fir and redwood forest in particular (Lenihan et al. 2008, PRBO 2011). Loss of moist forest habitat would likely be detrimental to *R. aurora*; however, most of the predicted vegetation changes occur farther inland from its range.

Disease has been repeatedly implicated in amphibian declines, but to date there is little evidence that disease has played a major role in determining *R. aurora* abundance. While *Bd* has been documented from a high proportion of sites examined in Humboldt County (11/13; Nieto 2004, Sun 2012), the prevalence of infected individuals is relatively low (15%; Nieto 2004, Sun 2012). Water mold infection of egg masses has been observed in the field (Cary 2010, M. van Hattem, unpublished data) but population consequences of infection are unknown. Terrestrial versus aquatic life stages may respond differently to fungal infection. Juvenile metamorphs infected with *Saprolegnia* in the lab did not have significantly higher mortality than uninfected individuals (Romansic et al. 2007), while two weeks of exposure was lethal to *R. aurora* larvae (Romansic et al. 2009a).

Status Determination

Rana aurora has a small range in California in a region that is undergoing continuing development, agricultural use, and timber harvest, making it a Priority 2 Species of Special Concern.

Management Recommendations

Management of *Rana aurora* should focus on addressing habitat degradation and loss due to development, timber harvest, and agriculture (including marijuana cultivation), introduction and spread of nonnative predatory fish and bullfrogs, and on minimizing unintended negative impacts due to salmonid restoration. Observations of higher abundance in breeding habitat with intact terrestrial vegetation nearby (though not excessively shading ponds; Cary 2010, Adams et al. 2011, Holcomb 2012) coupled with the terrestrial habitat use and long distances traveled by adults (Hayes et al. 2007) support the idea of maintaining vegetation buffers around breeding habitat in forested areas and setbacks between wetlands and development. Current regulations for development setbacks under the California Coastal Act of 1976 give distances from breeding wetlands of up to 30 m depending on land use. However, these setbacks are reducible upon request and we recommend that consistent, biologically based setbacks be developed. *Rana aurora* may experience less impact from timber harvesting methods that leave residual tree patches, particularly if multiple trees are included in patches between 0.8 and 1.5 ha in size and are near streams (Chan-McLeod and Moy 2007). Marijuana cultivation appears to pose a growing threat to maintenance of high-quality habitat for this species. Enforcement and regulation of marijuana cultivation is an ongoing issue in California and we suggest that the environmental impact of such activities be considered. Populations of introduced fish and bullfrogs should be prevented from invading *R. aurora* breeding habitat. While bullfrogs may already be widespread, intentional fish stocking should be restricted to avoid *R. aurora* habitat. Restoration projects for native salmonids should also take into consideration potential impacts to *R. aurora* that may be caused by converting freshwater wetlands to estuarine habitats and salt marshes.

Monitoring, Research, and Survey Needs

Monitoring of *Rana aurora* egg mass counts should continue in order to provide baseline data on distribution and abundance and to detect declines. *Rana aurora* management would benefit from additional study of movement and habitat use, life history, effects of marijuana cultivation, and impacts of introduced species in the field. Particularly as habitat becomes increasingly fragmented, data on connectivity among habitat patches, effects of road density, and use of terrestrial habitat away from breeding ponds can help inform appropriate setback distances and buffer configurations. Genetic studies may also be helpful for understanding patterns of frog movement across the landscape. Basic life history information overall and from the California range in particular is also lacking. Understanding saltwater tolerance of different life stages would be useful for predicting the extent of sea-level-rise effects on coastal populations. Field research on impacts of marijuana cultivation on amphibian populations would contribute to developing environmental regulations for this growing industry. Much of the concern for bullfrog impacts on *R. aurora* is from experimental mesocosm studies. Additional research that addresses the effects of bullfrogs and fish on *R. aurora* in the field is necessary to understand the community context of impacts, as the effects of bullfrogs in combination with fish may be greater than either singly (Kiesecker and Blaustein 1998), and fish may be facilitating bullfrog survival (Adams et al. 2003). Under the assumption that eradication of well-established introduced species is unlikely to be feasible at a large scale, a main goal of this work should be identifying factors that can potentially be manipulated to promote coexistence between *R. aurora* and nonnative predators, such as managing terrestrial and aquatic vegetation cover and hydroperiod.



FOOTHILL YELLOW-LEGGED FROG

Rana boylei Baird 1854

Status Summary

Rana boylei is a Priority 1 Species of Special Concern, receiving a Total Score/Total Possible of 83% (91/110). During the previous evaluation, it was also considered a Species of Special Concern, with varying levels of threat in different parts of the range (Jennings and Hayes 1994a).

Identification

Rana boylei is a small to medium-sized frog (up to 81 mm SVL) (Stebbins 2003). The skin usually appears rough and granular, with many tiny tubercles on the surface, including on the tympanum (Nussbaum et al. 1983). The dorsal coloration is variable and can be gray, brown, reddish, or olive, sometimes with extensive brick-red coloration around the weak dorsolateral folds (Nussbaum et al. 1983). Individuals can also change their overall coloration from relatively light to dark (Wheeler et al. 2005). An inverted triangle-shaped patch of buff coloration is usually present on the snout, but its dis-

tinctiveness varies (Stebbins 2003). The ventral coloration is typically yellow on the hind legs and posterior abdomen, with the rest of the venter mostly white with dark mottling on the throat and chest (Nussbaum et al. 1983).

Foothill Yellow-Legged Frog: Risk Factors

Ranking Criteria (Maximum Score)	Score
i. Range size (10)	5
ii. Distribution trend (25)	20
iii. Population concentration/ migration (10)	10
iv. Endemism (10)	7
v. Ecological tolerance (10)	7
vi. Population trend (25)	25
vii. Vulnerability to climate change (10)	10
viii. Projected impacts (10)	7
Total Score	91
Total Possible	110
Total Score/Total Possible	0.83




PHOTO ON PREVIOUS PAGE: Foothill yellow-legged frog, Del Norte County, California. Courtesy of Rob Scheil Photography.

Jennings and Hayes (2005) documented orange or red coloration on the ventral surfaces of the hind limbs in post-metamorphic animals from Glenn, Tehama, and Stanislaus Counties. Tadpoles reach a maximum size of 55 mm and are usually olive dorsally with dark spots or mottling that matches the stream substrate and a silvery venter (Nussbaum et al. 1983). Males call primarily underwater but will also call above (MacTague and Northern 1993).

Other species that *R. boylei* could potentially be confused with in California include the California and northern red-legged frogs (*R. draytonii* and *R. aurora*), the mountain and Sierra Nevada yellow-legged frogs (*R. muscosa* and *R. sierrae*), and juvenile bullfrogs. *Rana draytonii* and *R. aurora* have smooth skin, a prominent jaw stripe, distinct dorsolateral folds, and usually have red coloration under the hind limbs (although *R. boylei* can also have red ventral coloration, and young *R. draytonii* and *R. aurora* often have yellowish thighs) (Stebbins 2003, Jennings and Hayes 2005). *Rana muscosa* and *R. sierrae* have smoother skin, smooth tympana, and tend to lack the light patch on the snout (Stebbins 2003). Bullfrogs occasionally co-occur with *R. boylei* but tend to be greenish in color, with smoother skin, and large, smooth tympana (Stebbins 2003).

Taxonomic Relationships

Rana boylei has been recognized as a distinct species for a long period of time, although its phylogenetic placement among other North American ranids has been revised repeatedly (Baird 1854, Macey et al. 2001, Hillis and Wilcox 2005). Zweifel (1955) documented variation in color and morphology among California *R. boylei* populations. Recent phylogeographic studies have found that genetic variation among *R. boylei* populations is structured along hydrologic boundaries (Dever 2007, Peek 2010, Lind et al. 2011). In a range-wide phylogeographic study, Lind et al. (2011) identified some peripheral populations that are deeply divergent from populations within the core of the range. In California, populations in southern-

most Monterey County west and south of the Salinas River Valley and populations from the southern Sierra Nevada were found to be phylogenetically distinct from the rest of *R. boylei*, suggesting a long history of isolation. While extreme southern populations from Los Angeles County are now extirpated, Lind et al. (2011) hypothesized that animals from those localities may also have been genetically distinct.

Life History

As a stream-dwelling frog, the life history of *Rana boylei* coincides with seasonal patterns in river flows associated with California's Mediterranean climate. The most sensitive life stages (eggs and larvae) develop during relatively stable conditions when streams are at their lower stages (Kupferberg et al. 2009b). Breeding and oviposition occur in spring after flood waters recede, and tadpoles metamorphose in late summer through early autumn before winter rains (reviewed in Lind 2005, Haggarty 2006, Wheeler and Welsh 2008). Southern populations breed earlier than northern populations (Zweifel 1955), and the onset and duration of breeding can be influenced by water temperature, cessation of rainfall, water velocity and depth, and day length (Zweifel 1955, Kupferberg 1996a, Lind et al. 1996). Between 2002 and 2007 at a site in Del Norte County, breeding activity was initiated in early April and lasted for 19–52 days, with earlier breeding occurring in low-flow years (Wheeler and Welsh 2008). Breeding activity ceased briefly during rain events that increased flows (Wheeler and Welsh 2008).

Females lay a single cluster of up to 2000 eggs (Zweifel 1955) attached to pebble or cobble substrates (Fuller and Lind 1992) or to bedrock (M. van Hattem, pers. comm.). Eggs take 2–3 weeks to hatch, depending primarily on water temperature (Kupferberg 1996a). Major sources of natural egg mortality are desiccation through stranding in dry years, and scour from floods in wet years (Kupferberg et al. 2009b). Adults breed at 2 or 3 years of age depending on the geographic location, and this translates into

fluctuations in adult populations being determined by environmental conditions during recruitment 2–3 years prior (Kupferberg et al. 2009b). Metamorphosed animals captured in Tehama County were 1.2–7.2 years old based on skeletochronology (Bourque 2008), suggesting that they can be relatively long-lived.

Radiotelemetry studies are beginning to offer more insight into terrestrial movements. Adults aggregate at pools in the spring but become more difficult to find in the summer (Van Wagner 1996, Haggarty 2006, Wheeler and Welsh 2008). In one study in Tehama County, frogs used watercourses for movement and were rarely more than 12 m from the stream channel (Bourque 2008). Females tended to move upstream during spring and downstream during the fall and winter. Travel rates in this population were up to 1386 m/day, faster than previously thought. In other studies, the longest distances traveled have been closer to 500 m at rates of tens to a few hundred meters per day (Van Wagner 1996, Drennan et al. 2006, Wheeler et al. 2006). Females tend to move farther distances than males, with female movements up to 7 km documented in one study (Bourque 2008, Gonsolin 2010). At one locality in Del Norte County, 68% of males remained in one breeding site during the reproductive season, with average home range sizes of 0.58 m² (Wheeler and Welsh 2008). At a site where the availability of permanent water is a limiting factor in Santa Clara County, resident tributary frogs moved to the main stem to breed and moved greater distances than resident main stem frogs (Gonsolin 2010). Greater than 90% of movements were associated with movements to or from breeding sites, and all movements outside of the breeding season were made in response to the channel drying back or to rainfall (Gonsolin 2010).

Larvae appear to be herbivorous, while metamorphs and adults consume terrestrial and aquatic insects. Algae with epiphytic diatoms are a preferred food for larvae, and the abundance of floating algae indicates the quality of larval food resources (Kupferberg 1996b, Kup-

ferberg 1997). Metamorphosed animals primarily forage terrestrially (Zeiner et al. 1988, Van Wagner 1996, Haggarty 2006, Hothem et al. 2009). Spiders, beetles, and flies are common prey items (Haggarty 2006, Wiseman and Bettaso 2007, Hothem et al. 2009). Gut content analyses of adults collected from 22 sites in the Cache Creek watershed found that 98% of individuals contained terrestrial prey, 28% contained aquatic prey, and one animal contained mammal hair and bone fragments (Hothem et al. 2009). Two occurrences of adults cannibalizing juvenile conspecifics have been documented (Wiseman and Bettaso 2007).

Habitat Requirements

Rana boylei is primarily stream dwelling and requires shallow, flowing water in streams and rivers with at least some cobble-sized substrate (Hayes and Jennings 1988). Different life stages use different habitat types for development, foraging, and overwintering.

Breeding and oviposition occur at the margins of relatively wide and shallow channel sections, habitats that experience reduced flow variation (Storer 1925, Fitch 1936, Kupferberg 1996b, Lind et al. 1996). Breeding sites are often located near tributary confluences (Kupferberg 1996a, Bourque 2008). Egg masses are attached in low-flow locations behind and sometimes under rocks. The most commonly used substrates for breeding sites are cobble, boulders, and gravel (Fuller and Lind 1992, Kupferberg 1996a). Eggs have been found at water depths up to 87 cm (C. Bondi, S. Yarnell, and A. Lind, pers. comm.), in water velocities of 0–0.21 m/s, and up to 12.5 m from shore (Kupferberg 1996a, reviewed in Lind 2005). The critical thermal maximum for embryos is 26°C, and eggs have been found in water ranging from 9°C to 21.5°C (Zweifel 1955). Density of egg masses was highest in Eel River reaches when July mean temperatures were between 17.5°C and 19°C (Catenazzi and Kupferberg 2013). Egg mass surveys from 1991 to 2002 across 11 small and large streams in the Northern Coast Ranges and the Sierra Nevada found

that oviposition sites occurred in a very narrow range of microhabitat conditions that were different from randomly selected habitats, strongly suggesting active habitat selection by frogs (Lind 2005). High-quality breeding areas are often used over multiple years (Lind 2005). Larvae tend to stay in natal habitats until they metamorphose (Van Wagner 1996). Surveys in the Mattole Watershed in northern coastal California across different channel types found that tadpole presence was best predicted by relatively warmer water temperatures (Welsh and Hodgson 2011). Tadpoles were never found in water colder than 13°C, and tadpole abundance increased with water temperature (Welsh and Hodgson 2011). In choice experiments, tadpoles selected temperatures between 16.5°C and 22.2°C (Catenazzi and Kupferberg 2013).

Metamorphosed animals use a variety of aquatic habitats, including riffles, pools, and glides (reaches intermediate between riffles and pools) depending on the life stage and season (Van Wagner 1996, Yarnell 2000, Lind 2005, Yarnell 2005, Haggarty 2006). At Red Creek in Tehama County, post-breeding season adults and subadults preferred pool and riffle habitats, while young of the year metamorphs selected slower-moving glides and runs (Haggarty 2006). In Nevada County, all age classes used riffles after the breeding season (Van Wagner 1996). In the Sierra Nevada foothills, subadults chose fast-flowing sections of stream, while adults used slower-moving pool habitats (Yarnell 2000, Yarnell 2005). In the Mattole Watershed, the best predictor of adult presence in streams was canopy openness (Welsh and Hodgson 2011). Abundance of adults and larvae was positively associated with larger basin areas and finer substrates, conditions more typical of alluvial channels than other channel types (Welsh and Hodgson 2011).

Less is known about terrestrial habitat use. Adults typically occur along waterways with some degree of shading (Fitch 1938, Zweifel 1955, Moyle 1973, Hayes and Jennings 1988, Van Wagner 1996), although they also occur in open habitats (Welsh et al. 2005, Haggarty

2006, Welsh and Hodgson 2011). During the spring, radio-tracked males and females in Tehama County were often found on land near water (38% and 66% of the time, respectively; Bourque 2008). The average distance from water was less than 3 m in all seasons, although adults occasionally used habitat up to 40 m distant from streams (Bourque 2008). Adults move to tributaries or upland habitats to avoid floods following large rain events (Kupferberg 1996b, Van Wagner 1996, Yarnell 2000, Bourque 2008). Tributaries are also used for overwintering in early spring before adults are abundant on the principal channels (Kupferberg 1996b, Yarnell 2000). Juveniles will also move into tributaries, with maximum movements of 860 m from hatching site to upstream tributaries observed in Santa Clara County (Gonsolin 2010). Adults may aggregate above ground in terrestrial microhabitats on tributaries post-breeding (Leidy et al. 2009).

Distribution (Past and Present)

Historically, *Rana boylei* occurred in foothill and mountain streams from the San Gabriel River in Los Angeles County to southern Oregon west of the Sierra-Cascade crest (Nussbaum et al. 1983, Stebbins 2003), from sea level to 1940 m (Hemphill 1952). There is an isolated, unverified record from northern Baja California, Mexico, at ~2000 m (Loomis 1965).

Jennings and Hayes (1994a) considered *R. boylei* endangered in central and southern California south of the Salinas River, threatened in the west slope drainages of the Sierra Nevada and Cascades, and of special concern in the Coast Ranges north of the Salinas River. They estimated that *R. boylei* were extirpated from 45% of their historical localities in California, and 66% of historical localities from the Sierra Nevada. Building on that mapping effort, Lind (2005) looked at 394 historic localities in California and Oregon, and found that 201 localities (51%) were no longer occupied, with extirpations largely in southern California and northern Oregon. Kupferberg et al. (2012)

determined current occupancy of 310 randomly selected sites that were occupied prior to 1975. They found that half of the sites still had *R. boylei* populations, with frogs more likely to be present in sites without large dams.

Extirpations likely began in the second half of the twentieth century. Grinnell and Storer (1924) noted several sites in the Sierra Nevada foothills around Yosemite where *R. boylei* were common. In resurveys of those sites and surveys of additional sites in the early 1990s, Drost and Fellers (1996) did not find any *R. boylei*. Surveys by Moyle (1973) in the 1970s found *R. boylei* at only 30/95 sites in the southern and central Sierra Nevada foothills. Field surveys since 1993 have found at least one frog at only 213/804 sites in 28/40 California counties (Fellers 2005a). Fellers (2005a) estimated that extant populations occur in 40% of streams in the Pacific Northwest, 30% of streams in the Cascade Mountains, 30% of streams in the south Coast Range (south of San Francisco), and 12% of streams in the Sierra Nevada.

Trends in Abundance

Kupferberg et al. (2012) compiled egg mass density data from multiple sources on 27 Sierran and coastal populations in northern California between 1991 and 2010. The range of densities reported was between 1.9 and 105.7 clutches/km of reach sampled. Average density was higher in free-flowing rivers (31.1 clutches/km) than in rivers with dams (5.5 clutches/km), but no differences were detected between abundances in coastal versus montane watersheds (Kupferberg et al. 2012). Fellers (2005a) reported that only 30 of 213 occupied California sites had population sizes greater than 20 adults. In the Coast Ranges, population sizes of greater than 100 adult frogs occurred at six sites, and populations greater than 50 adult frogs occurred at nine sites (Fellers 2005a). Small population sizes are presumably due to population declines, leading to predictions that populations in the southern Sierra Nevada will not be viable for more than another decade (Fellers 2005a). Minimum viable population sizes

are unknown, however, and may vary across the range.

Nature and Degree of Threat

The main threats to and likely causes of *Rana boylei* decline are human activities that alter natural hydrologic regimes of streams and rivers, such as dams for hydroelectric power generation, water storage, and water delivery. Other potential stressors include land use changes that degrade or destroy riparian habitat (particularly urban and agricultural development), pesticides, disease, and invasive species.

Alterations to the natural flow regime, for example, through dam releases, can have direct mortality effects and indirect negative effects on *R. boylei* by altering habitat availability and quality. Kupferberg et al. (2009b) reviewed published literature and Federal Energy Regulatory Commission hydroelectric dam relicensing reports to assess the effects of pulsed flow releases on *R. boylei*. The data spanned 1997–2007 and included seven major river basins in California. Pulsed flows from dam releases after oviposition resulted in scouring of egg masses, while flow changes during oviposition led to stranding when water levels subsequently dropped and exposed egg masses. Similarly, tadpoles can be scoured and stranded due to pulsed-flow releases. The effect of releases on post-metamorphic animals is less clear, and the impact of flow changes on habitat availability is highly site specific. Reservoirs and dams may also disrupt patterns of connectivity among *R. boylei* populations. Comparisons of genetic structure within and among *R. boylei* populations in three pairs of regulated versus unregulated Sierran rivers found that regulated rivers exhibited lower genetic diversity and greater genetic drift compared to unregulated river populations (Peek 2010).

Kupferberg et al. (2009c) modeled *R. boylei* population growth under different flow scenarios. A major result was that populations in regulated rivers had 4–13-fold greater extinction risk than populations in unregulated rivers due to smaller population sizes. Kupferberg

et al. (2009c) simulated how an unregulated population would be affected by flows more typical of regulated rivers. When subjected to aseasonal flow conditions, modeled populations showed a doubling of extinction risk. Many different kinds of hydrologic changes can contribute to these negative effects, and when different stressors are combined, the impact on frog populations is greater than expected from simply adding up the effects of individual stressors.

Field and laboratory experiments conducted by Kupferberg et al. (2011) showed that tadpoles suffered negative effects including death at or below water velocities experienced during aseasonal pulsed flows. For example, most tadpoles could no longer swim or seek refuge at velocities of ≥ 20 cm/s or greater, and in the absence of refugia tadpoles reached exhaustion in ~ 7 min in a 5 cm/s current. Rates of flow in regulated reaches can be much higher than this. For example, in the North Fork Feather River, surface velocity measured in larval rearing habitat near channel edges can reach over 30 cm/s after releases for recreational purposes (Garcia and Associates 2005).

Smaller-scale hydrologic modification and loss or degradation of riparian habitat due to urban and agricultural use is also a threat to *R. boylei*. Analyses correlating *R. boylei* distribution with landscape characteristics demonstrated negative effects of urban and agricultural land use change and pesticides on *R. boylei* presence (Davidson et al. 2002, Davidson 2004, Lind 2005). Vineyard conversion can have impacts on small creeks, and the establishment of permanent ponds used for irrigation and frost protection can create habitat for bullfrogs (S. Kupferberg, pers. comm.). Marijuana cultivation practices that divert water from small creeks can lead to premature drying. Growers have been observed to construct plastic-lined impoundments in creeks and add fertilizers directly to creek water, as well as use pesticides and herbicides in and around frog habitat (Gonsolin 2010). These practices are suspected to have contributed to declines in

some populations near Gilroy (Gonsolin 2010). Similar impacts are likely in Humboldt, Mendocino, and Trinity Counties (CDFG 2013). The large-scale effects of such illegal operations are unknown, and potentially dangerous to study. While in-stream gravel and suction dredge gold mining may have been more of a concern in the past, current regulations protecting salmonids have likely largely reduced the direct impact of such activities on *R. boylei*. For example, in Humboldt County in-stream gravel mining occurs above (in elevation) and outside the wetted channel, and relatively high egg mass density has been documented in reaches where gravel mining occurs in the Mad River (M. van Hatten, pers. comm.).

The current distribution of *R. boylei* is strongly correlated with climate variables, which suggests that this species may be sensitive to future climate changes, particularly those that affect stream hydrology (reviewed in PRBO 2011). Comparisons of occupied and extirpated historic localities found that sites where *R. boylei* persists have higher mean annual precipitation, less variability in precipitation, and fewer dry years than extirpated sites (Davidson et al. 2002, Lind 2005). Within the range of *R. boylei*, warming temperatures are predicted to result in more precipitation falling as rain instead of snow, and consequently less storage of water as snowpack. Reductions of 30–80% in snowpack accumulation are predicted within the northwestern range of *R. boylei*, and up to 90% reduction in snowpack is predicted for the south coast hydrologic region (Snyder et al. 2004, Cayan et al. 2008b). In the Sierra Nevada, snowpack losses of 50–90% are predicted by the end of the twenty-first century, with greatest losses at low to mid-elevations (Knowles and Cayan 2002, Hayhoe et al. 2004, Knowles and Cayan 2004, Maurer 2007, Cayan et al. 2008b). Loss of snowpack is likely to result in earlier runoff and reduced spring and summer streamflows. Timing of spring snowmelt is predicted to shift earlier in the spring in the Sierra Nevada (Snyder and Sloan 2005), while in northwestern California the opposite

has occurred over the last 50 years (Stewart et al. 2005). How frogs will respond to these changes in hydrology is unknown, but negative effects due to anthropogenic changes in hydrology are well documented. Reduction in water availability may also lead to more conflict with human use of water and affect how regulated reaches are managed (reviewed in Franco et al. 2011). It is important to note, however, that predictions of changes in precipitation are much less certain than predictions for temperature (Franco et al. 2011, PRBO 2011). In addition, climate change may also affect disease dynamics. Outbreaks of nonnative parasitic copepods occurred during two recent warm years at a long-term study site, resulting in morphological abnormalities and smaller sizes at metamorphosis (Kupferberg et al. 2009a). The outbreak was likely caused by increased summer water temperature, decreased daily discharge, or a combination of these factors. These conditions may increase under a changing climate, but could also occur as a result of marijuana cultivation.

No declines to date have been associated with *Bd*, but the disease does infect *R. boylei* in the field. Padgett-Flohr and Hopkins (2009) examined museum specimens from 1890 to 2000, and found that *Bd* first appeared in *R. boylei* samples from the 1960s, with 10% of specimens infected. In all, 0–40% of specimens were infected with *Bd* in the following decades. In laboratory trials, *R. boylei* appeared to be protected by skin peptides against *Bd* and therefore may not be very susceptible to chytridiomycosis (Davidson et al. 2007). Chytrid infection did not affect survival, even in the presence of a co-applied pesticide, but did suppress growth of recently metamorphosed individuals by approximately 40% (Davidson et al. 2007).

Observational data and surveys have found that *R. boylei* is rare or absent in habitats with introduced fishes and bullfrogs (Hayes and Jennings 1986, Hayes and Jennings 1988, Kupferberg 1997, Lind et al. 2003, Fuller 2008). Breeding populations of *R. boylei* can be an

order of magnitude smaller when bullfrogs are present compared to uninvaded reaches (Kupferberg 1997). In field experiments in outdoor enclosures, bullfrog tadpoles caused a 48% reduction in survivorship of *R. boylei* tadpoles, and a 24% decline in mass at metamorphosis. The mechanism behind the negative impacts of bullfrogs was competition for food (Kupferberg 1997). Metamorphosed bullfrogs prey on *R. boylei*, including post-metamorphic individuals (Crayon 1988, Hothem et al. 2009), but the population-level consequences of this predation are unclear. Another nonnative predator, the signal crayfish (*Pacifastacus leniusculus*), has been introduced into several Sierra Nevada drainages from farther north where the two species co-occur (Wiseman et al. 2005). Signal crayfish have been observed eating and dislodging egg masses and attacking larvae (Wiseman et al. 2005). Within *R. boylei*'s range, signal crayfish have been documented at 30–40 sites, with ~25 invaded sites occurring in the Sierran foothills (G. Fellers, pers. comm.).

Status Determination

Documented declines and extirpations of *Rana boylei* populations combined with continuing threats to remaining populations result in a Priority 1 designation for this species.

Management Recommendations

Several aspects of the biology of *Rana boylei* can help inform management efforts. *Rana boylei* use a variety of stream and streamside habitats during different life stages; therefore, protected habitat needs to provide adequate habitat diversity. The timing and pattern of releases of water from dams during April through June should be managed to minimize egg scouring and stranding. For example, dam releases can be staggered to better mimic the natural spring recession in snowmelt-fed streams. Further recommendations for hydrologic management can be found in Kupferberg et al. (2009b, 2009c). Dam removal should be explored where appropriate and is likely to benefit *R. boylei* and other native taxa. River management for other taxa

needs to take *R. boylei* into account. For example, in-stream structures to improve habitat for fish such as steelhead can negatively impact *R. boylei* (Fuller and Lind 1992). Habitat restoration and possibly repatriation of southern Sierra Nevada populations should be considered. Southern populations in general should be priorities for conservation because of the degree of losses and distinctive genetic diversity represented in this part of the range (Lind et al. 2011). Removal or management of nonnative predators such as fish and bullfrogs may help restore *R. boylei* habitat. For example, projects that remove artificial pools (e.g., relict mine tailing ponds) by restoring linkages to main river channels would result in more natural hydrologic conditions and reduce breeding habitat for bullfrogs (Fuller et al. 2010). Finally, Lind et al. (2011) suggested that an approach using genetic analyses of *R. boylei* and co-distributed riverine taxa would help in prioritizing drainages for protection based on levels of diversity.

Monitoring, Research, and Survey Needs

Modeling of population dynamics and hydrology are highly site specific and limited by available demographic data, and acquiring those additional data should be a high priority for *Rana boylei*. More research is needed on survivorship of tadpoles and juveniles, especially during overwintering. The mechanisms underlying hydrological effects are currently best understood for egg masses, and we need to develop a better functional understanding of how hydrology affects different life stages (Kup-

ferberg et al. 2009b). More research is also needed on post-metamorphic stages. Post-metamorphic stages may be less at risk from aseasonal pulses in river flow because they are more mobile, but in regulated rivers the timing of pulsed flow events can be decoupled from climatic cues (such as the first appreciable fall rains) that would normally trigger movement to safer refuges (Kupferberg et al. 2009b). Caution should be taken in using radio telemetry to study post-metamorphic animals, as 62% of frogs in one study suffered skin injuries from transmitters (Bourque 2008). Modeling efforts would also be improved by monitoring a Sierra Nevada population in an unregulated reach for comparison with more regulated sites (Kupferberg et al. 2009b). Egg mass counts are commonly used to monitor *R. boylei* populations. Females only lay one mass/year, so egg mass counts accurately reflect the number of reproductive females. However, operational sex ratios are female biased; therefore, accurate population size estimates cannot be made based on egg counts alone (Wheeler and Welsh 2008). Lind et al. (2011) provided important range-wide phylogeographic data, but their study was limited by very low nuclear genetic diversity and relied primarily on mitochondrial data. Additional work could provide valuable additional data on levels of variation and genetic isolation among local hydrologic basins, as might be predicted for this stream-restricted anuran. Finally, efforts to find remnant *R. boylei* populations in the San Gabriel Mountains and upper Piru Creek in southern California should continue.



CASCADES FROG

Rana cascadae Slater 1939

Status Summary

Rana cascadae is a Priority 2 Species of Special Concern, receiving a Total Score/Total Possible of 65% (72/110). During the previous evaluation, it was also considered a Species of Special Concern (Jennings and Hayes 1994a).

Identification

Rana cascadae is a medium-sized (2.0–8.0 cm SVL) frog with drab-green, tan, or brown dorsal coloration and well-defined black blotches scattered across the back (Slater 1939, Stebbins 2003). The number of blotches varies from very few to about 50 (Slater 1939), and unmarked individuals occur rarely (Jennings and Hayes 1994a). Blotches appear to be on the surface of the frog's skin and are reminiscent of spattered ink (Stebbins 2003). The species has a prominent light stripe above the jaw and strong dorsolateral folds. The venter is cream or buff, usually with yellowish (sometimes reddish) areas posteriorly and on the undersides of the legs. Laterally, the sides are mottled and

fade into the ventral coloration (Slater 1939). The male advertisement call is a series of low chucks given in rapid succession, usually ending with one slightly drawn out chuck (Elliott et al. 2009).

Cascades Frog: Risk Factors

Ranking Criteria (Maximum Score)	Score
i. Range size (10)	10
ii. Distribution trend (25)	20
iii. Population concentration/ migration (10)	10
iv. Endemism (10)	0
v. Ecological tolerance (10)	3
vi. Population trend (25)	15
vii. Vulnerability to climate change (10)	7
viii. Projected impacts (10)	7
Total Score	72
Total Possible	110
Total Score/Total Possible	0.65




PHOTO ON PREVIOUS PAGE: Cascades frog, Trinity County, California. Courtesy of Adam Clause.

In California, this species could be confused with the California or northern red-legged frogs (*R. draytonii* and *R. aurora*), both of which it resembles in overall body shape. In adults, *R. aurora*/*R. draytonii* have extensive mottling on the venter with red pigmentation on the ventral thighs and groin, rather than the yellow that often characterizes *R. cascadae* (Dunlap 1955). However, the color of the thighs is variable in *R. cascadae* and may not be a reliable character to separate these taxa (S. Barry, pers. comm.).

Taxonomic Relationships

This species is closely related to *Rana aurora* and *R. draytonii* (Shaffer et al. 2004, Hillis and Wilcox 2005). It was proposed as a distinct species based on morphology (Slater 1939), and this interpretation has been repeatedly confirmed with additional morphological and genetic data (Dunlap 1955, Case 1978, Shaffer et al. 2004).

Based on genetic data, the species appears to show considerable differentiation among local populations that is consistent with an overall isolation-by-distance model of gene flow (Monsen and Blouin 2003, Monsen and Blouin 2004). In addition, the California populations appear to be strongly divergent in both mitochondrial and nuclear DNA from the remainder of the species' range in the Cascade and Olympic mountain ranges (Monsen and Blouin 2003). Populations of this species appear to have consistently small effective population sizes (<50; Phillipsen et al. 2011). The available data are primarily from outside of California, although the observed pattern is likely consistent throughout the range. Preliminary results based on mitochondrial DNA, nuclear DNA, and microsatellites suggested little divergence between Lassen and Klamath populations in California (Chang and Shaffer 2010). However, more extensive work with larger range-wide sampling is needed.

Life History

Rana cascadae breeds in the spring, soon after emerging from hibernation and the spring thaw that opens breeding pools (Nussbaum et al.

1983, Stebbins 2003). First-time breeders frequently disperse to new areas of suitable breeding habitat (51% of first-time breeders relative to only 7% of experienced breeders in Echo Lake Basin; Garwood 2009), which may help to connect local subpopulations into larger more stable metapopulations. Breeding occurs at the margins of waterbodies, with oviposition often occurring in large aggregations (Syte 1975, Nussbaum et al. 1983, Garwood 2009). Oviposition behavior appears to be variable throughout the species' range, with some authors reporting diurnal oviposition of largely unattached egg masses (Briggs 1987), and others noting that most egg masses are deposited at night and are attached to vegetation (Nussbaum et al. 1983; K. Pope, pers. comm.). Breeding at individual sites is relatively synchronous and occurs over a few days, although the timing of breeding across the range can vary widely with local weather conditions and elevation (Briggs 1987, Garwood 2009). Embryo development can occur at temperatures ranging from 6°C to 27°C (Syte 1975, Nussbaum et al. 1983). After hatching, larvae sometimes aggregate into dense clusters (generally fewer than 40 individuals) composed primarily of siblings (O'Hara and Blaustein 1981, O'Hara and Blaustein 1985, Blaustein and O'Hara 1987) and choose higher water temperature than those required during embryo development (up to ~28°C; Wollmuth et al. 1987, Bancroft et al. 2008). After metamorphosis, lower water temperatures are again preferred.

Rana cascadae appears to be largely diurnal. The diet of adult frogs is generalized and includes a wide variety of arthropods, as is the case for most other California ranids (Joseph et al. 2011). An analysis of stomach contents for 275 frogs documented the presence of 110 invertebrate taxa (Larson 2012). Frogs across all size classes generally avoided small prey items (<4 mm), and larger frogs more strongly preferred large prey items (Larson 2012).

Habitat Requirements

Rana cascadae utilizes a wide variety of aquatic habitats, including temporary and permanent

ponds, lakes, marshes, and streams, as well as adjacent vegetated terrestrial habitat (Nussbaum et al. 1983, Jennings and Hayes 1994a, Stebbins 2003, Pearl and Adams 2005, Garwood 2009). The species will also use wet meadows (often those that have formed from old sphagnum bogs) and can occasionally be found a large distance from water (Nussbaum et al. 1983). They require water year-round at all life stages and cannot tolerate habitats that freeze solid in the winter (K. Pope, pers. comm.). Montane lentic habitat is required for breeding and overwintering, with small, shallow, spring-fed ponds serving as the primary breeding habitat (Garwood 2009). Populations appear to be sustained by a matrix of varying habitat types that individual frogs disperse among throughout the year (Garwood and Welsh 2007, Garwood 2009), suggesting that habitat conservation needs to consider spatial scales larger than single lakes (or other patches of habitat). The presence of predaceous fish may limit their distribution (Welsh et al. 2006, Pope et al. 2008), although this alone cannot explain the broadscale pattern of population declines in this species (Fellers et al. 2008; also see the “Nature and Degree of Threat” section below).

Distribution (Past and Present)

In California, *Rana cascadae* occurs in two population segments. One is in the Lassen area and the extreme northern end of the Sierra Nevada (Lassen, Plumas, Shasta, and Tehama Counties) and is now nearly extirpated. The other occurs in the Trinity Alps and Siskiyou Mountains region. The species’ range in California extends from Siskiyou County south to the northern end of Butte County. Outside of California, the range of *R. cascadae* follows the Cascade Range nearly to the United States–Canadian border, with another disjunct population at high elevations on the Olympic Peninsula (Stebbins 2003).

Trends in Abundance

Populations of this frog have declined strongly in the Lassen area, where nearly all known

populations have disappeared in the last 30 years (Fellers and Drost 1993, Jennings and Hayes 1994a). More recent surveys in the Lassen region further confirm these declines. *Rana cascadae* was found at only 6 of 856 sites surveyed over 14 years, population sizes were small, and breeding was limited at these 6 sites (Feilers et al. 2008). Populations elsewhere, including the Klamath Mountains region in Siskiyou and Trinity Counties, are also fragmented, generally small, and at risk, although they are more intact overall than in the Lassen area (K. Pope, pers. comm.). Localized declines have also been detected elsewhere in the range (Pearl and Adams 2005, Fellers et al. 2008, Piovia-Scott et al. 2011). Welsh et al. (2006) found *R. cascadae* to be the most common anuran in the Klamath region. By contrast, more recent and ongoing surveys of eight populations in the Trinity Alps within the Klamath region find that only one of the populations is large and robust and that some of the threats present in the Lassen region are likely also operating there (K. Pope, pers. comm.). Pope and Larson (2013) report 11 remaining populations in the Lassen area and find that the number of young frogs was low at all sites that they surveyed.

Nature and Degree of Threat

Threats to this species appear to be complex and derived from multiple stressors. The largest factor contributing to declines in the Lassen region appears to be overall low recruitment due to changing hydrological conditions that lead to detrimentally high water temperatures and desiccation of egg masses and tadpoles, as well as impacts from *Bd* among subadult frogs (Pope et al. 2011). Extensive mark-recapture surveys in the Lassen region between 2008 and 2010 indicate widespread desiccation of egg masses and tadpoles and a lack of metamorphs relative to more stable populations in the Klamath area (Pope et al. 2011, Pope and Larson 2013). In comparisons between two of the remaining Lassen populations, the population with higher *Bd* prevalence and load in adult and

subadult frogs had lower survivorship for these two age classes. In both Lassen and the Klamath Ranges, subadult frogs had higher *Bd* prevalence and load than adult frogs, and the prevalence of *Bd* increased throughout the active season for subadult frogs but not for adult frogs. These results are consistent with previous studies of *Bd* in this species that suggest the pathogen has differential impacts depending on age class. Blaustein et al. (2005) examined the effect of *Bd* on larvae and found an increased incidence of mouthpart abnormalities but no effect on mortality or behavior. Garcia et al. (2006), however, found significant mortality in new metamorphs of *Rana cascadae* due to *Bd*.

Interestingly, *Bd* also appears to be widespread in the Klamath region where this species is currently much more stable than in the Lassen region (Piovia-Scott et al. 2011), suggesting more than one factor is playing a role in the declines. Ongoing characterization of *Bd* prevalence in these populations could help determine what factors are involved, although one hypothesis is that *Bd* achieves higher loads on frogs in declining populations than stable populations (J. Piovia-Scott, pers. comm.). Infection by the water mold *Saprolegnia* has also been implicated in *R. cascadae* declines. This pathogen is known to increase mortality in embryos, larvae, and metamorphs (Kiesecker and Blaustein 1999, Romansic et al. 2009a) and may have strong impacts on the outcome of competition between *R. cascadae* and sympatric *Pseudacris regilla* (Kiesecker and Blaustein 1999).

Habitat loss and modification is also a threat to continued persistence of populations in both the Klamath and Lassen regions. The species is highly associated with meadows, which have been impacted by cattle grazing, tree encroachment due to lack of wildfire, and changing hydrology associated with changes in the snowpack (K. Pope, pers. comm.; Pope et al. 2014).

Other possible contributors to *R. cascadae* declines that have been proposed include introduced fishes, environmental contaminants,

pathogens, and UV-B radiation. The presence of introduced trout appears to be inversely related to the distribution of *R. cascadae* (Welsh et al. 2006) and almost certainly impacts some populations. Aside from direct predation, introduced trout may affect *R. cascadae* indirectly by supporting higher populations of the aquatic garter snake (*T. atratus*), a predator on both trout and *R. cascadae* (Garwood and Welsh 2007, Pope et al. 2008), and by preemptive competition for aquatic prey (Joseph et al. 2011). However, trout have been present in the Lassen region for nearly a century and are also widely distributed in other areas where *R. cascadae* persists, making it unlikely that they alone can explain the declines over the last 30 years (Fellers et al. 2008). Nevertheless, populations appear to respond favorably to trout removal, showing marked increases in population size and recruitment following fish removal (Pope 2008).

Pesticide use is inversely correlated with the presence of *R. cascadae* (Davidson 2004). In particular, downwind transport of pesticides from intensively farmed areas in the Central Valley appears to be correlated with declines in several species of ranid frogs, including *R. cascadae* (Davidson et al. 2002, Davidson 2004). This hypothesis is attractive in that it explains the differential declines between the Trinity Alps region and the Lassen region because the Lassen region is directly downwind of areas that experience heavy agricultural use, whereas the Trinity Alps are not (Davidson et al. 2002). However, recent field measurements of contaminant residues in sediment and in *R. cascadae* and *P. regilla* tissue do not indicate higher levels in the Lassen compared to the Trinity Alps region, at least for the handful of different chemicals that have been analyzed to date, calling this hypothesis into question (Davidson et al. 2012). In addition, Sparling et al. (2001) measured the presence of cholinesterase levels in the non-declining *P. regilla* as a measure of the extent of pesticides that are locally deposited in an area and found strong effects in the Sierra Nevada but not in the Lassen area; these

results seem to indicate that pesticides may not be a major factor in the Lassen *R. cascadae* declines. Environmental contaminants at sub-lethal levels have also been shown to induce behavioral and morphological changes in *R. cascadae* (Marco and Blaustein 1999), suggesting that low-level agricultural residues may have important biological consequences. In summary, it appears that pesticides may be playing some role in *R. cascadae* declines in the Lassen region, but they are certainly not the entire story.

Finally, UV-B radiation may play a role, possibly in combination with other factors, in causing declines. Some studies have documented larval mortality and retinal damage due to UV-B, although the effect depends strongly on the intensity of UV-B, the duration of exposure, and possibly other factors including the presence of competitors, predators, or supplementary food (Fite et al. 1998, Hatch and Blaustein 2000, Belden et al. 2003, Garcia et al. 2006, Romansic et al. 2009b). The importance of these results has not yet been demonstrated in natural settings, however. Palen et al. (2002) found that dissolved organic matter in natural environments provided protection from UV-B at 89% of the sites examined for *R. cascadae*. Thus, it remains possible that UV-B is having an effect, although its importance in nature remains unclear.

Ultimately, it is likely that no one factor is solely responsible for the precipitous declines in Lassen region *R. cascadae* populations. Further, the causes of the initial range-wide declines may be distinct from the local factors that threaten the continued persistence of the few remaining populations. The most recent work suggests that the major factors playing a role in the range-wide declines are the presence of introduced fishes and *Bd*, while continued local persistence of the remaining populations is also threatened by low recruitment stemming from desiccation and detrimentally high water temperatures. Pope et al. (2014) present a recent and comprehensive review of both regional and local-scale threats to *R. cascadae*

throughout the range. The evidence that synergistic effects occur between several alternative mechanisms of decline is now widespread for a variety of amphibian species (Fellers et al. 2008). As declines have occurred, whatever the cause, it is likely that a breakdown of metapopulation dynamics will contribute to further declines as existing populations become more and more fragmented, decreasing the opportunity for population rescue via recolonization.

Status Determination

The catastrophic declines in the Lassen area are the primary reason for the SSC designation. *Rana cascadae* is nearly extirpated in the Lassen region, is undergoing local population declines elsewhere in its range, and appears to be susceptible to a wide range of threats. However, this frog is a moderate ecological specialist that appears to be relatively stable through much of its range, including a significant fraction of its range in California. The factors that caused declines in the Lassen area appear to not have operated in the Klamath area to date, leading us to project moderate future impacts on extant populations and a Priority 2 status. If strong declines begin to occur in the Klamath area, then a higher priority status will rapidly become justifiable.

Management Recommendations

Fellers et al. (2008), Pope et al. (2011, 2014), and Pope and Larson (2013) provide thorough reviews of threats to, and management recommendations for, *Rana cascadae*, and our recommendations largely follow those of these authors.

Habitat that supports this species in the Lassen area should be protected from modification that negatively impacts hydrology while further research is carried out. Pope et al. (2011) began some habitat restoration measures, and these efforts should be continued (coupled with ongoing monitoring to determine their effects). Fish removal in key populations has also been documented to increase recruitment and should be considered as a management strat-

egy, particularly in the Klamath where a larger number of existing populations might be stabilized before declines can occur. Pope et al. (2011) also proposed experimental treatment for *Bd* in newly metamorphosed frogs. Effective treatments for *Bd* may be essential for the long-term survival of many amphibian species, so these efforts should be further explored and potentially implemented if they are successful. At the same time, a captive colony of Lassen-area *R. cascadae* should be established, as the prospects for long-term survival in the wild appears to be low. If additional research can determine the causes of the declines and effective mitigation measures can be enacted, this captive population could eventually form the basis of a reintroduction program.

Monitoring, Research, and Survey Needs

Monitoring efforts should focus on the few remaining Lassen populations, with additional monitoring of stable populations elsewhere in the range as reference populations. Areas that have undergone habitat restoration or experimental treatments for *Bd* infection will require ongoing monitoring to quantify the long-term

effects of these efforts and to inform further work aimed at controlling the impact of these threats. If additional declines occur, this monitoring will facilitate early detection and, hopefully, provide the background data needed to understand the causes of declines.

As a reintroduction effort may eventually become necessary, it is important to further characterize the extent of intraspecific variation within this taxon now, before additional declines occur. Preliminary genetic work has been initiated, and it should form the basis of additional work that examines fine-scale population differentiation and structure. The obvious initial focus of such genetic work should be to assess the validity of the Lassen and Klamath regions as separate evolutionary units requiring their own management strategies. This work will also help to identify any potential population segments within either region that may qualify for independent management. Finally, additional studies that quantify the interactive effects among different causes of declines would be useful in providing a more complete picture of conservation threats in this taxon.



CALIFORNIA RED-LEGGED FROG

Rana draytonii Baird and Girard 1852

Status Summary

Rana draytonii is a Priority 1 Species of Special Concern, receiving a Total Score/Total Possible of 76% (84/110). During the previous evaluation, it was also considered a Species of Special Concern (Jennings and Hayes 1994a), and it has been listed as federally Threatened since 1996.

Identification

Rana draytonii is a relatively large (2.5–13.8 cm SVL) brown, gray, olive, or reddish-brown frog (Jennings and Hayes 1994a, USFWS 2002, Stebbins 2003). Prominent dorsolateral folds are usually present. Many small black flecks and larger irregular blotches are present on the back, and these occasionally form a network (Baird and Girard 1852). The larger black spots on the back often have a whitish or light center. The ventral surface is whitish or cream with extensive gray or black mottling, often overlain with red or reddish-orange coloration, particularly in the groin (Baird and Girard 1852, Steb-

bins 2003). In general, the red coloration in this species is individually and ontogenetically variable, with the undersides of the feet almost always red in adult animals, although the extent of red elsewhere on the legs and belly varies

California Red-Legged Frog: Risk Factors

Ranking Criteria (Maximum Score)	Score
i. Range size (10)	5
ii. Distribution trend (25)	20
iii. Population concentration/ migration (10)	10
iv. Endemism (10)	7
v. Ecological tolerance (10)	3
vi. Population trend (25)	25
vii. Vulnerability to climate change (10)	7
viii. Projected impacts (10)	7
Total Score	84
Total Possible	110
Total Score/Total Possible	0.76




PHOTO ON PREVIOUS PAGE: California red-legged frog, Alameda County, California. Courtesy of Adam Clause.

from extensive to absent (S. Barry, pers. comm.). The advertisement call is a series of low guttural chucks sometimes followed by a low groan (Elliott et al. 2009).

This species could be confused with the northern red-legged frog (*R. aurora*) where their ranges meet in southern Mendocino County (Shaffer et al. 2004). *Rana aurora* is about 3.5–4.0 cm (SVL) smaller than *R. draytonii*, generally lacks light areas in the centers of dorsal blotches, has proportionally smaller eyes, and lacks vocal sacs (Baird and Girard 1852, Hayes and Krempels 1986, Stebbins 2003).

Taxonomic Relationships

Rana draytonii was initially described as a distinct species, although the original description notes that it is similar in appearance to *R. aurora* (Baird and Girard 1852). Subsequent to the original description, Camp (1917) reclassified the two red-legged frogs as subspecies of a polytypic *R. aurora*. This arrangement persisted, occasionally also including the Cascades frog (*R. cascadae*) as a third subspecies, until the mid-1980s. At this time, a series of studies emerged suggesting that a substantial amount of differentiation between the two forms was present in allozymes, morphology, calling behavior, and oviposition behavior, leading several authors to suggest that they may be distinct lineages with a broad zone of contact (Hayes and Miyamoto 1984, Green 1986a, Green 1986b, Hayes and Krempels 1986). Subsequent analyses of mitochondrial DNA variation supported this view but characterized a narrow zone of contact in southern Mendocino County (Shaffer et al. 2004). Based on both DNA and morphological differentiation, Shaffer et al. (2004) suggested that the two be recognized as distinct species, and since then *R. draytonii* has increasingly been recognized as a species distinct from both *R. aurora* and *R. cascadae*.

Life History

Few data are available on seasonal activity patterns, but coastal populations are probably

active throughout much of the year due to the moderating effect that the Pacific Ocean has on temperature. The timing of reproduction varies from year to year and according to site but occurs from late November to late April (Storer 1925, Fellers 2005b). Breeding occurs in the water, and eggs are attached to emergent vegetation (in clusters of 300 to >4000; Storer 1925, Hayes and Miyamoto 1984). Hatching occurs in 6–14 days depending on water temperature, after which larvae metamorphose in 3.5–7 months (Storer 1925, Wright and Wright 1949). Larvae are known to overwinter at several sites, metamorphosing the following spring (Fellers et al. 2001). *Rana draytonii* is a generalist predator that feeds predominantly on invertebrates but has also been documented to take vertebrate prey including Pacific treefrogs (*Pseudacris regilla*), western toads (*Bufo boreas*), and California mice (*Peromyscus californicus*) (Hayes and Tennant 1985, Arnold and Halliday 1986, USFWS 2002, Davidson 2010). The prey types taken appear to be determined by the size of the frogs, with individual frogs taking most prey types that they can successfully swallow (Jennings and Hayes 1994a) and large frogs taking proportionally more vertebrate prey than small ones. Baldwin and Stanford (1987) reported a large adult preying upon California tiger salamander (*Ambystoma californiense*) larvae. *Rana draytonii* feed both in the water and by foraging in dense riparian vegetation. *Rana draytonii* is active both diurnally and nocturnally, although adults are generally more active at night (Hayes and Tennant 1985; G. Fellers, pers. comm.).

Wading birds, raccoons, and garter snakes (*Thamnophis sirtalis* and *T. hammondi*) are important native predators on this species (Cunningham 1959b, Jennings and Hayes 1994a). Nonnative fishes are also important predators on larvae and recent metamorphs (Schmieder and Nauman 1994, USFWS 1999).

Habitat Requirements

Rana draytonii chiefly inhabits ponds, although it also uses marshes, streams, lagoons, and other waterways throughout most of its range.

In southern California (from Ventura County southward) it seems to favor slow-flowing streams rather than ponds or pools. Breeding takes place primarily in ponds (at least in central and northern California) and less frequently in quiet pools in streams (Stebbins 2003, Fellers 2005b). This species will also utilize ephemeral water bodies for breeding, although nearby permanent water is probably required to maintain populations over the long term (Jennings 1988a). After breeding, adults often disperse along nearby shaded streams. Similar to *R. boylei*, whose vulnerable early life stages (embryos and tadpoles) are susceptible to ill-timed flow fluctuations controlled by upstream dams and diversions, *R. draytonii* populations breeding in stream habitats suffer from decreased recruitment after anthropogenic perturbation of natural flow regimes (S. Kupferberg, pers. comm.).

Optimal aquatic habitat has traditionally been thought to include dense riparian vegetation overhanging deep (>0.7 m) slow-moving pools (Hayes and Jennings 1988). More recent work has documented an additional, more complex relationship between aquatic vegetation and introduced bullfrogs. D'Amore et al. (2009) documented that *R. draytonii* spend more time in vegetative cover when bullfrogs are present and more time in the open when bullfrogs are removed from ponds, suggesting that the optimal amount of vegetation is somewhat context-dependent for *R. draytonii*. In addition, surveys of 85 ponds occupied by *R. draytonii* in the East Bay Regional Park District showed that there were no significant differences in adult frog density among ponds with 0%, $\leq 15\%$, or $>15\%$ emergent vegetation, but tadpoles and metamorphs were more abundant in the most open ponds (Bobzien and DiDonato 2007). Outside of the breeding season when conditions are wet, and especially during rainfall, adult frogs will disperse from the breeding habitat and will move to upland sites, where they are often found under logs, rocks, and other debris (USFWS 2002, Bulger et al. 2003, Fellers and Kleeman 2007). At some

sites, populations appear to consist of both migratory (11–22% of the adult population) frogs that move 200–2800 m and resident frogs that remain at the breeding site (Bulger et al. 2003). Fellers and Kleeman (2007) found that adult female frogs were more frequently migratory than males, although migration behavior did not differ between the sexes among those individuals that did migrate.

Distribution (Past and Present)

Historically, *Rana draytonii* ranged throughout the Sierra Nevada foothills and the Coast Range mountains south of Elk Creek in southern Mendocino County, California, southward to the Arroyo Santo Domingo, Baja California Norte, Mexico (Hayes and Krempels 1986, Jennings and Hayes 1994a, Crismer 2002, Shaffer et al. 2004). In California, this taxon historically ranged through at least 46 counties, but it is now apparently extirpated from 24 of these (USFWS 1996). It is unclear whether reproductive populations of *R. draytonii* were present in most of the Central Valley, and it is possible that the few valley records represent waifs washed downstream from Sierran populations (G. Fellers, pers. comm.; S. Barry, pers. comm.). If they were present in the Central Valley, they were extirpated before 1960. Populations in the Sierra Nevada may have been connected to the largest remaining populations of the species in the Coast Ranges through the lower Cascade and Tehachapi Ranges (S. Barry, pers. comm.), but today they are isolated (USFWS 2002). A recent comprehensive survey of museum specimens and historical records identified 21 historical localities for this species in the Sierra Nevada. Follow-up surveys at 20 of these 21 sites found that the species persists in large numbers in at least 1 site, there are populations at 6 additional sites, and at least a single individual documented at 3 more sites (Barry and Fellers 2013).

Strong overall declines have clearly occurred across most of the large range of this species, particularly in the southern portion of the range. In the Bay Area and Coast Ranges,

populations are more robust, although severe localized declines have been documented (reviewed in USFWS 2002). In southern California, *R. draytonii* has declined drastically through the Transverse and Peninsular Ranges, and very few populations now persist in Los Angeles, Riverside, and Ventura Counties (USFWS 2002, and references therein). One population is known from Santa Cruz Island, although this apparently is an introduction (Sweet and Leviton 1983, Jennings 1988b). The known elevational range of *R. draytonii* occurs from near sea level to 1500 m, although most populations occur below 1050 m (USFWS 2002, Barry and Fellers 2013). Some higher-elevation populations may be introductions (unpublished data reported in Jennings and Hayes 1994a).

Trends in Abundance

Drastic and ongoing declines have been documented throughout parts of this species' range. Many of these declines have resulted in extirpation of populations, and in many areas where this taxon persists, declines in abundance have occurred. Food market collection in the late 1800s apparently drove much of the initial declines (Jennings and Hayes 1985). By 1879, the species had already become rare around San Francisco due to the market trade (Lockington 1879). Population trends of the species in the Sierra Nevada foothills are somewhat unclear, since several new, large (>100 breeding adults) populations have recently been discovered (e.g., in Placer County). However, in southern California, population densities are uniformly low (<25 adults frogs) and generally declining (USFWS 2002). The sole remaining population known in Riverside County at the Santa Rosa Plateau, which was at least somewhat genetically distinctive (Shaffer et al. 2004), is now extirpated.

Nature and Degree of Threat

The largest threat facing *Rana draytonii* is probably habitat loss and alteration, resulting from urbanization and agriculture. The large-scale conversion of habitat to agricultural uses has

also resulted in an increase in pesticide exposure, which may have strong negative impacts on this species (Davidson et al. 2002). This effect is particularly strong for cholinesterase-inhibiting pesticides (Davidson 2004), although the species still persists in some heavily agricultural settings in Monterey and Santa Cruz Counties. Additional and ongoing fragmentation of habitats, conversion of wetlands to other uses, and modifications to the hydrology of wetlands also likely have detrimental impacts.

The effect of introduced species, in particular bullfrogs, has been studied both empirically (Moyle 1973) and from a modeling perspective (Doubledee et al. 2003). There is a strong overall negative impact of bullfrogs on native *R. draytonii*, although coexistence of the two species can occur in nature. Human-modified aquatic and terrestrial habitats in central California (Elkhorn Slough, Monterey County) favor introduced bullfrogs compared to native *R. draytonii* (D'Amore et al. 2010). The bullfrog is also a strong competitor with, and predator on, multiple life stages of *R. draytonii*. In addition, crayfish, mosquitofish (Lawler et al. 1999), and other introduced predaceous fishes likely have negative impacts on this species, although this also needs further study (Hayes and Jennings 1986, Fisher and Shaffer 1996, Fellers 2005b).

Chytrid fungus (*Bd*) is known to have caused serious declines in many amphibian species and has been detected in *R. draytonii* in nature. However, the direct impact *Bd* has on *R. draytonii* appears to be relatively slight. In a laboratory setting, *R. draytonii* is susceptible to chytrid infection, but frogs can clear their infections, do not die from the infection, and suffer no growth consequences when they have access to unlimited food (Padgett-Flohr 2008). In nature, across a landscape of ponds where *Bd* presence and absence fluctuated between wet and dry years, *R. draytonii* were generally uninfected and found to be significantly associated with uninfected ponds (Padgett-Flohr 2010).

Predicted climate change over much of California will affect *R. draytonii*, as well as most

other pond- and stream-breeding amphibians. In particular, warmer average temperatures, generally reduced levels of precipitation, and increased variability in the timing of rainfall are all predicted to occur (PRBO 2011). While the precise effects of these shifts will vary regionally and at the watershed level, the permanence and reliability of breeding sites are generally predicted to decrease under climate change predictions.

Status Determination

Rana draytonii automatically qualifies as a Species of Special Concern because it is listed under the federal but not state Endangered Species Act. However, sharp declines in both range and abundance, coupled with a variety of ongoing threats to long-term survival, also combine to warrant a Priority 1 Species of Special Concern status.

Management Recommendations

Management of *Rana draytonii* should mirror the guidelines in the USFWS recovery plan for this taxon (USFWS 2002). As further management needs are defined and existing management strategies are refined (through 5-year reviews or other avenues), state-level management should be adjusted accordingly.

The most important management needs for this taxon currently are the protection of habitat that supports the species, reduced pesticide exposure, and elimination of nonnative predators. Land conversion and additional fragmentation should be avoided wherever possible, and adequate, complex upland habitat should be available in order to allow migration to occur naturally. Fellers and Kleeman (2007) found that the median distance of movement away from breeding ponds was 150 m and that there were some long-distance movements up to 1400 m. Unpublished radiotelemetry observations from the East Bay Regional Park District (S. Kupferberg, pers. comm.) demonstrated that ground squirrel burrow density, sometimes more than 100 m from the aquatic habitat, was also a key component of habitat quality.

Taken together, these studies indicate that large tracts of terrestrial habitat are important (to accommodate both short- and long-distance dispersal) and that a healthy population of ground squirrels (and possibly other burrowing rodents) may be essential for long-term population viability.

Finally, pesticide use should be curtailed in areas where this species occurs, including areas upwind where pesticides are likely to be blown into areas that support this species. Unpublished data from the East Bay Regional Park District (S. Kupferberg, pers. comm.) indicate that cattle-grazing does not appear to negatively impact this species.

Monitoring, Research, and Survey Needs

Further research is needed to determine what the precise impacts many of these threats identified above are having on *Rana draytonii*. Surveys of private land in the Sierra Nevada are slowly revealing the presence of extant populations that were previously missed (S. Barry, pers. comm.), suggesting that this may be a fruitful strategy elsewhere in the range as well. Managers should partner with private landowners to gain access and survey for remaining populations of this species in areas where it has previously been thought to be extirpated, and these populations, which may be very small in size, should be monitored regularly.

Finally, the only range-wide genetic analysis of the species thus far conducted was based purely on mitochondrial DNA (Shaffer et al. 2004), and supporting data from a large set of nuclear DNA markers is badly needed. In particular, the potential genetic break between populations north and south of Santa Barbara County, and the genetic affinities of remnant populations from southern California and Baja California, Mexico, will form an important part of future management.

Additional monitoring, research, and survey needs are covered in depth in the USFWS recovery plan for this taxon. We refer the reader to this document for more information (USFWS 2002).



NORTHERN LEOPARD FROG

Rana pipiens Schreber 1782

Status Summary

Rana pipiens is a Priority 1 Species of Special Concern, receiving a Total Score/Total Possible of 73% (80/110). During the previous evaluation, it was also designated as a Species of Special Concern (Jennings and Hayes 1994a).

Identification

Rana pipiens is a medium-sized ranid frog with strong, continuous dorsolateral folds that do not angle inward posteriorly. Its dorsal coloration is green to brown with large well-defined black or dark-brown oval or round spots. Each spot is ringed with a narrow band of white or cream. The ventral coloration is white or cream with no mottling or other dark markings (Stebbins 2003). The call is a low, snore-like trill, often followed by low chuckling and/or grunts (Stebbins 2003, Elliott et al. 2009).

Within its range in California, this species can potentially be confused with the Oregon spotted frog (*R. pretiosa*). However, *R. pretiosa* has much smaller, more irregular spots, which

often have diffuse borders and are not ringed in white. It also has conspicuous red or salmon markings on the underside, which *R. pipiens* lacks. Other members of the leopard frog complex in California, the lowland leopard frog

Northern Leopard Frog: Risk Factors

Ranking Criteria (Maximum Score)	Score
i. Range size (10)	10
ii. Distribution trend (25)	20
iii. Population concentration/ migration (10)	10
iv. Endemism (10)	0
v. Ecological tolerance (10)	0
vi. Population trend (25)	20
vii. Vulnerability to climate change (10)	10
viii. Projected impacts (10)	10
Total Score	80
Total Possible	110
Total Score/Total Possible	0.73




PHOTO ON PREVIOUS PAGE: Northern leopard frog, Washington County, Utah. Courtesy of William Flaxington.

(*R. yavapaiensis*) and the Rio Grande leopard frog (*R. berlandieri*), have dorsolateral folds that are discontinuous and angle inward posteriorly. In addition, both are yellow ventrally. The Cascades frog (*R. cascadae*) has more numerous, small, irregular black dots that are not ringed in white.

Taxonomic Relationships

The taxonomic history of the leopard frog species complex, and *Rana pipiens* in particular, is complicated (Hillis 1988) and remains incompletely understood. The name *R. pipiens* previously included all members of the leopard frog complex from Canada south to Panama, including *R. yavapaiensis*, also native in California, and the introduced *R. berlandieri*. However, this concept of a single wide-ranging leopard frog species changed in the last several decades, and over a dozen species are recognized at present. The current taxonomy of the *R. pipiens* complex was initially based on variation in morphology and vocalizations (Pace 1974). Subsequent work including molecular analyses recognized several additional taxa and clarified relationships among the contained species (Platz and Mecham 1979, Hillis et al. 1983, Platz and Frost 1984, reviewed by Hillis 1988).

Frost et al. (2006a) recommended placing this species and many other North American ranids in the genus *Lithobates*, although this proposal and the analyses that support it are controversial (Crother 2009, Frost et al. 2009a, Pauly et al. 2009). We retain the traditional taxonomy here to maintain stability pending further analyses.

Life History

No life history data for California populations have been published. Because *Rana pipiens* in California are a mixture of introduced and presumably native populations (see the "Distribution" section) and live on the extreme western edge of the species' range, we are reluctant to use information from more easterly populations as a proxy for those that occur in California. In Colorado, breeding occurs during the

first spring nights that have relatively "mild" temperatures near or above freezing (Corn and Livo 1989), and this presumably is also the case in California. Tadpoles are present through the summer months and are not known to overwinter, suggesting a late summer or fall metamorphosis. Further east, adults and juveniles are known to range far from water and breeding sites (Dole 1971), although it is unknown if this also characterizes California populations. Range-wide, *R. pipiens* is a generalist predator, feeding on a wide variety of arthropods and small vertebrates (Knowlton 1944, Linzey 1967, Harding 1997), and this presumably also characterizes the species in California.

Habitat Requirements

Despite the paucity of records from California, this species is known from a variety of habitats, including small streams, rivers, and lakes (Storer 1925, Stebbins 1951, Jennings and Hayes 1994a). *Rana pipiens* occupies a wide variety of habitat types throughout its range, so we are hesitant to speculate on microhabitat requirements in California. Generally, the species hibernates underwater and requires aquatic habitats that do not freeze solid during winter (Emery et al. 1972, Licht 1991), and this presumably is also the case for California populations. Nearby damp upland habitat is utilized for foraging during the active season (Dole 1967). The species has been found in a variety of open grassy areas and meadows, although heavily grazed areas and cultivated fields do not appear to be suitable (Pope et al. 2000). In the Midwestern United States, the presence of quality upland foraging habitat seems to affect the abundance of this species. When grasslands were restored around suitable pond-breeding habitat, the density of frogs increased markedly (K. Mierzwa, pers. comm., in Pope et al. 2000).

Distribution (Past and Present)

Outside of California, *Rana pipiens* ranges widely across North America, from Nova Scotia and Newfoundland, Canada, west to Washing-

ton and Nevada. In California, *R. pipiens* populations that may be native are known from Modoc and Siskiyou Counties, the Lake Tahoe basin, and the upper Owens Valley (Jennings and Fuller 2004), although some workers question whether the latter two regions constitute natural, as opposed to purely introduced, populations (S. Barry, pers. comm.). Numerous introductions have occurred throughout the state, including some within the putative native range. The vicinity of Fallen Leaf Lake in the Lake Tahoe Basin is one such example (Bryant 1917). It is also possible that putatively native populations of this frog are all the result of human introductions, and determining their status is an important research priority. The upper Owens Valley supports tiger salamander populations that were recently shown to be introduced (Johnson et al. 2010), demonstrating that similarly distributed nonnative species have been established in this region. The tiger salamander introductions occurred as a consequence of the fishbait industry (Riley et al. 2003), which also sometimes sells leopard frog tadpoles and adults.

We are not aware of any additional recent records in California beyond those reported by Jennings and Hayes (1994a), though an unverified sight record of a "spotted frog" in Surprise Valley, Modoc County, California, could have been *R. pipiens*. However, the circumstances and description of this frog make it more likely that it was *R. pretiosa*, another California Species of Special Concern (see that species account for additional information).

Trends in Abundance

Trends in abundance for California populations of *Rana pipiens* are difficult to interpret because of the uncertainty regarding which populations are native or introduced. However, assuming that historical California populations are native, severe declines have clearly occurred. We are aware of only scattered sight records for the species over the last two decades. Jennings and Hayes (1994a) reported two relatively recent sight records in the early 1990s from Siskiyou

and Inyo Counties. Macey and Paperfuss (1991a) reported that leopard frogs occurred on the east side of the White Mountains below Boundary Peak, though they failed to detect the species in follow-up surveys (T. Paperfuss, pers. comm.). More recent surveys of historical localities in the Owens River also did not detect this species and found that much of the habitat currently appears to be unsuitable (Becker and Henderson 2010). We are not aware of any presumed-native populations of this species occurring in the state since these records. Elsewhere in its range, *R. pipiens* has undergone severe declines and localized extirpations, particularly in the western parts of the United States (reviewed by Rorabaugh 2005).

Nature and Degree of Threat

Habitat modification is probably the most important threat for *Rana pipiens* in California. *Rana pipiens* forages in upland habitat having moderately tall vegetation with a moist substrate. Livestock grazing in these habitats tends to reduce vegetation height, which leads to drying of the substrate, apparently rendering this habitat unsuitable for the frog. It is likely that this process contributed to the declines observed in both the Owens Valley and the Modoc Plateau areas where most California records for *R. pipiens* are concentrated. Changing hydrology elsewhere in the range has led to the extirpation of some local populations (Corn and Fogleman 1984). Given that California populations are at the western range limit of the species, projected climate changes may have a strong effect in the state. Current models project warmer summer and winter temperatures, decreases of 8–21% of annual precipitation, and a 34% decrease in snowpack (PRBO 2011). Taken together, these climate projections indicate that the moist soil and wetland complexes favored by this species will probably decrease in the Great Basin of California, further reducing the already sparse habitat for this species.

Some studies have detected significant negative impacts from pesticides on *R. pipiens*,

although the importance of this threat in nature is not well understood. In other parts of their range, *R. pipiens* are known to be sensitive to herbicides and pesticides used in agriculture (Relyea 2008, Relyea and Jones 2009), and mixtures of these chemicals can result in 99% mortality rates (Relyea 2008). However, the evidence on this topic is complex and dependent on the specific chemicals tested. A popular herbicide consisting of a mixture of glyphosate and POEA (commonly marketed under the commercial name Roundup®) is one such example. Some studies have found limited impacts from these chemicals and concluded that direct mortality in wild populations from this herbicide is unlikely (e.g., Wojtaszek et al. 2004), while other studies have found very strong direct lethal effects (e.g., Relyea 2005b). When direct lethal effects were not found, several studies demonstrated that chemical contaminants can have lethal impacts when combined with other stressors (e.g., predator cues; Relyea 2005a) or sublethal detrimental effects such as decreased immune system functionality (Christin et al. 2003, Gilbertson et al. 2003, Rohr et al. 2008). These seemingly unpredictable effects of agrochemicals may depend on specific populations and conditions in a local area (Relyea 2005b). Although these results are both complex and sometimes contradictory, substantial evidence exists that environmental contaminants are likely to have significant impacts on *R. pipiens* and other amphibians in California (e.g., Davidson et al. 2002, Davidson 2004).

Other potential threats to *R. pipiens* include introduced exotic bullfrogs and predatory fishes, and extensive habitat modification associated with agriculture (Hayes and Jennings 1986).

Status Determination

Rana pipiens' small range in California coupled with severe declines drives the high score for this species. None of these threats are currently being reversed, so it is reasonable to expect additional declines in the future, assuming that native populations still exist in California.

Rana pipiens is sensitive to localized extirpation due to drought (Corn and Folgeman 1984), and the expected increase in temperature and decrease in precipitation due to climate change are likely to have additional negative impacts. The combination of these factors justifies a Priority I status.

Management Recommendations

The development of an effective management strategy will largely depend on finding remnant populations in the state, carrying out research on the life history of those specific populations to determine their habitat needs, and then taking a proactive management and habitat restoration approach to recover it in its native range. A key first step with any remnant population is to determine whether it is native or introduced. Researchers can most easily accomplish this using DNA markers, and we recommend that larval tail tips be collected for any population that is discovered. A considerable amount of phylogenetic work, particularly using mitochondrial DNA markers, has been published for this species, and straightforward DNA sequencing of California animals should allow them to be placed into a phylogenetic context with other *Rana pipiens* from across the species' range. This approach was used by Johnson et al. (2010) and demonstrated that potentially native populations of tiger salamanders (*Ambystoma tigrinum*) were in fact nonnative introductions. If native populations of *R. pipiens* are found, the habitat supporting them should be protected in order to reduce potential threats such as nonnative predators, agricultural disturbance, grazing, off-highway vehicle use, pesticide applications, and changes to local hydrology. If nonnative populations are found, managers should evaluate their potential to spread and pose a threat to other native taxa. In certain cases, removal programs could be effective at mitigating threats posed by nonnative *R. pipiens*.

Monitoring, Research, and Survey Needs

Comprehensive surveys of historical localities as well as the Modoc Plateau area, including the

Goose Lake Basin and the Warner Mountains, should be conducted to determine whether any viable populations persist in California and to identify areas of potential habitat for ongoing surveys. The most recent records for this species come from the vicinity of Owens Valley, and all drainages flowing into the valley should be carefully surveyed. It is critically important that tissue samples be collected from any extant populations that are found so that frogs can be genetically characterized with respect to their introduced or native status.

Given our current lack of information about the life history of this species in California, basic ecological research is a key priority for any native populations that remain in the state.

Information about habitat preferences and requirements, demography, and timing of key life history events would all improve our ability to conserve remnant populations of *Rana pipiens*.

Finally, if remnant populations are found, multi-locus microsatellite or single nucleotide polymorphism DNA data should be analyzed to estimate the effective population size and potential connectivity with other remaining populations. If populations are determined to be native, small, and genetically isolated, *R. pipiens* could be a prime candidate for human-mediated translocations to establish new populations in currently unoccupied habitat patches.



OREGON SPOTTED FROG

Rana pretiosa Baird and Girard 1853b

Status Summary

Rana pretiosa is a Priority 1 Species of Special Concern, receiving a Total Score/Total Possible of 0.82 (82/100). During the previous evaluation, it was also designated as a species of special concern (Jennings and Hayes 1994a) and it was listed as federally Threatened in 2014 (USFWS 2014). We are aware of only two unverified site records for this species in California in the last 25 years.

Identification

Dorsally, *Rana pretiosa* is a dark-brown, reddish, or greenish frog with black spots or blotches (McAllister and Leonard 1997). The dorsal blotching is usually irregular around the edges, rather than sharply demarcated, and has a small light spot in the center of the larger spots. The venter is usually mottled and has a base color that changes from cream white at the chin to orange more ventrally (Dunlap 1955, Stebbins 2003). The ventral coloration often appears to be superficial or “painted on” (Dun-

lap 1955, Nussbaum et al. 1983). Like many California ranids, this species has a prominent light stripe below the eye (particularly so in juveniles) and thin dorsolateral ridges that

Oregon Spotted Frog: Risk Factors

Ranking Criteria (Maximum Score)	Score
i. Range size (10)	10
ii. Distribution trend (25)	20
iii. Population concentration/ migration (10)	10
iv. Endemism (10)	0
v. Ecological tolerance (10)	7
vi. Population trend (25)	25
vii. Vulnerability to climate change (10)	Data deficient
viii. Projected impacts (10)	10
Total Score	82
Total Possible	100
Total Score/Total Possible	0.82




PHOTO ON PREVIOUS PAGE: Oregon spotted frog, Lane County, Oregon. Courtesy of Troy Hibbitts.

dissolve into a series of raised dots two-thirds to three-quarters of the way down the back. The call consists of a series of faint clicks, repeated roughly seven times in rapid succession (Briggs 1987, Stebbins 2003, Elliott et al. 2009).

Within its California range, this species is most likely to be confused with the Cascades frog (*R. cascadae*). Although similar, *R. cascadae* spots tend to have sharply defined edges, no light centers, and appear to be on the surface of the skin, reminiscent of black ink being splattered on the frog (Stebbins 2003). In addition, the underside of the legs are yellow tan in *R. cascadae* (reddish in *R. pretiosa*), the eyes are oriented dorsally when viewed from above in *R. pretiosa* (oriented outwardly in *R. cascadae*), and *R. pretiosa* has full, rather than partial webbing between the toes of the rear legs. The Columbia spotted frog (*R. luteiventris*) may also occur in California, and it could also be confused with *R. pretiosa* (see the "Distribution" section).

Taxonomic Relationships

Green et al. (1996, 1997) divided *Rana pretiosa* into two species, *R. pretiosa* and *R. luteiventris*, based on morphology and allozyme variation. The two taxa are morphologically similar (usually distinguishable in the field based on the ventral mottling in *R. pretiosa*; M. Hayes, pers. comm.), but preserved specimens can usually be differentiated with a series of head measurements (Green et al. 1997). The two species are also diagnosable using allozymes (Green et al. 1996) and mitochondrial DNA cytochrome-*b* sequence (Funk et al. 2008).

Life History

No data on life history of California populations exist and much of the data from elsewhere in the range occurred before the partitioning of *Rana pretiosa* and *R. luteiventris*. As California populations of *R. pretiosa* are at the extreme southern edge of the species' range, the timing of life history events may occur earlier relative to those reported from more northerly sites, although the high elevation of California sites may compensate for any potential latitudinal

gradient. California populations were geographically closest to Oregon frogs from the Klamath basin, and those populations may serve as the best models for California.

Frogs emerge from hibernation as soon as the winter thaw permits (Stebbins 2003) and water temperatures rise to about 6°C (C. Pearl, pers. comm.). *Rana pretiosa* breeds explosively soon after emergence, usually over a 1- or 2-week period. Males often congregate in shallow water and begin to call (Licht 1969, Nussbaum et al. 1983). Egg masses are deposited together in large groups in vegetated margins of large permanent aquatic habitats, usually at the high-water mark. The species can experience high egg mass mortality when waters recede rapidly, leading to stranding, desiccation, and/or freezing (Licht 1971, Briggs 1987). However, eggs from multiple sites in Oregon were found to resist near-freezing temperatures as long as they remained beneath the water surface (Bowerman and Pearl 2010). Artificially incubated egg masses hatch in as few as 72 hours to as many as 400 hours, depending on temperature (25°C and 10°C, respectively), followed by metamorphosis in approximately 4 months (Licht 1971).

Males appear to have lower survivorship than females, presumably due to the longer periods of time that they spend in breeding congregations and the resulting exposure to predation (Licht 1974, Chelgren et al. 2008). Post-metamorphic frogs consume a wide variety of invertebrate prey including insects, occasional mollusks, and crustaceans, as well as small vertebrates including anurans (Nussbaum et al. 1983, Licht 1986b, Pearl and Hayes 2002, Pearl et al. 2005b).

Habitat Requirements

Information on habitat utilization in California is very limited, although habitat requirements are better studied elsewhere in the range. The species appears to seasonally use different habitat types (Watson et al. 2003, Chelgren et al. 2008). *Rana pretiosa* is highly aquatic and rarely found away from the water (Licht 1986a). It frequently uses temporary pools, ditches, and

other shallow water sources, but nearby deep permanent water is always required and serves as a refuge for adult frogs during dry parts of the year and during drought (McAllister and Leonard 1997, Watson et al. 2003). Breeding occurs in shallow water with aquatic vegetation (Licht 1971, Watson et al. 2003). In Oregon, oviposition sites occurred, on average, 14.1 m (range 0.08–35.0 m) from the shore in water that was 18.5 cm deep (range 1–57 cm) (Pearl et al. 2009). At one site in Washington, the species overwintered in shallow water, where it buried itself at the base of emergent plants (Watson et al. 2003). Overwintering in flowing springs has also been documented (Chelgren et al. 2008). Overland dispersal appears to be quite limited, and the species may require habitat where the shallow-water breeding and overwintering habitats are connected to deep-water refuge habitat by intervening water during early spring and late fall to allow inter-habitat migrations (Watson et al. 2003).

The habitat requirements for *R. pretiosa* have likely contributed to its declines. The diversity of habitat types that are used, coupled with the requirement that they are connected by intervening stretches of water, is fairly specific and is probably only common in large, relatively intact wetland complexes. These complexes are becoming increasingly rare throughout the species' range as landscapes are drained and converted to agriculture and grazing.

Data are limited on effects of grazing on this species. At one site in western Washington where reed canarygrass (*Phalaris arundinacea*) forms dense stands, Watson et al. (2003) suggested that grazing could help open patches and make them suitable for *R. pretiosa*. However, grazing also has the potential to reduce water quality and cover from predators. Additional work is needed on how the timing and intensity of grazing affect frog behavior and habitat use.

Distribution (Past and Present)

Few localities for *Rana pretiosa* have been documented in California, and all known localities

appear to be extirpated. Historically, *R. pretiosa* occurred in the northeastern corner of California, ranging south to Plumas and Tehama Counties and west to the eastern portions of Siskiyou, Shasta, and Tehama Counties (Slevin 1928). Within this range, the species has been found in scattered localities in Modoc, Shasta, and Siskiyou Counties (Stebbins 1972, Jennings and Hayes 1994a), with the last documented record occurring in a woodpile in Cedarville, Modoc County, in 1989 (Jennings and Hayes 1994a). This last record is somewhat anomalous, since the frog was found in a heavily modified area near the town center of Cedarville, in habitat that seems to be unsuitable for the frog. Given the very specific habitat requirements of *R. pretiosa*, the fact that no specimen from the site was ever examined by a herpetologist and no vouchers exist, it is possible that this is a misidentified or human-introduced specimen (L. Groff, pers. comm.; M. Hayes, pers. comm.). It remains possible that isolated populations still persist, particularly in remote portions of the Warner Mountains and on private land in Surprise Valley, Modoc County. Fairly recent surveys in the Warner Mountains, Modoc Plateau, and Pitt River drainage failed to locate any individuals (Jennings and Hayes 1994a, Groff 2011). There is an unverified sighting of a "spotted frog" in Surprise Valley from November 2008 (L. Gray, pers. comm.), but a follow-up survey at this locality revealed only *Psuedacris regilla*. A more recent survey comprising 18 localities selected using a species distribution model for this species did not detect *R. pretiosa* in California (Groff 2011), although the southernmost extant locality in Oregon is only about 10 km from the state border. Between 2012 and 2013, USFWS biologists conducted additional surveys at 12 sites within the Pit River watershed and Warner Mountains. Again, no evidence of *R. pretiosa* was found (USFWS-Klamath Falls Field Office, unpublished data, 2013).

Outside of California, *R. pretiosa* is patchily distributed from extreme southwestern British Columbia, south through Washington and

Oregon (Green et al. 1997). This distribution is fragmented, and the species has undergone severe declines through most of its range (McAllister et al. 1993, Green et al. 1997). Declines are thought to have occurred disproportionately in lowland areas, and over two-thirds of the remaining populations occur along the crest and eastern slopes of the Cascade Range (Pearl et al. 2009).

It is possible that some *R. pretiosa* in California, particularly those east of the Warner Mountains in Modoc County, could actually be *R. luteiventris*. There are known *R. luteiventris* populations approximately 16 km north of the California border on the eastern slopes of the Warner Mountains, making the presence of *R. luteiventris* in California plausible (Funk et al. 2008; M. Hayes, pers. comm.). However, the species has not been documented in California.

Trends in Abundance

No abundance data for California populations exist. Reports from parts of the Willamette Valley, Oregon, and Puget Lowlands, Washington, suggest that *Rana pretiosa* was common in those areas around the 1930s. Declines are thought to have been occurring for a large part of the twentieth century (Dumas 1966, McAllister et al. 1993, Pearl and Hayes 2005). At one time, the species was apparently common in Warner Valley, Oregon, immediately north of Surprise Valley in California (Cope 1883). Any remaining populations in California are likely to be isolated and on private land that has not been surveyed. A recent species distribution model generated a set of potential sites, some of which were surveyed, but no California populations were found (Griff 2011).

Nature and Degree of Threat

At least four major factors have likely contributed to the decline of *Rana pretiosa* in California. First, the species has been strongly impacted by the loss of the extensive wetland complexes that were once common in northern California. As land has been drained and modified for livestock grazing and agriculture, the

overall amount of available acreage that provides the precise suite of habitat types used by this species has declined. This loss of wetland habitat is further exacerbated by climate projections for northeastern California, which predict increasing temperatures, strongly decreasing precipitation, and reduced snowpack (PRBO 2011); all of these changes will reduce permanent wetlands and place increasing demands on the remaining aquatic habitat. Second, *R. pretiosa* appears to be sensitive to relatively low levels of nitrates and nitrites resulting from agricultural runoff (i.e., those meeting EPA allowances for drinking water; Marco et al. 1999). This observation is consistent with the precipitous declines observed in lowland Oregon and Washington populations, which have been more heavily impacted by agriculture than higher-elevation populations. Application of the pesticide DDT was also correlated with die-offs in the closely related *R. luteiventris* in northern Oregon (reported as *R. pretiosa*; Kirk 1988). Third, the species appears to be sensitive to introduced exotic predators, particularly bullfrogs and exotic fishes. Some data indicate that it is likely more sensitive to the presence of bullfrogs than other native ranid frogs. In areas where *R. aurora* and *R. pretiosa* are sympatric, stronger declines were observed in *R. pretiosa* than *R. aurora* in areas where bullfrogs have invaded (Pearl et al. 2004). Laboratory experiments also demonstrate a differential impact of bullfrogs on *R. pretiosa* relative to *R. aurora*, likely due to *R. pretiosa*'s more strongly aquatic life history (Pearl et al. 2004). Bullfrogs have also been hypothesized to negatively impact small *R. pretiosa* populations via reproductive interference (Pearl et al. 2005c). In combination with the well-documented effects of non-native fishes on western ranid frogs (Adams 1999, Lawler et al. 1999, Adams 2000, Joseph et al. 2011), this suite of nonnative predators is likely to have a strong negative effect on *R. pretiosa* populations. Finally, *Bd* has been found to be present in remaining populations of *R. pretiosa* (Pearl et al. 2007, Hayes et al. 2009), although experimental work suggests that the

species may be resistant (Padgett-Flohr and Hayes 2011). However, given the importance of *Bd* in some anuran declines, further work on its impact on *R. pretiosa* is warranted.

Given the rarity of *R. pretiosa* records from California and our lack of historical population parameters, it is impossible to differentiate between these causes. However, it is reasonable to assume that several or all of these factors were involved in the decline of the species in California.

Status Determination

The limited California range of *Rana pretiosa* and its apparent extirpation from the few known historic localities are the main drivers for its high score. The paucity of historical records in California suggests that this taxon may have historically been rare in the state, and its specialized ecological requirements (large permanent wetlands, specialized sub-habitats for breeding, hibernation, and growth) make it inherently sensitive to declines. Together, these factors justify a Priority 1 designation for this species.

Management Recommendations

Ongoing management efforts for this species should be coordinated through the range-wide conservation strategy that the Washington Department of Fish and Wildlife is leading and the California Department of Fish and Wildlife is participating in (B. Bolster, pers. comm.). Cushman and Pearl (2007) recently assessed *Rana pretiosa* conservation needs and provided a detailed roadmap for management of this species. Our recommendations largely follow theirs. If the surveys outlined below identify any remaining populations of this species in the state, the wetland habitat supporting the population should be protected from fragmentation and modification, including the introduction of exotic fishes and amphibians. Captive populations of this species should also be established to serve as assurance colonies, should the last wild populations go extinct. If continued surveys suggest that the species is

extirpated from California, captive breeding and reintroduction programs could be initiated with Oregon animals if appropriate habitat can be identified and protected. Given the very high levels of genetic differentiation and population structure found among extant Oregon and Washington populations (Blouin et al. 2010), populations from the southern Klamath Basin genetic unit are probably the best candidates for such a reintroduction in California. Beyond these two steps, effective management of this taxon in California will require additional research into the causes of decline.

Monitoring, Research, and Survey Needs

Comprehensive surveys throughout *Rana pretiosa*'s known historic range should be conducted to determine if any populations persist in the state. Surveys of remaining large wetland complexes are particularly important, as are surveys of potential habitat on private property. A recent species distribution model (Groff 2011) identified and surveyed some, but not all, of the predicted localities that may support this species in California, and this study provides an excellent starting point for additional surveys. Significant habitat that has not yet been surveyed remains on private property, particularly east of the Warner Mountains (although *R. luteiventris* may replace *R. pretiosa* in this area). The aforementioned recent surveys made a particular effort to gain access to private land, but permission was only granted in approximately 15% of cases (Groff 2011). Future surveys should continue to build partnerships with private stakeholders and survey large wetland complexes on private lands. If any populations are found, nonlethal tissue samples should be collected so that species identification can be verified with molecular data.

Should any populations be located, a monitoring program in conjunction with life history research should immediately be initiated with the goal of quantifying population sizes and connectivity (if multiple adjacent populations are found) and to allow for a better understanding of habitat requirements and causes of decline in this species. Molecular genetic

studies using microsatellite and/or single nucleotide polymorphism data from multiple nuclear markers can provide valuable insights into historical population declines/expansions and should be conducted if any native populations are discovered. In addition, given the very

high levels of population structure found among extant Oregon and Washington populations, any California populations should be surveyed for genetic variation and integrated into the existing species-wide genetic dataset (Blouin et al. 2010).



LOWLAND LEOPARD FROG

Rana yavapaiensis Platz and Frost 1984

Status Summary

Rana yavapaiensis is a Priority 1 Species of Special Concern, receiving a Total Score/Total Possible of 74% (63/85). During the previous evaluation, it was also considered a Species of Special Concern (Jennings and Hayes 1994a). *Rana yavapaiensis* has not been confirmed to occur in California since 1965 (Jennings and Hays 1994a).

Identification

Rana yavapaiensis is a medium-sized ranid frog (4.6–8.7 cm SVL) with prominent dorsolateral folds that are discontinuous and angle inward posteriorly (Platz and Frost 1984). The coloration is variable, but is generally gray green, gray brown, or tan with irregular blotches above and cream or white on the venter. The ventral pelvic region is yellow, and this sometimes extends onto the legs. In older individuals, there is also dark mottling on the chin (Jennings and Hayes 1994a; Stebbins 2003). A cream-colored supralabial stripe is present that fades anteriorly in front of the eye (Platz and Frost 1984).

In California, this frog is most likely to be confused with the closely related, nonnative Rio Grande leopard frog (*R. berlandieri*). The distinguishing characters for the two species widely overlap, and positive identification is therefore

Lowland Leopard Frog: Risk Factors

Ranking Criteria (Maximum Score)	Score
i. Range size (10)	10
ii. Distribution trend (25)	20
iii. Population concentration/ migration (10)	10
iv. Endemism (10)	0
v. Ecological tolerance (10)	3
vi. Population trend (25)	Data deficient
vii. Vulnerability to climate change (10)	10
viii. Projected impacts (10)	10
Total Score	63
Total Possible	85
Total Score/Total Possible	0.74




PHOTO ON PREVIOUS PAGE: Lowland leopard frog, Cochise County, Arizona. Courtesy of Brian Freiermuth.

difficult. *Rana berlandieri* attains larger body sizes (up to 11.4 cm SVL) and has proportionately larger eyes than *R. yavapaiensis*. Coloration of the two species is similar, but *R. yavapaiensis* generally has more extensive reticulation between the blotches on the hind legs, and its ventral coloration is often less dusky than *R. berlandieri* (Stebbins 2003). *Rana berlandieri*'s call consists of a low trill often followed by grunts, whereas *R. yavapaiensis* calls with higher-pitched notes that are given in rapid succession, often followed by lower-pitched chucks (Stebbins 2003, Elliott et al. 2009). Given that there are no known extant *R. yavapaiensis* localities remaining in California and that it is similar in appearance to the nonnative species *R. berlandieri*, positive identifications should be made cautiously. The species are readily distinguishable using molecular data (Hillis and Wilcox 2005, Frost et al. 2006a), which should be used to confirm any potential *R. yavapaiensis* specimens from California.

Taxonomic Relationships

Rana yavapaiensis was recognized as a distinct species in the leopard frog complex primarily on the basis of morphology, reproductive isolation, and allozyme variation (Platz and Platz 1973, Platz 1976, Platz and Frost 1984). The species is morphologically similar to other species of leopard frogs in the southwest. Jaeger et al. (2001) distinguished relict leopard frogs (*R. onca*) from *R. yavapaiensis* using genetic and morphological data. Based on a mitochondrial DNA dataset, Hillis and Wilcox (2005) confirmed a close relationship between these two species to the exclusion of other leopard frog taxa, including several geographically nearby members of the complex.

Frost et al. (2006a) recommended placing this species and many other North American ranids in the genus *Lithabates*, although this proposal and the analyses that support it are controversial (Crother 2009, Frost et al. 2009a, Pauly et al. 2009). We retain the traditional taxonomy here to maintain stability and pending further analyses.

Life History

Life history characteristics of California populations of *Rana yavapaiensis* are poorly known. The species apparently breeds opportunistically during winter rains (Stebbins 1972), and breeding has been documented to occur from late December through March in California (Storer 1925, Ruibal 1959). Elsewhere in the range, breeding has been documented from October to April (Platz and Platz 1973, Collins and Lewis 1979, Frost and Platz 1983, Sartorius and Rosen 2000). The reproductive biology of *R. yavapaiensis* has only been studied in Arizona. There, the species is known to experience at least two reproductive peaks within a year (once in the fall, once in the winter or spring), and tadpoles may overwinter (Collins and Lewis 1979, Sartorius and Rosen 2000). However, some authors have observed among-population variation in the occurrence of multiple breeding peaks, and it is unknown whether California populations had one or two breeding peaks per year.

Rana yavapaiensis undergoes marked year-to-year fluctuations in population size throughout its range (Clarkson and Rorabaugh 1989, Sredl et al. 1997, Sartorius and Rosen 2000), which renders isolated populations susceptible to extirpation. This also makes it difficult to confirm the absence or extirpation of populations with single-year surveys, emphasizing the importance of multiyear surveys for this species.

Habitat Requirements

Habitat requirements for *Rana yavapaiensis* are poorly understood, particularly in California. The species was historically found in slow-moving water along the San Felipe Creek drainage and the Lower Colorado River (Storer 1925, Stebbins 1972). The species has been found predominantly in marshy areas with bulrushes, cattails, and grasses with a willow overstory (Storer 1925, Jennings and Hayes 1994a, Jennings and Hayes 1994b), but it is unknown whether this vegetation type is required for population persistence. The species also expanded into artificial canals

and ditches in the Imperial Valley as agriculture developed in the region (Storer 1925, Klauber 1934), as is the case currently for *R. berlandieri* in Imperial County. It is unknown whether *R. yavapaiensis* can persist in these artificial habitats or whether they represent non-sustaining sink habitat requiring immigrants from nearby source populations.

Aquatic dissolved salt levels probably limit the distribution of this species, at least in some situations. Ruibal (1959) examined salt tolerance in adults and eggs from the San Felipe Creek drainage and found that salinities observed throughout most of the drainage were lethal to eggs (though not to adults) and that suitable areas for breeding were limited to the springs and seeps that fed the drainage. Whether salt concentration was always a limiting factor in California, or agricultural practices led to unnaturally high salt levels in some water bodies, is unknown.

Distribution (Past and Present)

No extant populations are presently known in California (Jennings and Fuller 2004). The distribution of *Rana yavapaiensis* was historically patchy, even before recent declines. In California, the species was historically present in suitable habitat along the Lower Colorado River, the Imperial Valley, and the San Felipe Creek drainage (Platz 1988, Stebbins 2003). Outside of California, the species historically ranged along the Lower Colorado River from northern Mexico to Arizona, from near sea level to 1700 m (Platz and Frost 1984, Platz 1988, Jennings and Hayes 1994a, Jennings and Hayes 1994b, Stebbins 2003). The last confirmed record in California is from 1965 in an irrigation ditch east of Calexico, Imperial County (Jennings and Hayes 1994a).

Trends in Abundance

Severe declines have occurred throughout the known California range of *Rana yavapaiensis*, and currently there are no known extant populations. Repeated surveys since 1965 have failed to locate this species (Vitt and Ohmart 1978,

Clarkson and Rorabaugh 1989, Jennings and Hayes 1994b). In addition, in 1976 Hurricane Kathleen apparently modified the surface drainage patterns around San Sebastian Marsh, Imperial County, eliminating the wetland habitat that supported the species previously (E. Ervin, pers. comm.). *Rana yavapaiensis* also appears to be declining through parts of its range outside of California (Clarkson and Rorabaugh 1989, Stebbins 2003).

Nature and Degree of Threat

The declines in *Rana yavapaiensis* occurred before extensive collections were made or studies were carried out. As a consequence, threats to this species in California are poorly understood, with few actual data supporting any of the potential threats considered here. Possible threats that contributed to its decline include direct impacts from agricultural runoff, which has been shown to be highly detrimental to other species in the leopard frog complex (Relyea 2008), habitat alteration, including water availability and/or flow regimes (Hayes and Jennings 1986), and predation by or competition with introduced bullfrogs, predaceous fishes, and invertebrates (Clarkson and Rorabaugh 1989). Some recent declines in the closely related *R. onca* appear to be linked to encroachment of dense emergent vegetation into open water habitats (Bradford et al. 2004), and this process could plausibly also affect *R. yavapaiensis*. All of these factors were occurring simultaneously within the range of *R. yavapaiensis* along with declines, making it difficult to disentangle their effects (Hayes and Jennings 1986). In addition, over 13,000 km of ditches in the Imperial Valley were burned and subsequently sprayed with oil during this time, and this presumably adversely affected these frogs (Twining and Hensley 1943).

Chytridiomycosis has been documented as contributing to declines in *R. yavapaiensis* populations in Arizona (Bradley et al. 2002), and this disease is a concern for any remaining California populations. An additional concern is the possibility of competition or hybridiza-

tion with *R. berlandieri* in California. *Rana berlandieri* was introduced into California well after *R. yavapaiensis* declined (Platz et al. 1990), so it is presumably not involved in the initial decline of the species. However, as it continues to expand its range in southern California, *R. berlandieri* may pose a risk to any remaining *R. yavapaiensis* populations (Rorabaugh et al. 2002). Hybridization has been documented between other species pairs of the leopard frog complex, including rare natural hybridization between *R. yavapaiensis* and the Chiricahua leopard frog (*R. chiricahuensis*) (Platz and Frost 1984). Molecular phylogenetic analyses suggest that *R. berlandieri* is more closely related to *R. yavapaiensis* than to *R. chiricahuensis*, implying that natural hybridization between *R. berlandieri* and *R. yavapaiensis* may be possible. Because *R. berlandieri* is now far more common in California than *R. yavapaiensis*, ongoing hybridization, should it occur, may result in genetic swamping of any remaining populations.

Status Determination

Rana yavapaiensis has undergone severe declines and has not been documented in California in over 40 years, and there is a strong possibility that the species is already extirpated statewide. However, it remains possible that the frog is present in scattered isolated localities that have not been surveyed, or that frogs have gone undetected despite surveys.

If any populations persist, it is likely that they are vulnerable to the causes of initial decline throughout most of the California range of this species. Such populations, which are almost certainly small and/or isolated, would also be vulnerable to the natural fluctuations in population size that occur in this species. This natural vulnerability could be exacerbated by changing precipitation regimes in the southeastern part of California, where increasing temperatures, declines in precipitation, and greater year-to-year variation in rainfall are expected to occur due to climate change (Cayan et al. 2008b, PRBO 2011).

Management Recommendations

If new surveys locate remaining populations of this species, the habitat supporting these frogs should be protected while further study is carried out. Without a better understanding of this species' life history in California, establishing an effective management program will be difficult. If native California populations are not found, *Rana yavapaiensis* is a potential candidate for assisted reintroductions from nearby populations in Arizona, particularly in areas where introduced *R. berlandieri* are not present or have been eliminated. More generally, such future introductions should be attempted in habitats that are as pristine as possible, and are free of introduced anurans of any species, introduced predatory fishes (including mosquitofish), and pathogenic fungi.

Monitoring, Research, and Survey Needs

Survey efforts need to be renewed along the San Felipe Creek drainage, the Imperial Valley, and the Lower Colorado River. Although the most likely areas for remnant populations are those that have been the least impacted by agriculture and development, even degraded agricultural habitat can be utilized by *Rana yavapaiensis*, and therefore should be surveyed. Because populations are prone to large yearly fluctuations, surveys should be repeated over multiple years in both the wet and dry seasons. Surveys for larvae should also be undertaken since tadpoles are often more reliably detected than adults. If any remaining populations are located, the habitat surrounding these areas should be protected, and researchers should begin a monitoring program to quantify and track population sizes. Any suspected *R. yavapaiensis* populations should be confirmed using a set of molecular markers, both to firmly establish species identity and to check for hybridization between *R. yavapaiensis* and *R. berlandieri*. Because hybridization is a concern, both mitochondrial and nuclear markers should be used. Given the difficulty in distinguishing the two species, we recommend that populations of presumptive *R. berlandieri* be

sampled for genetic material using nonlethal means (such as toe clips) and checked for diagnostic molecular markers to confirm that no native *R. yavapaiensis* DNA is present.

Should surveys discover extant populations of *R. yavapaiensis*, research into the basic life

history and the causes of decline in California will be a prerequisite to developing an effective management program. Life history studies with a particular focus on habitat suitability should be undertaken on any populations that are located or reestablished.



COUCH'S SPADEFOOT

Scaphiopus couchii Baird 1854

Stotus Summary

Scaphiopus couchii is a Priority 3 Species of Special Concern, receiving a Total Score/Total Possible of 56% (62/110). During the previous evaluation, it was also considered a Species of Special Concern (Jennings and Hayes 1994a).

Identification

Scaphiopus couchii is a medium-sized (5.7–9.1 cm SVL) anuran with a black keratinized spade on the heel of each hind foot and a vertically elliptical pupil (Stebbins 2003). The dorsal coloration is variable, ranging from green or greenish-yellow to brownish-yellow with a pattern of darker markings forming lines, spots, or a reticulating network (Grismer 2002, Stebbins 2003). Males are generally greener and have less conspicuous dorsal patterning than females (Grismer 2002, Stebbins 2003). The ventral surface is whitish (Grismer 2002). The call is a short (~1 s) low groan that declines in pitch and has been described as sounding simi-

lar to the bleating of a sheep (Elliott et al. 2009). Within its range, *S. couchii* can be distinguished from all other frogs by the presence of a conspicuous black spade on the hind feet and a vertically oriented pupil. Specimens that

Couch's Spadefoot: Risk Factors

Ranking Criteria (Maximum Score)	Score
i. Range size (10)	10
ii. Distribution trend (25)	0
iii. Population concentration/ migration (10)	10
iv. Endemism (10)	0
v. Ecological tolerance (10)	10
vi. Population trend (25)	15
vii. Vulnerability to climate change (10)	10
viii. Projected impacts (10)	7
Total Score	62
Total Possible	110
Total Score/Total Possible	0.56




PHOTO ON PREVIOUS PAGE: Couch's spadefoot, Cochise County, Arizona. Courtesy of Rob Schell Photography.

have been unearthed from burrows, or have newly emerged, may be covered in a dark hardened layer of skin that soon sloughs off (Mayhew 1965).

Taxonomic Relationships

Scaphiopus couchii was initially described based on morphology, and its distinctiveness has not been questioned since this time (Baird 1854). Intraspecific variation has only been examined in a small portion of the range in Arizona and New Mexico (Chan and Zamudio 2009). Little genetic structure was observed among the populations studied.

Life History

Scaphiopus couchii is xeric-adapted and primarily fossorial, spending the majority of its life in self-constructed burrows and emerging only during and immediately after intense summer rains (Mayhew 1965, McClanahan 1967, Stebbins 2003). This species has been found to be active on the surface after summer monsoon rainstorms in August and September, but not after winter rainstorms in October, December, or January (Mayhew 1965), suggesting that the activity period is limited to the summer in California as it is throughout the rest of its range. However, California differs from the rest of the range in that it receives the majority of its rainfall during the winter, suggesting that little local adaptation has occurred in this species (Mayhew 1965). Outside of California, surface activity is also tied to rain events, although *S. couchii* has (rarely) been found active on the surface during periods of high humidity, even when no recent rainfall has occurred (Mayhew 1962).

Emergence behavior is elicited by the low-frequency sound of rain falling on the desert soil, not from the rain itself (Dimmitt and Ruibal 1980a). Frogs emerge from deep (20–90 cm) burrows on the first night following the first heavy summer rain (Shoemaker et al. 1969, Dimmitt and Ruibal 1980a). Most breeding for a season usually occurs on this first night of activity (Woodward 1982). Following

this, the species may forage intermittently for up to 2 months, although much of this time is also spent in shallow (2–10 cm) burrows, which the frogs dig to avoid desiccation (Dimmitt and Ruibal 1980a). One feeding event can likely provide enough energy to allow an individual to persist for at least one year (McClanahan 1967, Dimmitt and Ruibal 1980b). Females deposit their eggs in ephemeral pools that form following intense summer rains (Woodward 1982). The development rate of this species is remarkably fast, with eggs hatching in as little as one day and metamorphosis occurring within 8–10 days if sufficient food is available (Mayhew 1965, Newman 1989, Morey and Janes 1994). Tadpoles are tolerant of a wide range of water temperatures (up to 39–42.5°C) such as are frequently encountered within the breeding pools (Brown 1969). This species likely does not breed every year in California and may skip reproduction and remain underground in sub-optimal years (Mayhew 1962).

Habitat Requirements

Scaphiopus couchii requires soils that are soft enough to allow burrowing. The species appears to prefer areas that contain at least some vegetation, although burrowing in completely open areas is also known (Mayhew 1965, McClanahan 1967). This taxon also requires the presence of temporary desert rain pools that retain water for at least 8 days to allow sufficient time for metamorphosis. The area in California in which *S. couchii* occurs receives an average of about 6.5 cm of rainfall per year, and its fine-scaled distribution may be linked to the amount of runoff that collects in localized areas (Mayhew 1965). The distances traveled between upland retreats and breeding sites are not known, nor are the precise terrestrial habitat requirements of adults or juveniles.

Distribution (Past and Present)

Scaphiopus couchii ranges throughout much of Texas, Mexico, southern Arizona, and southern New Mexico, from near sea level to 1800 m (Stebbins 2003). Scattered, localized

populations also occur in central Arizona and southern Colorado (Stebbins 2003). In California, this species ranges from the Colorado River west at least to the vicinity of the Algodones Dunes, ranging as far north as Chemehuevi Wash (~9 km north of Vidal Junction) and south to the vicinity of the United States–Mexico Border (Mayhew 1962, Tinkham 1962, Jennings and Hayes 1994a). A few observational records exist in the vicinity of the Salton Sea, and although these appear to be credible, verification is needed that populations are extant in this area.

This taxon's range in California is likely relictual from more mesic periods and is probably more fragmented now than it once was (Mayhew 1965). This species was not known to occur in the state until 1962 (Mayhew 1962, Tinkham 1962), and no significant declines have been documented since that time.

Trends in Abundance

No historical or current abundance data are available for this taxon within California. Human activities have both created and destroyed breeding sites for the species (S. Morey, pers. comm.), but no quantitative studies have documented the overall impacts of these activities on the species across California.

Nature and Degree of Threat

Scaphiopus couchii is likely persisting closer to its physiological limits in California than it is elsewhere within its range (Mayhew 1965). The California range is both hotter and drier than most of the rest of the range, and most of the limited rainfall occurs outside of the monsoon, during a time when *S. couchii* is usually inactive. The current populations in California likely persist due to the presence of local conditions that allow for the collection of sufficient quantities of water, such as the presence of basins on the eastern base of the Algodones Dunes and pools that form along desert washes. The relatively fragmented nature of the species' California distribution and the physiological

conditions under which it lives make it susceptible to localized extirpations due to habitat modification that destroys temporary pools and due to the effects of climate change. Recent models (PRBO 2011) indicate that average temperature will increase significantly, by more than 2°C in most months in the Sonoran/Colorado Desert of California. Given that *S. couchii* may already be near its physiological temperature limits, this may have an enormous impact on its viability in the state. In addition, some precipitation projections include an overall decrease of up to 45% (PRBO 2011), and increased variation in year-to-year precipitation (Cayan et al. 2008b), which could have severe detrimental impacts on this species by decreasing the number of years in which enough rainwater collects to allow breeding. Essentially, if the interpretation is correct that the California population exists at the physiological limits of the species' capacity, then predicted changes in rainfall and temperature may seriously reduce its range in the state.

Off-highway vehicle usage in the Algodones Dunes has degraded habitat in many areas (R. Fisher, pers. comm.). Noise generated by off-highway vehicle usage has been implicated in eliciting emergence in this species by mimicking the sound of falling rain that it uses as an emergence cue (Brattstrom and Bondello 1979). Temporary and permanent anthropogenic water sources associated with livestock (cattle ponds) and perhaps agriculture may help to provide suitable breeding habitat that is important to the persistence of this species.

Status Determination

The small and fragmented range of this taxon, coupled with its sensitivity to habitat disturbance through off-highway vehicle use and predicted climate change, justifies its Priority 3 status.

Management Recommendations

The primary, immediate management goal for *Scaphiopus couchii* is to protect existing habitat from further impact. Off-highway vehicle use

and larger modifications (solar projects, mining) may negatively alter both the hydrology of breeding pools and the suitability of soil for burrowing. In particular, if pools are modified such that they dry faster (through either more rapid draining or overall smaller size), their hydroperiod may become too short to allow metamorphosis. Specific areas requiring protection should be determined by the surveys outlined below. In the future, the impacts of projected climate change may seriously threaten this species in California, and proactive management may be required to counteract this threat; such management could include relocating populations to cooler or more mesic sites, deepening and maintaining the hydroperiod of natural breeding sites, and potentially creating completely novel breeding pools that can hold water if the climate changes.

Monitoring, Research, and Survey Needs

Range-wide surveys need to be undertaken for this taxon to identify suitable remaining habitat, determine the sizes of extant breeding populations, and to further characterize the species' range in California. To our knowledge, the northernmost population at Chemehuevi Wash has not been resurveyed since its original description in 1962 (R. Fisher, pers. comm.), and this is an important area in need of surveys. As the species distribution in California is patchy, largely in remote regions of the state, and given that the species does not emerge every year, care should be taken to search desert pool habitats even in areas where this anuran has not yet been documented. Surveys should ideally take place during the first night following the first major summer (monsoonal) rain event. Surveyors should be experienced with this frog's call (Elliott et al. 2009), as this will likely be the easiest way to find populations, and pools should be surveyed for tadpoles

within a few days after they fill during summer rains.

The movement ecology of this taxon and its potential to recolonize previously extirpated areas are unknown and are a topic in need of further study, particularly so in California where populations appear to be fragmented. Additional study of its physiological limits would also be helpful in establishing a more informed management plan, now and in the face of future climate changes. In particular, the severity of drought and the number of years between breeding events that can be tolerated are critical pieces of information for the long-term management of this species. Landscape ecological information, including the amount of terrestrial habitat needed, the relationship between population size and pool basin size, inundation duration and frequency, and the movement frequency of animals between breeding sites would all be valuable for future management considerations. Additional information on habitat use itself, including the extent that ongoing railroad and water diversion projects within the range subsidize or detract from potential habitat for this species, is also a critical research need that would inform ongoing management of this species.

Finally, given the spotty distribution of the species and the potential for genetic isolation among sites, multi-locus population genetic studies using microsatellites or single nucleotide polymorphisms of all extant California populations would provide a badly needed estimate of the extent to which populations are subdivided and therefore the optimal management strategies to protect genetic diversity. In addition, given how widespread the species is across the southwestern United States, genetic data comparing the uniqueness of the California population is essential for range-wide management.



WESTERN SPADEFOOT

Spea hammondi (Baird 1859)

Status Summary

Spea hammondi is a Priority 1 Species of Special Concern, receiving a Total Score/Total Possible of 69% (76/110). During the previous evaluation, it was also considered a Species of Special Concern under the name *Scaphiopus hammondi* (Jennings and Hayes 1994a).

Identification

Spadefoot toads as a group have catlike eyes with vertical pupils, a single black spade on each hind foot, and indistinct paratoid glands (Stebbins 2003). *Spea hammondi* is dusky green or gray dorsally, often with irregular markings (Stebbins 2003). Tubercles on the skin are tipped with orange or red, and the irises are usually pale gold (Jennings and Hayes 1994a, Stebbins 2003). The ventral surface is white to light gray without markings (Stebbins 2003). Adults are 4–6 cm SVL (Stebbins 2003). Larvae can reach approximately 7 cm in TL and their eyes are set close together when viewed from above (Stebbins 2003). This species is unlikely to be confused with other sympatric anurans.

Taxonomic Relationships

North American spadefoots have had a confusing taxonomic history. Studies using allozymes and morphology (Wiens and Titus 1991) and mitochondrial DNA (Garcia-Paris et al. 2003) support the species status of *Spea hammondi*.

Western Spadefoot: Risk Factors

Ranking Criteria (Maximum Score)	Score
i. Range size (10)	5
ii. Distribution trend (25)	20
iii. Population concentration/ migration (10)	10
iv. Endemism (10)	7
v. Ecological tolerance (10)	10
vi. Population trend (25)	10
vii. Vulnerability to climate change (10)	7
viii. Projected impacts (10)	7
Total Score	76
Total Possible	110
Total Score/Total Possible	0.69




PHOTO ON PREVIOUS PAGE: Western spadefoot, Sacramento County, California. Courtesy of Robert Thomson.

placing it sister to a clade consisting of the Great Basin spadefoot (*S. intermontana*) and the Plains spadefoot (*S. bombifrons*). This arrangement is consistent with Kluge (1966) and Sattler (1980). Relationships within *Spea* are still unresolved however, and cryptic taxa may exist within *S. hammondi* (Garcia-Paris et al. 2003) and *S. intermontana* (Wiens and Titus 1991). Ongoing phylogeographic work should clarify the extent of intraspecific variation in the species. Preliminary data indicate that some mitochondrial introgression has occurred between *S. intermontana* and *S. hammondi* in southern California, but not the Central Valley portions of the species' range (P. Spinks, unpublished data).

Life History

Adult *Spea hammondi* are terrestrial, moving from summer refugia to ephemeral water bodies to breed in the spring following warm late winter or spring rains (Storer 1925, Burgess 1950, Stebbins 1954, Feaver 1971, Brown 1976, Morey 1998). Breeding aggregations can consist of more than 1000 individuals (Jennings and Hayes 1994a). Breeding occurs over a 2–3 week period, during which males can be heard chorusing intermittently (Brown 1976, Morey and Reznick 2004). Additional bouts of breeding can occur, and pools can contain cohorts of different ages (Morey 2005). Onset of breeding activity varies depending on rainfall and region. For example, heavy rains in 1991 resulted in breeding occurring only in March (San Luis Obispo and Riverside Counties; Morey and Reznick 2004). In the two following years, breeding occurred between January and March (Morey and Reznick 2004). Breeding has also been documented in August, and from October to December in San Diego County (Ervin et al. 2005, Ervin and Cass 2007). It is unknown how common early-breeding behavior is, but the October 2004 events may have been in response to very dry conditions, and many of these larvae ultimately succumbed to desiccation. The previous year, 2003, set a record rainless period, and the breeding in 2004 occurred

after the first measurable rain in 181 days (Ervin et al. 2005).

Females lay 300–500 eggs in clusters of 18–25 (Stebbins 1951, Stebbins 1985) that usually hatch in 3–4 days (Morey 2005). Morey and Reznick (2004) surveyed vernal pools in San Luis Obispo and Riverside Counties and found that the larval period lasted an average of 58 days. In the laboratory, the minimum time for larval development was estimated to be 14 days (Morey and Reznick 2004). Males raised experimentally under high food conditions developed secondary sexual characters by the beginning of their first breeding season after metamorphosis, while females of the same age had adult coloration but underdeveloped ovaries (Morey and Reznick 2001). It is unknown how long it takes to reach maturity in the field, but based on this experimental work males probably mature 1–2 years after metamorphosis and females at least 2 years after metamorphosis. Most individuals are mature at 4–4.5 cm SVL (Storer 1925, Morey and Guinn 1992).

Larvae are frequently at risk of desiccation due to pools drying before development is complete. In Fresno County, 17 out of 23 vernal pools dried before larvae metamorphosed (Feaver 1971). Across 20 populations in San Luis Obispo and Riverside Counties, Morey and Reznick (2004) observed that 15% of ponds dried before larvae metamorphosed. As pools dry, larvae experience increased daily variation in temperature, increased ammonia levels, increased water hardness, and decreased depth (Morey and Reznick 2004). These factors lead to crowding and decreased growth rate. While several cues are operating simultaneously, water reduction alone is sufficient to trigger accelerated development within 24 hours (Denver 1997a, Denver 1997b, Denver et al. 1998, Boorse and Denver 2003). In the field, there is a positive correlation between hydroperiod and mass at metamorphosis (Morey and Reznick 2004). In the lab, animals reared at low density were larger (4.96 g) at metamorphosis and metamorphosed sooner (77.8 days) than animals maintained at high density (2.9 g, 87.8 days; Morey and Reznick

2001). Survivorship of metamorphs was also higher for animals that were larger at metamorphosis, regardless of larval density (Morey and Reznick 2001). Effects of the larval rearing environment persisted for several months after metamorphosis, but small metamorphs were able to catch up in growth if terrestrial food availability was high.

Little is known about terrestrial activity, although most movement and surface activity is thought to be nocturnal (Morey 2005). Juveniles leave natal pools shortly after metamorphosis in April–June presumably seeking refugia, although their terrestrial habitat is unknown (Morey 2005). Adults and juveniles retreat to burrows by late summer, with juveniles capable of digging burrows 10–20 cm deep even in hard, dry soil (Morey and Reznick 2001). Mammal burrows may also be used (Stebbins 1951).

Larval diet has not been studied, although larvae of other spadefoot species are generalists, consuming animals, plants, and organic detritus (Pomeroy 1981, Pfennig 1990). Cannibal morph larvae with broad heads and enlarged jaw muscles are known from San Luis Obispo and Riverside counties, but it is unknown how common they are throughout the species' range (Morey 2005). Adults are generalized predators on terrestrial arthropods and other prey, including beetles, moths, flies, and earthworms (Morey and Guinn 1992).

Habitat Requirements

Spea hammondi occurs in grasslands, oak woodlands, coastal sage scrub, and chaparral vegetation in washes, floodplains, alluvial fans, playas, and alkali flats (Stebbins 2003, Morey 2005). Temporary pools are used for breeding, but *S. hammondi* will also readily breed in artificial water bodies such as cattle ponds (Morey 2005). Vernal pools used by *S. hammondi* for breeding had an average ponding duration of 81 days (range 36–127, $n = 9$, San Luis Obispo and Riverside Counties) (Morey and Reznick 2004). Pools with at least some successful recruitment lasted on average 3 weeks longer than larval development time (Morey and Reznick 2004).

Pool temperature during larval development ranged from 11°C to 32°C (Morey and Reznick 2004). Brown (1967) found that water temperatures between 9°C and 30°C were necessary for larval development (eggs collected from Riverside County).

Perennial pools containing introduced predators such as crayfish, fish, or bullfrogs are often unsuitable for successful recruitment (Jennings and Hayes 1994a). However, in southern California, ephemeral pools utilized by introduced species with predatory aquatic stages, such as the African clawed frog (*Xenopus laevis*), can still function as breeding habitat for *S. hammondi* (confirmed by the presence of dispersing metamorphs), but the effects these introduced species have on overall recruitment levels are unknown (Ervin and Fisher 2001, Ervin and Burkhardt 2006).

Distribution (Past and Present)

Spea hammondi occurs in the Central Valley and bordering foothills across southern California from Shasta County south into northwestern Baja California, including the Coast Ranges south of Monterey, from sea level to 1365 m (Jennings and Hayes 1994a, Ervin et al. 2001, Stebbins 2003; S. Barry, pers. comm.). Jennings and Hayes (1994a) concluded that as of the 1990s, over 80% of historically occupied habitat in southern California and 30% of habitat in northern California were no longer suitable due to development and habitat conversion. In surveys throughout the Central Valley, Fisher and Shaffer (1996) reported *S. hammondi* as virtually extirpated from the Sacramento Valley and at a reduced density in populations of the eastern San Joaquin Valley.

Trends in Abundance

Current or historical abundance data are largely unavailable or anecdotal, and little recent data is available. Recent surveys of Mather Airport (formerly Mather Air Force Base) in Sacramento County estimated that breeding adults numbered in the few dozens, although this was based on short-duration surveys and limited

data (A. Chang, unpublished data). Morey and Guinn (1992) reported an average of 1.16 individuals/km of roadway during a relatively wet winter (1982–1983) and 0.68 individuals/km during a drier winter (1984–1985) in the San Joaquin Valley.

Nature and Degree of Threat

The major threat to *Spea hammondi* is habitat loss and fragmentation due to agriculture and urban development. Other threats include invasive species and climate change. Davidson et al. (2002) found that currently occupied sites had less surrounding urban development than extirpated sites. Extant populations also occur at higher elevations than extirpated sites on average, possibly due to invasive species being more common at lower elevation (Fisher and Shaffer 1996, Davidson et al. 2002). *Spea hammondi* is sensitive to invasive species such as crayfish, bullfrogs, and mosquitofish; however, many of these species cannot persist in the highly ephemeral breeding habitats *S. hammondi* uses (Jennings and Hayes 1994a, Morey 2005). Bullfrogs have been documented to prey on *S. hammondi* (Morey and Guinn 1992, Balfour and Ranlet 2006), although the impact of this predation on overall abundance is unknown.

Spea hammondi may be at risk from climate change because breeding is dependent upon temperature and rainfall cues, and larval development requires ephemeral pools to persist long enough to complete development (Morey and Guinn 1992, Jennings and Hayes 1994a). Mean annual temperatures are projected to increase throughout the range of *S. hammondi*, with warmer winters and summers and earlier spring warming expected (reviewed in PRBO 2011). The frequency of extremely hot days is predicted to increase by up to 25 days per year in some parts of the range (Bell et al. 2004). There is less certainty about future precipitation patterns, with estimates ranging from little change to roughly 30% decreases in rainfall (Snyder and Sloan 2005, PRBO 2011). Changes in temperature and precipitation will likely affect vernal pool hydrology (e.g., Pyke 2005) and may

also affect the timing of breeding, though how *S. hammondi* will respond to these changes needs further study. The largely unsuccessful early breeding observed by Ervin et al. (2005) may be indicative of the kinds of mismatches in environmental cues and breeding behavior that this species may suffer under climate change. The probability of large (>200 ha) wildfires is expected to change very little in the Central Valley (Westerling and Bryant 2008). In the more northern coastal part of the range, the probability of large fires is expected to increase (Westerling and Bryant 2008), and the area burned is expected to increase by up to 50% (Lenihan et al. 2008). In the southern part of the range where wildfire is common, there is little consensus on future fire dynamics because of the difficulty in modeling Santa Ana weather events (Westerling et al. 2004, Westerling and Bryant 2008). The largely subterranean lifestyle of *S. hammondi* may make it relatively resistant to the effects of fire. However, wildfires occurring during dispersal may be particularly detrimental due to direct mortality and habitat degradation and this issue requires more study. Vegetation shifts due to climate change are expected to be modest in the Central Valley, where land use is a more important determinant of habitat type (PRBO 2011). Elsewhere in the range, chaparral and shrublands are expected to decrease in area, while grassland is expected to increase (Lenihan et al. 2008, PRBO 2011). The impact of these shifts may be modest as *S. hammondi* uses all of these habitat types.

Status Determination

Ongoing habitat loss and extirpations throughout the range of *Spea hammondi* warrant Priority 1 Species of Special Concern status.

Management Recommendations

Remaining sites should be protected from urban and agricultural development, with emphasis on larger habitat blocks that allow for more natural metapopulation dynamics to persist. The fact that *Spea hammondi* readily breeds in anthropogenic structures can be exploited to

create breeding habitat in response to habitat loss and potentially also to climate changes that affect natural vernal pool phenology. Terrestrial habitat is likely not so easily restored, and minimizing or eliminating disturbance around breeding habitat would help protect adults (see the "Monitoring, Research, and Survey Needs" section). Efforts to remove introduced predators from breeding habitat should be considered. In some cases, cattle grazing operations may be beneficial to *S. hammondi*. Over 3 years in Sacramento County, Marty (2005) found that experimentally grazed vernal pools experienced fewer drying and refilling cycles within a season, and had a longer maximum inundation period (115 days) than ungrazed treatments (65 days) or treatments where grazing occurred seasonally (65–78 days).

Monitoring, Research, and Survey Needs

Research is needed into terrestrial habitat use (Jennings and Hayes 1994a, Morey 2005), including juvenile dispersal, adult migration patterns and distances, and the importance (if any) of rodent burrows for all age classes. This

information is important for determining how much and what kinds of terrestrial habitat to protect around breeding sites. For example, Morey and Reznick (2001) found that the quality of juvenile terrestrial habitat in terms of food availability compensated for stressful larval conditions. Additional study on which environmental conditions promote post-metamorphic survival will aid in management planning. It is also unknown what proportion of adults breed each year and how long individual adults spend at breeding sites (Morey 2005). Underground habitat use is poorly known, including feeding and dormancy patterns. Remaining populations are likely highly fragmented, and research is needed into connectivity among populations at both the local and the regional levels; additional landscape ecology and genetic studies would help determine patterns of differentiation (Jennings and Hayes 1994a). Finally, comparative studies of this species in the Central Valley and southern California would help determine the extent of biological variation in life history patterns across this ecological gradient.



SOUTHERN LONG-TOED SALAMANDER

Ambystoma macrodactylum sigillatum Ferguson 1961

Status Summary

Ambystoma macrodactylum sigillatum is a Priority 2 Species of Special Concern, receiving a Total Score/Total Possible of 66% (73/110). It was not considered a Species of Special Concern during the previous evaluation (Jennings and Hayes 1994a).

Identification

Ambystoma macrodactylum sigillatum is a medium-sized (4.1–8.9 cm SVL) salamander with a broad head and large eyes (Stebbins 2003). The dorsal ground coloration is black or dusky brown with a yellow dorsal stripe that is usually divided into blotches on the body and into fine spotting on the head and tail (Ferguson 1961, Petranka 1998, Stebbins 2003). Small whitish-blue flecks are present on the sides of the body, and the ventral surface is dark brown (Stebbins 2003). The larvae have large bushy gills and a dorsal fin that extends to near the forelimbs (Petranka 1998).

Metamorphosed individuals of this species are unlikely to be confused with any other salamanders within its range. Other subspecies of *A. macrodactylum* have similar body proportions but differ in the size, extent of blotching,

Southern Long-Toed Salamander: Risk Factors

Ranking Criteria (Maximum Score)	Score
i. Range size (10)	5
ii. Distribution trend (25)	15
iii. Population concentration/migration (10)	10
iv. Endemism (10)	3
v. Ecological tolerance (10)	3
vi. Population trend (25)	20
vii. Vulnerability to climate change (10)	10
viii. Projected impacts (10)	7
Total Score	73
Total Possible	110
Total Score/Total Possible	0.66




PHOTO ON PREVIOUS PAGE: Southern long-toed salamander, Butte County, California. Courtesy of Robert Hansen.

and coloration of the dorsal stripe, and their ranges do not overlap in California. Differentiating larvae from co-occurring newts (*Taricha granulosa*, *T. torosa*) requires careful attention. Newt larvae generally have small, narrow heads and few gill rakers (5–7 on the anterior side of the third gill arch), whereas *A. macrodactylum* larvae have broad heads and 9–13 gill rakers on the anterior side of the third arch (Stebbins 2003).

Taxonomic Relationships

Ambystoma macrodactylum sigillatum is one of five currently recognized subspecies of long-toed salamander (Petranka 1998, Stebbins 2003). *Ambystoma macrodactylum* has been widely recognized as a distinct species since its initial description by Baird (1854). Since this time, a number of different species and subspecies have been described. The current five-subspecies arrangement stabilized after the work of Ferguson (1961), which described *A. m. columbianum* (eastern long-toed salamander) and *A. m. sigillatum*, as well as the work of Russell and Anderson (1956), which described the geographically isolated *A. m. croceum* (Santa Cruz long-toed salamander) from Santa Cruz and Monterey Counties. Ongoing genetic studies indicate that several of these subspecies may warrant full species status (Savage 2008). *Ambystoma macrodactylum sigillatum* was described based on the size, color, and pattern of the dorsal band, as well as vomerine tooth counts (Ferguson 1961). Although it intergrades morphologically with *A. m. columbianum* at the northern edge of its range (Ferguson 1961), ongoing genetic analyses support recognition of *A. m. sigillatum* as a distinct species (Savage 2008).

Life History

Ambystoma macrodactylum sigillatum is a pond-breeding salamander that often has a prolonged larval stage. The life history of this taxon varies widely depending on elevation and climate (Petranka 1998). Here we have summarized data for *A. m. sigillatum*, where possible, and

described the variation present across the species where the life history is highly variable and/or uncertain.

Adults emerge from hibernation and migrate to breeding habitat after the first thaw. Mating begins shortly after adults enter the breeding habitat, usually in May or June, with lower-elevation populations usually being able to breed earlier than higher-elevation populations (Anderson 1967, Howard and Wallace 1985). Elsewhere in the *A. macrodactylum* range, primarily at low elevations where the climate is mild, breeding is not delayed by winter freezes, so reproduction starts with the onset of fall rains (Ferguson 1961, Nussbaum et al. 1983). As in other *Ambystoma* species, mating follows a pattern of courtship and spermatophore deposition. Females oviposit on vegetation, rocks, sticks, or directly on the pond bottom 2–3 days following courtship and mating (Anderson 1961, Stebbins 2003). The eggs are laid singly or in clumps of up to 100 eggs (Petranka 1998, Stebbins 2003). The pattern of egg deposition varies geographically in this species: *A. m. sigillatum* tends to lay eggs singly or in long loose clusters in relatively deep water (Anderson 1967), although this is variable. Eggs hatch in 2–5 weeks, with longer incubation periods required at higher elevations and lower water temperatures (Anderson 1967, Nussbaum et al. 1983, Petranka 1998). The larval period can be as short as 50 days in temporary pools at lower elevations but may last 2 years in the highest elevations in permanent pools (Nussbaum et al. 1983, Pilliod and Fronzuto 2005). Size at metamorphosis varies widely from 2.3 to 4.8 cm SVL (Howard and Wallace 1985). This species is able to tolerate a relatively wide range of water temperatures, with larvae overwintering under the ice at near freezing temperatures but then selecting the warmest areas available throughout the summer (up to 24.5°C). Presumably these temperatures allow for more rapid larval growth and development (Anderson 1968b).

Ambystoma macrodactylum sigillatum is a generalist predator, as both larva and post-

metamorph, that feeds on a variety of small insects, crustaceans, and spiders (Anderson 1968a). Larvae and males in the aquatic environment will prey on zooplankton, insect larvae, and small snails (Anderson 1968a, Nussbaum et al. 1983). In the lab, larvae are also known to take frog (primarily *Pseudacris*) tadpoles and conspecific larva (Anderson 1968a, Nussbaum et al. 1983). Females apparently do not feed in the aquatic environment, which may simply reflect the short amount of time they spend there during the breeding season (Anderson 1968a).

Habitat Requirements

Ambystoma macrodactylum, as a species, occurs in a larger variety of habitat types than any other salamander in the Northwestern United States (Ferguson 1961, Nussbaum et al. 1983). Suitable habitats for *A. m. sigillatum* include arid grassland and sagebrush communities, dry woodlands, coniferous forests, alpine meadows, and a wide variety of intermediate habitat types (Ferguson 1961, Petranka 1998, Pilliod and Fronzuto 2005). In some areas, this species is abundant in disturbed agricultural areas (Nussbaum et al. 1983). Elsewhere in the range, landscape genetic studies indicate that populations that persist in highly modified habitats do so with increased population isolation, probably increasing susceptibility to local extirpations (Goldberg and Waits 2010).

At high elevations (above 2450 m in the Sierra Nevada and 2100 m in the Klamath Mountains), where breeding occurs late and larval development is prolonged, some populations of *A. m. sigillatum* require permanent water bodies for breeding because larvae overwinter prior to metamorphosis (Anderson 1967; K. Leyse, pers. comm.). If these overwintering sites are shallow (1–2 m in depth), as is common in the Tahoe region of the Sierra Nevada, few larvae seem to survive the winter (K. Leyse, pers. comm., unpublished data). Spring-fed water bodies may increase the likelihood of successful overwintering, though more data are required to verify this. This subspecies also per-

sists far more readily in fishless water bodies (see the “Nature and Degree of Threat” section).

The species is known to utilize hardwood forests, meadows, and granite slopes for upland habitat. Further study on the extent and types of upland habitat that this species requires are needed.

Distribution (Past and Present)

Ambystoma macrodactylum sigillatum ranges from southwestern Oregon (south of the Calapooya divide, Lane and Douglas Counties) through the Trinity Alps, Warner Mountains, Sierra Nevada, and adjacent areas of northwestern California reaching as far south as Carson Pass (Ferguson 1961, Brode 1967, Bury 1970a, Pilliod and Fronzuto 2005). The known elevational range for this taxon is from near sea level to 3000 m (Stebbins 1966, Nussbaum et al. 1983), although the distribution in California is restricted to the higher end of this range. The presence of isolated populations of the species *A. macrodactylum* in Santa Cruz and Monterey Counties, California (*A. m. croceum*), and in southeastern Oregon suggests that the species may have been historically distributed more broadly throughout the west. If so, the present-day range likely reflects a range contraction as climate has changed over the last several thousand years.

Localized, present-day changes in distribution appear to be ongoing in several parts of California. In the historically fishless Klamath–Siskiyou bioregion, *A. m. sigillatum* are 44 times more likely to be present in lakes without fish than lakes that contain fish. Because these fish have been introduced during the last 150 years, it is likely that some lakes where *A. m. sigillatum* does not occur represent localized extirpations as a result of fish predation (Welsh et al. 2006). A similar pattern occurs in the north central Sierra Nevada near Lake Tahoe. Here, *A. m. sigillatum* are present in 92.3% of fishless sites, but only 37.5% of fish-containing sites (Leyse 2005). In the Klamath Mountains, *A. m. sigillatum* was documented at 25 of 118

sites in surveys conducted between 1999 and 2001. Salamanders were present at only 15 of these sites when they were resurveyed in 2008 (K. Pope, pers. comm.). The overall geographic extent of the *A. m. sigillatum* range appears to still be intact, but it is clear that localized extirpations are occurring in several areas.

Trends in Abundance

Abundances of *Ambystoma macrodactylum sigillatum* have declined throughout relatively large areas of the California range. The Klamath Mountain surveys described above documented 4126 individuals at 25 occupied sites in 1999–2001 but only 569 individuals at the 15 occupied sites in 2008 (K. Pope, pers. comm.). Few historical abundance data are available, but overall current abundance of larvae at lower-elevation sites appears to be low (K. Leyse, pers. comm.). Population genetic estimates of population trends suggest that regional populations exchange few migrants and that effective population sizes are small (Savage et al. 2010).

Nature and Degree of Threat

Trout introductions are the largest threat to remaining populations of *Ambystoma macrodactylum sigillatum*. Welsh et al. (2006) found that the absence of introduced fish was a major predictor of *A. m. sigillatum* presence even after controlling for other environmental variables. Aside from the local effect of fish on individual water bodies, fish introductions appear to affect *A. macrodactylum* populations at the scale of entire watershed basin. In Idaho, basins with higher introduced fish densities had significantly lower densities of *A. macrodactylum* (Pilliod and Peterson 2001). The authors postulated that much of the remaining fishless habitat in fish-containing basins is too shallow for most larvae to successfully overwinter and that the deeper, fish-containing pools no longer acted as stable source populations for the basin. This led to a destabilization of normal source–sink dynamics, causing declines throughout the entire basin. These results suggest that the presence of fish at the

basin scale is a significant conservation risk, irrespective of whether patches of fishless habitat remain within the basin (Pilliod and Peterson 2001). Where *A. m. sigillatum* persist in the presence of fish, larval densities are very low both in deeper fish-containing pools and in adjacent fishless pools (K. Leyse, pers. comm.). When larvae are found in fish-containing pools, they tend to hide under rocks or are only captured in overnight trapping, indicating that they may alter their behavior in response to the presence of predators (K. Leyse, pers. comm., though see Tyler et al. 1998). Declines due to the presence of fish have also been documented elsewhere in *A. macrodactylum*'s range (Liss and Larson 1991, Liss et al. 1995, Tyler et al. 1998). In Montana, introduced trout were linked to *A. m. krausei* extirpations. Salamander recolonization following local trout extirpations strongly indicated that trout were the actual causal agent of declines (Funk and Dunlap 1999).

Climate change also poses a threat for *A. m. sigillatum*. Many of the remaining pools that this species utilizes are shallow. Projected shifts to earlier and faster snowmelt in the Sierra Nevada could have complex and possibly negative effects on this species by changing the hydrology of lakes and ponds (Cayan et al. 2008b, Franco et al. 2011, PRBO 2011). As many of these pools appear to be spring fed, any changes to hydrology of the springs could also have severe impacts (Leyse 2005).

Disease and environmental contaminants may also pose threats for remaining populations of *A. m. sigillatum*. Lethal ranavirus infections of *A. m. sigillatum* were recently detected in Lassen Volcanic National Park (Bunck et al. 2009). This species is also susceptible to iridovirus infection and exposure to atrazine, a commonly used herbicide (Forson and Storfer 2006). *Bd* has been detected in a single adult salamander at Carter Meadow in Lassen National Forest, although the load was low. Prevalence of *Bd* appears to be low for this species and no evidence of die-offs or illness due to this pathogen is known (K. Pope and J. Piovita-Scott, unpublished data).

Status Determination

Ongoing serious declines in distribution and abundance are the primary reasons for this Priority 2 status.

Management Recommendations

The presence of relatively deep fishless pools appears to be important to the continued persistence of this species, particularly at the highest elevations. As such, fish stocking should be limited in areas where *Ambystoma macrodactylum sigillatum* occurs. Where stocking does occur, mitigation strategies outlined by Appendix K of California Department of Fish and Wildlife hatchery and stocking program environmental impact report should be followed (ICF Jones and Stokes 2010).

Monitoring, Research, and Survey Needs

Declines due to fish predation have now been amply demonstrated, so continued monitoring on the effects of fish predation is less important than work related to fish removal. If predaceous fish can be successfully removed from areas supporting this species, occasional monitoring

should be undertaken to detect unauthorized reintroductions, particularly in areas that experience high human impact and to document recolonization dynamics by the salamanders. An important management question centers on the relative importance of permanent and temporary pools to metapopulation dynamics across elevations. That is, it may be that at lower elevations, temporary fish-free pools are the primary source of successful recruitment, and deeper lakes can therefore be maintained as fishing resources, whereas at the highest elevations, the species can only persist if permanent, fish-free habitats are common. The type and extent of upland habitat utilized by this species is also in need of further study. In particular, the extent of upland habitat that populations require in order to persist has not been studied in this taxon. Climate change could also have different impacts on the upland phase of the life cycle, in addition to the impacts that are projected for the aquatic part of the life cycle. In addition, populations are still under considerable risk from disease, and monitoring efforts focused on detecting the presence of ranavirus and *Bd* should be continued.



SANTA CRUZ BLACK SALAMANDER

Aneides flavipunctatus niger Myers and Maslin 1948

Status Summary

Aneides flavipunctatus niger is a Priority 3 Species of Special Concern, receiving a Total Score/Total Possible of 48% (53/110). This taxon was not previously considered a Species of Special Concern (Jennings and Hayes 1994a).

Identification

Aneides flavipunctatus niger is a medium-sized plethodontid salamander (5.1–9.5 cm SVL) (Stebbins 2003). The adult dorsal coloration is either solid black or black with a few small white flecks (Myers and Maslin 1948). Juveniles (<4.0 cm SVL) have brassy dorsal pigmentation with white to blue-white spots (Lynch 1981). The ventral coloration is black or dark gray (Myers and Maslin 1948). The nasolabial grooves and costal grooves are well defined, and most individuals (95%) have 17 costal grooves (Lynch 1981, Stebbins 2003). *Aneides flavipunctatus niger* has rounded toe tips, counter to the squared toe tips typical of *Aneides*. Its limbs are short relative to the trunk, with

3–5 costal grooves between adpressed limbs. The heads of males are larger than those of females, and are roughly triangular with prominent, protruding upper jaw teeth (Stebbins 2003).

Santa Cruz Black Salamander: Risk Factors

Ranking Criteria (Maximum Score)	Score
i. Range size (10)	10
ii. Distribution trend (25)	10
iii. Population concentration/migration (10)	0
iv. Endemism (10)	10
v. Ecological tolerance (10)	7
vi. Population trend (25)	10
vii. Vulnerability to climate change (10)	3
viii. Projected impacts (10)	3
Total Score	53
Total Possible	110
Total Score/Total Possible	0.48




PHOTO ON PREVIOUS PAGE: Santa Cruz black salamander, Santa Cruz County, California. Courtesy of William Flaxington.

Aneides flavipunctatus niger could be confused with the co-occurring arboreal salamander (*A. lugubris*). Adult *A. lugubris* are grayish to brownish above with yellow flecks that are often concentrated on the sides, squarish toe-tips, and a pale whitish venter (Stebbins 2003). Juvenile *A. f. niger* have green pigmentation, while *A. lugubris* juveniles do not.

Taxonomic Relationships

Aneides flavipunctatus niger is recognized as a subspecies based on geographic isolation from other populations, morphological and color variation, and ecology (Myers and Maslin 1948). Allozyme studies by Larson (1980) and subsequent reanalysis by Highton (2000) suggested that *A. f. niger* is a distinct lineage. More recent analyses of mitochondrial DNA data supported the allozyme analyses and identified another potentially distinct lineage of *A. flavipunctatus* in the Mount Shasta Region (Rissler and Apodaca 2007). Further genetic studies are ongoing and should help resolve these taxonomic issues. Current work is expanding sampling throughout the range of *A. f. niger* and includes both mitochondrial and nuclear markers (S. Reilly, pers. comm.).

Life History

Little is published on the life history of *Aneides flavipunctatus niger*, and we therefore rely on information from the northern subspecies, the speckled black salamander (*A. f. flavipunctatus*) when data from *A. f. niger* are lacking (see the "Distribution" section). *Aneides flavipunctatus niger* is a terrestrial salamander that can be active year-round in streamside microhabitats (Lynch 1974). Like the majority of salamanders, it is most active on the surface at night, and more so during rain events. Females lay eggs in July or early August (Petranka 1998). In the laboratory, field-collected *A. f. flavipunctatus* from Mendocino County stayed with clutches until the young hatched (N. Staub, pers. obs. in Staub and Wake 2005), but it is unknown whether *A. f. niger* females also attend eggs in the field. Lynch (1981) examined 112 adult

females across the range of *A. flavipunctatus* (including *A. f. niger* populations) and found that females carried 5–25 enlarged ovarian follicles, with fecundity increasing with body size. In the southern populations sampled in this study (which would contain *A. f. niger* samples), an average-sized female was 63 mm SVL, with an estimated clutch size of 9 (Lynch 1981). One record of a natural clutch of *A. f. niger* eggs was found more than 20 cm belowground (Van Denburgh 1895). Like many plethodontid salamanders, eggs undergo direct development, and fully formed, small juveniles appear at the surface shortly after the onset of fall rains, often in October or November (Lynch 1981).

No diet information has been published on *A. f. niger*. We presume that it is a generalized predator of small arthropods and other invertebrates. *Aneides flavipunctatus flavipunctatus* in northern coastal California are generalized predators that eat small invertebrates, including millipedes, beetles, termites, hymenopterans, flies, and collembolans (Lynch 1985).

Habitat Requirements

Aneides flavipunctatus niger is restricted to mesic forests in the fog belt of the outer Coast Range (Myers and Maslin 1948). While salamanders in the genus *Aneides* are sometimes quite arboreal, *A. f. niger* is a ground-dweller (Myers and Maslin 1948). *Aneides flavipunctatus niger* occurs in moist streamside microhabitats and is frequently found in shallow standing water or seeps (Myers and Maslin 1948, Lynch 1974; S. Barry pers., comm.). In these moist microhabitats, *A. f. niger* has been found under stones along stream edges and under boards near creeks (Myers and Maslin 1948). *Aneides flavipunctatus niger* also occurs in talus formations or rock rubble (S. Reilly, pers. comm.).

Distribution (Past and Present)

Aneides flavipunctatus niger is endemic to California and has a small range in the woodlands of the Santa Cruz Mountains in western Santa Clara, northern Santa Cruz, and southernmost San Mateo Counties. *Aneides flavipunctatus*

flavipunctatus occurs from Sonoma County north along the coast into southwestern Oregon and east to Shasta County (Stebbins 2003). Museum specimens exist for the Santa Lucia Mountains (LACM 141882-141883); however, we are unaware of other records for this region, and recent searches in this area have not been successful (S. Reilly, pers. comm.). Lynch (1981) reported that almost all localities of *A. flavipunctatus* (including sites within the range of *A. f. niger*) occurred below 600 m elevation in mesic forests that do not experience sustained freezes.

Some populations of *A. f. niger* have presumably been lost to development. Such losses are most likely to have occurred along the east slope of the Santa Cruz Range as older ranchland has been converted to subdivisions (S. Barry, pers. comm.). However, there is very little documentation of the historical distribution of this taxon.

Trends in Abundance

As for many plethodontids, documenting abundances is exceedingly difficult because *Aneides flavipunctatus niger* spends the majority of its time underground. No reliable population estimates exist for any sites, and therefore no declines in population abundance have been quantitatively documented. Some declines are likely to have taken place due to development and disturbance within the limited geographic range of this taxon. *Aneides flavipunctatus niger* is reported to have been abundant and easily found in the late 1950s, relatively abundant in the 1970s, and difficult to find in recent years (D. Wake, pers. comm.). Range-wide sampling efforts over the last few years have yielded only a handful of specimens (<15) at a few sites, including the UC Santa Cruz campus (S. Reilly, pers. comm.). This anecdotal evidence suggests that declines may have occurred and are possibly ongoing.

Nature and Degree of Threat

Aneides flavipunctatus niger habitat is vulnerable to the effects of logging, spring capping,

and roadbuilding. The Peninsula Open Space Trust has acquired some of the vulnerable property in the northern part of the range, but there is still some risk of further ranchland subdivision (<http://www.openspacetrust.org>; S. Barry, pers. comm.). Climate change may pose some threats to this taxon, particularly given its small range and habitat specificity. Within the range of *A. f. niger*, mean annual temperatures are predicted to increase, though little change is expected in precipitation (reviewed in PRBO 2011). If conditions become significantly warmer and drier, this may affect opportunities for surface activity, although use of moist streamside microhabitats may minimize this effect. The frequency and size of fires in the Coast Ranges is expected to increase up to 50% by the end of the century, although impacts on the forested habitats used by *A. f. niger* are likely to be less severe than in more open habitats (Fried et al. 2004, Lenihan et al. 2008, Westerling and Bryant 2008). The extent of grassland vegetation is predicted to increase, and forested areas are predicted to decrease within the range of *A. f. niger*, which may negatively affect habitat availability (Lenihan et al. 2008).

Status Determination

Aneides flavipunctatus niger is an endemic salamander with a small geographic range in an area with some risk of additional development. However, ongoing declines and population losses have not been well documented, resulting in a Priority 3 designation.

Management Recommendations

Further protection of habitat is key for managing this taxon. In particular, special attention should be given to preserving forests, streamside and spring microhabitats, and natural talus formations within the Santa Cruz Mountains and to maintaining and enhancing connectivity between habitat patches.

Monitoring, Research, and Survey Needs

Basic ecological and life history information is almost entirely lacking for this taxon, as are

estimates of current population abundances, limiting our ability to make more specific management recommendations. Surveys of microhabitats such as streams and seeps in forested areas should be conducted, though disturbance of microhabitat in order to find animals needs to be balanced with concerns regarding continuing decline. These surveys may be more effective if artificial cover objects are placed in suitable habitat, allowing for more comparable survey efforts among localities and increased detectability. Animals are most likely to be

encountered at night when surface conditions are moist. Surveys are needed to establish estimates of abundance and to monitor population sizes over time. Upland terrestrial habitat usage is poorly known, and upland surveys would be useful for determining whether riparian buffers would be beneficial for *Aneides flavipunctatus niger*. Ecological and/or genetic studies of movement ecology and landscape genetics would be useful for understanding connectivity among populations and the permeability of different vegetation types.



INYO MOUNTAINS SALAMANDER

Batrachoseps campi Marlow, Brode, and Wake 1979

Status Summary

Batrachoseps campi is a Priority 3 Species of Special Concern, receiving a Total Score/Total Possible of 50% (55/110). During the previous evaluation, it was also considered a Species of Special Concern (Jennings and Hayes 1994a).

Identification

Batrachoseps campi is one of the largest and most robust members of the diverse plethodontid genus *Batrachoseps* (to 6.1 cm SVL) (Stebbins 2003). The head is relatively broad, and the tail is short compared to other *Batrachoseps* species. The body coloration is dark brown to blackish, with grayish or silvery dorsal spotting which ranges from very sparse to a continuous network. Individuals sometimes have a silvery or greenish cast overall (Stebbins 2003).

This species is the only salamander within its range and thus is unlikely to be confused with other species in the field. With the exception of the Kern Plateau salamander (*B. robus-*

tus) and the largest individuals of the Tehachapi slender salamander (*B. stebbinsi*), other nearby *Batrachoseps* species are noticeably less robust and do not occur east of the Sierra crest. *Hydromantes* species may appear superficially similar

Inyo Mountains Salamander: Risk Factors

Ranking Criteria (Maximum Score)	Score
i. Range size (10)	10
ii. Distribution trend (25)	5
iii. Population concentration/ migration (10)	0
iv. Endemism (10)	10
v. Ecological tolerance (10)	10
vi. Population trend (25)	10
vii. Vulnerability to climate change (10)	0
viii. Projected impacts (10)	10
Total Score	55
Total Possible	110
Total Score/Total Possible	0.50




PHOTO ON PREVIOUS PAGE: Inyo Mountains salamander, Inyo County, California. Courtesy of Adam Clause.

but have five toes on the hind feet rather than four, as is the case in *Batrachoseps* (Stebbins 2003).

Taxonomic Relationships

This species is a member of the *Plethopsis* subgenus of *Batrachoseps*, which also includes the Oregon salamander (*B. wrighti*) from north central Oregon, and *B. robustus* from the Kern Plateau and western margins of Owens Valley in eastern California (Wake et al. 2002). *Plethopsis* can generally be characterized as a stout, robust group of *Batrachoseps* with relatively broad heads. *Batrachoseps campi* is morphologically distinguishable from other *Plethopsis* based on the presence of silvery iridophores, lack of dorsal stripe, and lack of white flecks ventrally (Marlow et al. 1979, Wake et al. 2002, Stebbins 2003). In addition, the species is genetically distinct at allozyme and mitochondrial loci (Yanev 1978, Yanev and Wake 1981, Jockusch and Wake 2002).

Life History

The life history of *Batrachoseps campi* is in need of further study. Its habitat differs somewhat from other closely related *Batrachoseps* species (e.g., *B. robustus*, *B. wrighti*), but information from these taxa is still likely to apply to *B. campi* in several respects. Surface activity occurs at night (Macey and Papenfuss 1991a) during which time the species presumably feeds on a variety of small insects. A life history study of the species is likely to provide important information for future management.

Habitat Requirements

Batrachoseps campi appears to be largely restricted to small patches of riparian habitat associated with perennial springs and limestone fissures in canyons of the Inyo Mountains. Localities where this species has been found contain wet rocks and fissures in close proximity to perennial water (Hansen and Wake 2005a). Salamanders are usually found under wet rocks or in clumps of moist ferns or other cover (Hansen and Wake 2005a). The

species retreats into fissures and rock crevices when surface conditions are not favorable. Habitat surrounding these localized springs consists of Mojave Desert and Great Basin vegetational associations, which are unsuitable for the species. Individuals have only been found away from immediate proximity to flowing water at high-elevation sites in areas of pinyon–juniper woodland (Giuliani 1996, Hansen and Wake 2005a).

Distribution (Past and Present)

Batrachoseps campi is known from a small number of localities on the eastern and western slopes of the Inyo Mountains (Jennings and Hayes 1994a), although additional populations (presumably few) may be discovered in currently unsurveyed sites (Hansen and Wake 2005a). The known elevational range of the species extends from 490 to 2600 m (Macey and Papenfuss 1991a, Hansen and Wake 2005a).

Trends in Abundance

Populations may have declined or been extirpated at a few sites due to habitat modification, though population abundance data are essentially lacking (Papenfuss and Macey 1986). Although data are scarce, most known populations appear to be stable.

Nature and Degree of Threat

The primary threat to this taxon is habitat modification. The overall species range is very small (<20 ha total occupied habitat) and within that range consists of very small, isolated patches of suitable habitat (Hansen and Wake 2005a). The populations in each of these patches are isolated, so recolonization following extirpation is unlikely (Yanev and Wake 1981). Flash floods have scoured the canyon bottoms at some localities, destroying the riparian habitat, though salamander populations appear to persist and slowly recover (Giuliani 1996, Hansen and Wake 2005a). Damage to the sensitive riparian microhabitat from the capture and containment of springs (spring capping), mining, water diversion, and feral burro activity has

occurred at other localities (Papenfuss and Macey 1986). Much of the species' range is unprotected and is vulnerable to further modification.

Status Determination

Due to its small range size and isolated populations, this species is inherently vulnerable to decline. The springs that are essential to its existence are scarce within the species' range and are vulnerable to impacts from water diversion and habitat degradation from humans, livestock, and feral mammals. There are few data on the habitat requirements of this species and the extent to which the isolated population can withstand these impacts. For all of these reasons, a Priority 3 status is justified.

Management Recommendations

The primary management priority for *Batrachoseps campi* is to protect existing habitat. Restoration of degraded habitat would be helpful. However, given the dearth of information on habitat requirements, it is very difficult to know what kinds of restoration would most benefit the species. Thus, restoration efforts need to be informed by the research and monitoring efforts outlined below. Until that time, the riparian areas around desert springs should be protected from modification, specifically with respect to changes in hydrology and vegetation. Some populations, such as the one at Barrel Spring, Inyo County, California, are likely to be

sensitive to relatively minor changes in hydrology (D. Wake, pers. comm.).

Monitoring, Research, and Survey Needs

While the key management priority for this species is simply to protect habitat and minimize disturbances, restoration efforts would require basic research on the size, habitat requirements, and occupancy of sites throughout the species' limited geographic range. In the course of this work, surveyors would need to undertake basic life history research to gather information on population sizes (both census and genetically determined effective population sizes), yearly activity cycles, habitat occupancy, and basic ecological data. Because habitat protection alone is likely to be sufficient to safeguard this species, it may be best to carry out this work only in areas where disturbance to the habitat can be minimized.

Additional desert spring habitat near the known distribution needs to be searched during times when surface moisture is high enough to bring salamanders to the surface, although minimizing damage to these rare habitats is a critical priority. Higher-elevation populations may be more dispersed across the landscape, and surveys should take this into account. Monitoring efforts need to be initiated at localities that have experienced habitat degradation to quantify the ability of *Batrachoseps campi* to tolerate habitat changes that occur as springs are managed for human or livestock needs.



LESSER SLENDER SALAMANDER

Batrachoseps minor Jockusch, Yanev and Wake 2001

Status Summary

Batrachoseps minor is a Priority 1 Species of Special Concern, receiving a Total Score/Total Possible of 71% (78/110). This taxon had not yet been described at the time of the previous Species of Special Concern revision and was therefore not evaluated.

Identification

Salamanders in the genus *Batrachoseps* are generally characterized as elongate, slender plethodontid salamanders with extremely reduced limbs, elongate, worm-like bodies, and extremely long tails that are often longer than the SVL of the animal. Many species have been identified in the last two decades, many of which are morphologically cryptic and some of which have extremely small ranges. *Batrachoseps minor* is the smallest species of *Batrachoseps* (up to 3.4 cm SVL). The coloration is dark blackish brown on the sides and dorsum, sometimes with a lighter brown or tan dorsal stripe along the back (Stebbins 2003). Dense

white speckles are present on the ventral surface (Jockusch et al. 2001).

This species is morphologically similar to the more common and microsympatric black-bellied slender salamander (*B. nigriventris*), though its limbs and feet are relatively more

Lesser Slender Salamander: Risk Factors

Ranking Criteria (Maximum Score)	Score
i. Range size (10)	10
ii. Distribution trend (25)	10
iii. Population concentration/migration (10)	0
iv. Endemism (10)	10
v. Ecological tolerance (10)	3
vi. Population trend (25)	25
vii. Vulnerability to climate change (10)	10
viii. Projected impacts (10)	10
Total Score	78
Total Possible	110
Total Score/Total Possible	0.71




PHOTO ON PREVIOUS PAGE: Lesser slender salamander, San Luis Obispo County, California. Courtesy of William Flaxington.

robust (Hansen and Wake 2005b). Subadults, in particular, can be difficult to tell apart in these species, particularly in some preserved specimens. Molecular identification may be required in some of these cases.

Taxonomic Relationships

Batrachoseps minor was previously included in *B. pacificus* (sensu lato). Populations now regarded as *B. minor* were recognized largely on the basis of mitochondrial DNA and allozymes, though some morphological features distinguish this species from other members of the *B. pacificus* complex (Jockusch et al. 2001). *Batrachoseps minor* is closely related to the San Simeon slender salamander (*B. incognitus*), and the garden slender salamander (*B. major*) (Jockusch et al. 2001, Jockusch and Wake 2002).

Life History

The life history of *Batrachoseps minor* has not been studied. The species presumably feeds on very small insects and other terrestrial invertebrates and exhibits similar ecological characteristics as other members of the *B. pacificus* complex.

The species is microsympatric throughout the entirety of its range with *B. nigriventris*, which is both more widespread and more common than *B. minor* within the range (Hansen and Wake 2005b). It is possible the *B. nigriventris* ecologically replaces *B. minor* at lower elevations (Hansen and Wake 2005b), though the extent or effects of competition between these species has not been studied.

Habitat Requirements

Batrachoseps minor is found on steep north and east-facing mesic slopes within its known range (Jockusch et al. 2001). Known localities have a canopy of oak, tanbark, madrone, and laurel with a poison oak thicket understory (S. Sweet, pers. comm.). These sites remain damp much longer than surrounding slopes, and are 2–3°C cooler at the litter/soil interface (S. Sweet, pers. comm.). Very few localities are

known, and habitat requirements need further study.

Distribution (Past and Present)

Batrachoseps minor is found only in north central San Luis Obispo County. It is present in the southern part of the San Lucia Range above 400 m, ranging from the vicinity of Black Mountain south and east into the Paso Robles and Santa Rita drainages (Jockusch et al. 2001). Populations farther south have been assigned to this species based on morphology and molecular information (E. Jockusch, pers. comm.).

Trends in Abundance

This species was apparently once common within its range. Many specimens were collected throughout the 1970s before the species was described, but the species subsequently became much more difficult to find (Jockusch et al. 2001; D. Wake, pers. comm.). Few specimens have been reported in the literature in the last decade, although several unreported sightings are known, and populations may now be increasing to some degree (Hansen and Wake 2005b; E. Jockusch, pers. comm.; D. Wake, pers. comm.; S. Sweet, pers. comm.). During 1971–1975, field crews associated with the Museum of Vertebrate Zoology undertook 10 field trips that collected 265 *Batrachoseps* from sites known to support *B. minor*. This collection comprised 206 *B. minor* (77% of the total) and 59 *B. nigriventris* (S. Sweet, pers. comm.). In 12 surveys conducted since 2011, 27 *B. minor* have been found along with 60 *B. nigriventris* (31% of the total; S. Sweet, pers. comm.), suggesting that the frequency with which *B. minor* is detected relative to *B. nigriventris* has decreased and that the total number of *Batrachoseps* found is smaller today than it was previously. No obvious changes in habitat or plant cover between the early 1970s and the present that might explain these changes have been observed (S. Sweet, pers. comm.).

Nature and Degree of Threat

Little information is available concerning any aspect of the biology of this species, making

threats difficult to characterize with certainty. Some habitat modification resulting from land conversion to vineyards has occurred within the range, and the invasion of exotic plants has caused changes to the understory in some areas (Hansen and Wake 2005b; D. Wake, pers. comm.); both of these factors are presumably detrimental to the species' persistence. That said, the extent to which such land conversion has occurred has been disputed (S. Sweet, pers. comm.) and a large amount of apparently suitable habitat still remains in the general region. The species was formerly detected in large numbers at wineries (Hansen and Wake 2005b; E. Jockusch, pers. comm.; D. Wake, pers. comm.). Other factors contributing to the declines deserve further study. As this species seems to be limited to relatively mesic areas within its range, changing hydrology and temperature associated with climate change has the potential to render much of the current habitat unsuitable for this species. The marked declines in abundance over the last few decades may indicate a degree of sensitivity to habitat or climatic conditions or, alternatively, may simply represent a temporary and cyclical decline associated with moderate-term changes in climate (rainfall specifically; S. Sweet, pers. comm.). Here, we interpret the observed pattern with precaution in mind, treating the documented declines in abundance as real and noncyclical, but acknowledging that an alternative possibility exists and that further study and published data are needed.

Status Determination

Batrachoseps minor is a California endemic and has an exceedingly small geographic range. Large apparent declines have occurred since the 1970s, and the threats to this taxon are poorly understood, leading to a Priority 1 status.

Management Recommendations

Given what is currently known about this species, little can be done in terms of management. Few sites have been confirmed (using

molecular data) to support *Batrachoseps minor*, and these sites should be protected from further modification that is likely to be detrimental to salamander populations. Additional information on the range, habitat requirements, and environmental sensitivity of the species is needed to help guide future management.

Monitoring, Research, and Survey Needs

Batrachoseps minor is poorly known biologically, and published accounts of even the most basic habitat and ecological data are largely lacking for the species. Additional and ongoing surveys for this taxon are needed to help determine its range, both geographically and ecologically. However, careful attention needs to be paid to effective identification of specimens that are found. Because *B. minor* is so similar in appearance to *B. nigriventris*, and the two species occur in microsympatry, surveyors need to have extensive experience distinguishing different *Batrachoseps* species from each other. Subadult specimens of *B. minor* may require molecular identification unless and until field-validated morphological characters can be identified. As the status of remaining populations is unknown, a reasonable management policy would be that no *Batrachoseps* from the known or suspected range of *B. minor* be removed from the wild unless the collector has extensive experience identifying these species. Rather, individuals should be photographed and non-destructively sampled, preferably by removing a small portion from the end of the tail (~2 mm) and genotyped to establish identification. If a few replicate DNA sequences from both the nuclear and mitochondrial genomes could be established as reliable barcoding genes, DNA typing could be accomplished quickly and inexpensively. Surveys should take place when surface conditions are appropriately moist to enhance the likelihood of finding populations of this elusive salamander. The chances of finding *B. minor* without disturbing its natural habitat would likely be increased by establishing a transect of artificial cover objects

(plywood boards) throughout the known range. Nighttime surveys during rain events might also be productive. In addition, nearby areas should continue to be surveyed for this species, as its distribution could potentially be larger, both ecologically and geographically, than is presently known. Higher-elevation areas, such as those in the vicinity of Santa Rita and Old Creek Road, San Luis Obispo County, should be surveyed if access to private land in these areas can be established. It is possible that the known localities occur near the lower elevational range of the species, and larger populations exist at higher elevations (E. Jockusch, pers. comm.). Recent and

repeated surveys in some of these areas have failed to detect this species, which suggests elevation may not be an important factor (S. Sweet, pers. comm.). Nevertheless, the species is clearly less detectable than it was decades ago and additional published data are needed to better characterize the known distribution and abundance. Additional research into potential causes of the declines in detectability should also be pursued. In particular, screens of museum specimens for the presence of pathogenic fungi might be fruitful (D. Wake, pers. comm.), as could study of decadal scale climate and rainfall patterns within the species known range (S. Sweet, pers. comm.).



RELICTUAL SLENDER SALAMANDER

Batrachoseps relictus Brame and Murray 1968

Status Summary

Batrachoseps relictus is a Priority 1 Species of Special Concern, receiving a Total Score/Total Possible of 60% (66/110). It was also considered a Species of Special Concern during the previous evaluation (Jennings and Hayes 1994a); however, the range of the species has since been greatly reduced as a consequence of taxonomic revisions.

Identification

As is typical of its genus, *Batrachoseps relictus* is a small, elongate, worm-like salamander with a slender body, long tail, and tiny limbs. The dorsal coloration is blackish brown with a lighter, often indistinct dorsal stripe that may be reddish, yellowish, or dark brown (Stebbins 2003). *Batrachoseps relictus* is one of the smallest members of its genus. SVLs of mature animals collected at the type locality in the lower Kern River Canyon (see the "Distribution" section) averaged 30.2 mm, while those from Breckenridge Mountain averaged somewhat larger at

39 mm SVL (Jockusch et al. 2012). *Batrachoseps relictus* also has relatively few trunk vertebrae, with a modal number of 17 from the type locality (Brame and Murray 1968) and counts as low as 17 occurring with low frequency in the

Relictual Slender Salamander: Risk Factors

Ranking Criteria (Maximum Score)	Score
i. Range size (10)	10
ii. Distribution trend (25)	10
iii. Population concentration/ migration (10)	0
iv. Endemism (10)	10
v. Ecological tolerance (10)	7
vi. Population trend (25)	15
vii. Vulnerability to climate change (10)	7
viii. Projected impacts (10)	7
Total Score	66
Total Possible	110
Total Score/Total Possible	0.60




PHOTO ON PREVIOUS PAGE: Relictual slender salamander, Kern County, California. Courtesy of William Flaxington.

Breckenridge Mountain populations (Jockusch et al. 2012).

Several other species of *Batrachoseps* occur in the same region of the southern Sierra Nevada, and geographic range is the best way to distinguish animals in the field. Individuals from the upper Kern River Canyon (Greenhorn Mountains slender salamanders, *B. altasierrae*) that were previously considered a part of *B. relictus* (see the "Taxonomic Relationships" section) have relatively longer trunks, smaller heads, shorter limbs, and smaller feet (Jockusch et al. 2012). Female *B. altasierrae* have fewer maxillary teeth, and the vomerine teeth in both sexes are patchily distributed, compared to being arranged in rows in *B. relictus* (Jockusch et al. 2012).

In the lower Kern River Canyon, the range of *B. relictus* overlaps with Kern Canyon slender salamanders (*B. simatus*) and the yellow-blotched ensatina (*Ensatina eschscholtzii croceator*; Brame and Murray 1968). Unlike *B. relictus*, *B. simatus* is not closely associated with water, and populations of *B. relictus* at elevations where *B. simatus* occurs are likely extirpated (see the "Distribution" section). *Ensatina eschscholtzii croceator* is a larger, more robust salamander and is easily distinguished by conspicuous yellow blotches on the dorsum and a much larger body form (Stebbins 2003).

Taxonomic Relationships

The populations included in *Batrachoseps relictus* have changed considerably since its original description. Brame and Murray (1968) considered several geographically disjunct populations as belonging to *B. relictus*, most of which are now recognized as distinct species (Yanev 1978, Yanev 1980, Jockusch et al. 1998, Wake and Jockusch 2000, Jockusch et al. 2001). Populations in the Sierra Nevada from the Merced River to the Kern River were considered a part of the *relictus* group (Yanev 1980) and were split into four allopatric species by Jockusch et al. (1998). At that time, *B. relictus* was thought to range from the Tule River drainage to the lower Kern River Canyon, including populations in the Greenhorn Mountains (Jockusch et al. 1998,

Jockusch and Wake 2002). Since then, populations from the upper Kern River have been found to be morphologically distinct from salamanders at the *B. relictus* type locality, and have been described as the new species *B. altasierrae*, the Greenhorn Mountains slender salamander (Jockusch et al. 2012). Populations of *Batrachoseps* on Breckenridge Mountain were discovered in 1979. Jennings and Hayes (1994a) designated this putative taxon as a Species of Special Concern. Recent morphometric analyses have shown that populations from Breckenridge Mountain are most similar to *B. relictus* from the type locality, and are now included as *B. relictus* (Jockusch et al. 2012). Given the description of the new taxon *B. altasierrae*, the classification of Breckenridge Mountain populations as *B. relictus*, and the presumed extirpation of the type locality (see the "Distribution" section), extant *B. relictus* only occur on Breckenridge Mountain under the current taxonomic arrangement.

While we follow the recommendations of Jockusch et al. (2012) to recognize *Batrachoseps* from Breckenridge Mountain as *B. relictus*, it is important to note that their phylogenetic analyses of mitochondrial DNA show these populations as nested within *B. simatus*, the Kern Canyon slender salamander. Jockusch et al. (2012) argued that allozyme data and unpublished nuclear data recovered a different pattern that corroborated the distinctiveness of *B. relictus*, and that the mitochondrial DNA results were potentially explained by introgression from *B. simatus* into *B. relictus*. This interpretation appears to be reasonable. However, given the complexity of this group, it remains possible that additional work may lead to further taxonomic revisions.

Life History

Very little is known about the natural history of *Batrachoseps relictus*, and much of the ecological literature published under this name refers to what is now classified as *B. altasierrae*. *Batrachoseps relictus* on Breckenridge Mountain (1700–2000 m elevation) have been found surface active under cover objects from May to early

October (Jockusch et al. 2012). At lower elevations in the Kern River Canyon, animals have been collected between January and May, suggesting that surface activity is possible over most of the year and varies with elevation. Association with aquatic microhabitats likely facilitates extended periods of surface activity (see the "Habitat Requirements" section). Like other plethodontid salamanders, *B. relictus* is a direct developer that lays terrestrial eggs. Females have been found with yolked ova or eggs in May and June (Jockusch et al. 2012). A communal nest with roughly 125 eggs and 20 adults was discovered beneath a rock in a seep during June 1979 at the high-elevation site on Breckenridge Mountain (R. Hansen, pers. obs., in Jockusch et al. 2012; observation incorrectly ascribed to *B. simatus* in Stebbins 1985). Diet has not been studied in *B. relictus*. Presumably they use their projectile tongues to catch small invertebrates, as do other *Batrachoseps* species (Hansen and Wake 2005c).

Habitat Requirements

Individuals from the type locality in the lower Kern River Canyon have been found associated with perennial springs, seeps, and small creeks in oak woodland below 750 m (Hilton 1948, Brame and Murray 1968). This close association with water was described as "semiaquatic" by Brame and Murray (1968). Animals have been found under cover objects with water beneath them and observed in the water (Hilton 1948, Jockusch et al. 2012). On Breckenridge Mountain the dominant vegetation type at extant localities is pine–fir forest (Jockusch et al. 2012). East of Squirrel Meadow at 2000 m elevation, *Batrachoseps relictus* is typically associated with a small seep and sandy or gravel substrate (Jockusch et al. 2012). Use of upland habitat away from water is unknown, but two adults were found 45 m upslope from seep habitat at the Squirrel Meadow site (Jockusch et al. 2012). At Lucas Creek, the lower-elevation extant locality on Breckenridge Mountain (1665 m), all *B. relictus* to date have been found under cover objects along a 750 m stretch of stream (Jockusch et al. 2012).

Distribution (Past and Present)

The type locality is in the lower Kern River Canyon, 150 yards above the junction of state Highway 178 and the road turnoff to Democrat Hot Springs and Resort (Brame and Murray 1968). Despite repeated, careful searches, *Batrachoseps relictus* have not been seen at the type locality since 1970 (Jockusch et al. 2012; incorrectly reported as 1971 elsewhere). Extirpation of the type locality may have been caused by the degradation of the sensitive seep and spring habitat due to the construction of Highway 178 (Hansen 1988). With the presumed extirpation of the type locality, *B. relictus* is now thought to be restricted to two localities on Breckenridge Mountain, and has the smallest known range for any described species of *Batrachoseps*. Populations north of the Kern River including the Greenhorn Mountains are no longer considered a part of *B. relictus* (see the "Taxonomic Relationships" section). The known elevation range is from 480 m in the Lower Kern Canyon River up to 2000 m on Breckenridge Mountain (Jockusch et al. 2012).

Trends in Abundance

Declines are suspected at one extant site, the area east of Squirrel Meadow on Breckenridge Mountain. The locality was first discovered in 1979 but later degraded by construction of a logging road through *Batrachoseps relictus* habitat. Salamanders were not seen at this site for two decades, with declines presumed to be due to habitat degradation from road construction, wildfire, and timber harvest (Jockusch et al. 2012). More recent surveys of the site have found that populations appear to be rebounding to some degree (Jockusch et al. 2012). Whether such variation in abundance over time is typical, due to detection difficulty, or actual anthropogenic declines is unknown.

Nature and Degree of Threat

The major threat to *Batrachoseps relictus* is habitat degradation, particularly of sensitive spring and seep habitat. Climate change is expected to increase temperatures in the Sierra Nevada,

although changes in precipitation and fire regime are highly uncertain and large regional variation is expected across the mountain chain and at different elevations (reviewed in PRBO 2011). If conditions become warmer and drier, this would presumably negatively affect *B. relictus* populations, although microhabitat characteristics are likely key to determining surface activity and population stability. Large reductions in snowpack are predicted for the Sierra Nevada (reviewed in PRBO 2011), which may decrease the availability of streamside habitat for *B. relictus*.

Status Determination

The extremely limited geographic range of *Batrachoseps relictus*, the small number of known extant populations, and apparent extirpation of the type locality contribute to a Priority 1 Species of Special Concern designation for the species.

Management Recommendations

Protecting the two remaining localities from habitat degradation is critical to the persistence of *Batrachoseps relictus*. Given the extremely sensitive and restricted range of the species, any habitat modification should be avoided where the species still occurs. Road construction should be avoided, and road use and maintenance activities should be restricted, or ideally eliminated altogether. Timber harvest and use of heavy equipment in or near seeps and streams should be eliminated. If the type locality is confirmed to be extirpated, then repatriation of the species to the type locality may be appropriate. However, the lack of genetic infor-

mation from this site (no genetic samples exist) and the overall state of flux in the classification of southern Sierra Nevada *Batrachoseps* may argue against such reintroductions pending further molecular systematics work on the group as a whole. Although the extent and use of upland habitat is unknown, protection of riparian buffers would almost certainly benefit this species in disturbed areas. In addition, it is probably reasonable to assume that livestock grazing should be eliminated from areas where the species still occurs, at least until field ecological studies indicate that grazing is compatible with the salamander's habitat requirements.

Monitoring, Research, and Survey Needs

Basic life history and population biology information is severely lacking for this species, and represents a critical research need. A key survey need is to attempt to locate additional populations, particularly at mid-elevations on Breckenridge Mountain, which are largely unexplored (Jockusch et al. 2012). High-priority sites for surveys include streamside and seep habitats on the north face of the mountain. Monitoring should continue at the lower Kern River Canyon localities to confirm extirpation. Populations at the higher-elevation Breckenridge Mountain locality went undetected for many years, and it remains possible that animals could be rediscovered at the type locality. If so, the collection of genetic samples would be invaluable to support or refine the current taxonomy of the species, and to help determine patterns of connectivity among remaining populations.



CALIFORNIA GIANT SALAMANDER

Dicamptodon ensatus (Eschscholtz 1833)

Status Summary

Dicamptodon ensatus is a Priority 3 Species of Special Concern, receiving a Total Score/Total Possible of 66% (56/85). This species was not previously considered a Species of Special Concern (Jennings and Hayes 1994a).

Identification

Dicamptodon ensatus is a large (6.3–17.3 cm SVL) robust salamander with a very large head and stout limbs. The dorsal coloration is a coppery tan to dark brown irregular marbled pattern on a tan to light reddish brown background. The venter is paler and usually unmarked, although marbling often extends onto the chin, throat, and under the legs. The marbling coloration is often brighter in young metamorphs compared to adults. The tail is laterally compressed, the skin is smooth, and post-metamorphic juveniles and adults lack tubercles on their feet (Stebbins 2003).

Larvae are of the stream type, with short bushy gills and a tail fin that begins at the inser-

tion of the hind limbs and extends posteriorly to the tail tip. Larval dorsal coloration is light brown, and ventral coloration is white to yellowish white (Nussbaum 1976). There is also a pale

California Giant Salamander: Risk Factors

Ranking Criteria (Maximum Score)	Score
i. Range size (10)	10
ii. Distribution trend (25)	10
iii. Population concentration/ migration (10)	10
iv. Endemism (10)	10
v. Ecological tolerance (10)	10
vi. Population trend (25)	Data deficient
vii. Vulnerability to climate change (10)	3
viii. Projected impacts (10)	3
Total Score	56
Total Possible	85
Total Score/Total Possible	0.66




PHOTO ON PREVIOUS PAGE: California giant salamander, Santa Cruz County, California. Courtesy of Nicholas Hess.

eye stripe behind each eye, and the snout is depressed (Petranka 1998). The toe tips of larvae are black and cornified (Petranka 1998).

In California, *D. ensatus* is largely indistinguishable from the more widely distributed coastal giant salamander (*D. tenebrosus*) based on morphology alone. However, both geographic range and genetic markers distinguish these two species.

Taxonomic Relationships

Good (1989) split California *Dicamptodon* into two species, *D. tenebrosus* in the north and *D. ensatus* in the south, on the basis of allozyme data. A 4.7 km hybrid zone exists between the two species approximately 10 km north of Gualala in Mendocino County (Good 1989). Otherwise, the two species are allopatric.

Life History

Adult *Dicamptodon ensatus* are terrestrial and return to streams to breed during the fall rainy season (Kessel and Kessel 1943a) and in the spring (Stebbins 2003). One *D. ensatus* nest of approximately 70 eggs was found under a submerged wooden plank in a rapidly flowing stream in the Santa Cruz Mountains, San Mateo County, during June (Henry and Twitty 1940). Female *D. tenebrosus* guard nests through hatching (Nussbaum et al. 1983), and an adult female *D. ensatus* was found near the Santa Cruz Mountains nest (Henry and Twitty 1940), suggesting that both species may guard their eggs. Eggs in early developmental stages are pure white and approximately 5.5 mm in diameter (Petranka 1998). The larval stage lasts approximately 18 months, with larvae growing 8–12 mm in TL per month during the warmer months in their first year. Larvae reach 10 cm TL within a year of hatching and metamorphose in late summer at 13–14 cm TL (Kessel and Kessel 1943a, Kessel and Kessel 1943b, Kessel and Kessel 1944). The prevalence of pedomorphosis in this species is unknown, although it can be quite common in *D. tenebrosus*. A pedomorphic population of *D. ensatus* has been reported from caves on the UC

Santa Cruz campus (B. Sinervo, unpublished data).

Bury (1972) reported gut contents of 12 adults from Del Norte, Humboldt, and Marin Counties (i.e., a mix of *D. ensatus* and *D. tenebrosus*). Eight out of 12 specimens contained one or more vertebrates, including California slender salamanders (*Batrachoseps attenuatus*), lizards, mice, shrews, and voles. Other prey included large invertebrates such as land snails and smaller invertebrates such as beetles and crickets (Bury 1972). Cannibalism has been documented in adults (Anderson 1960). No diet data from larvae are available for this species, though they are presumed to have similar diets to larval *D. tenebrosus* (Petranka 1998), which primarily consume aquatic insects and other invertebrates (Parker 1994).

Habitat Requirements

Dicamptodon ensatus occurs in mesic coastal forests (oak woodland and coniferous forest; Petranka 1998), and coastal chaparral habitat is used in southern Marin County and San Mateo County (N. Waters, pers. comm.). Very little is known about terrestrial habitat use by adults and metamorphs, although adults are occasionally found surface active or under cover objects in wet conditions (Petranka 1998). One unusual record exists of an adult *D. ensatus* in a tree vole (*Arborimus pomo*) nest 2.4 m off the ground, the only account of arboreality in this species (D. Hamilton and W. Roberts, unpublished data in Forsman and Swingle 2007).

Breeding and larval development occurs in cold permanent and semipermanent streams (Petranka 1998). Larval habitat use is poorly studied. In one stream, small larvae were found in slow-moving water near the banks during heavy flows, and as flows decreased they moved into the main stream channel where larger larvae occurred (Kessel and Kessel 1943a, Kessel and Kessel 1943b).

Distribution (Past and Present)

Dicamptodon ensatus is endemic to California, occupying a small range from sea level to

900 m in elevation along the coast in two isolated areas near San Francisco Bay (Stebbins 2003). North of the Bay, they occur in the outer Coast Ranges from near the southern border of Mendocino County south through Marin County, and the inner Coast Ranges in Napa, Sonoma, Lake, and Solano Counties (Good 1989). South of the Bay, they occur in the Santa Cruz Mountains in San Mateo, Santa Clara, and Santa Cruz Counties (Good 1989; N. Waters, pers. comm.). *Dicamptodon ensatus* has not been recorded in the East Bay (Stebbins 2003). Nussbaum (1976) mentioned an unconfirmed sight record from the Santa Lucia Mountains in Monterey County. Multiple surveys by several researchers over the decades have attempted to verify this account with no individuals detected (N. Waters, pers. comm.). While extirpations have not been documented, urbanization, agriculture, and timber harvest have likely resulted in some population losses, particularly due to development in the southern part of the range (Bury 2005; S. Barry, pers. comm.)

Trends in Abundance

Given the paucity of information, this species is currently considered data deficient for the population trend metric. However, it is likely that abundance has been reduced in habitats disturbed by urbanization, roadbuilding, logging, or water diversions (Bury 2005).

Nature and Degree of Threat

The Santa Cruz Mountains isolate is currently largely contained within a network of public parkland, though the extent of possible losses in this region due to past development is poorly understood (N. Waters, pers. comm., S. Barry, pers. comm.). Coast Range populations in the north are likely subject to negative effects from timber harvest and development, though this area is less urbanized than the southern part of the range. Disturbances such as clear-cutting and road construction can lead to lower abundances in *Dicamptodon tenebrosus* (Corn and Bury 1989, Welsh and Ollivier 1998). Other threats include fragmentation of riparian habi-

tat, water diversions for municipal and agricultural use, and road mortality (N. Waters, pers. comm.).

Climate change may negatively impact *D. ensatus*, although uncertainty in climate projections coupled with limited ecological information makes assessing risk difficult. Mean annual temperature is expected to increase while projected changes in precipitation are likely modest, leading to warmer and possibly drier conditions in northwestern and central California (reviewed in PRBO 2011). At the same time, upwelling is expected to intensify (Snyder et al. 2003, Lebaszi et al. 2009). This may increase fog development and contribute to cooler, moister conditions along the coast, potentially ameliorating effects of warming or drying within the range of *D. ensatus*. The frequency and extent of wildfire is expected to increase in the region encompassing the southern part of the range, with predicted increases in area burned of up to 50% (Fried et al. 2004, Lenihan et al. 2008, Westerling and Bryant 2008). How fire regime will change in the northern part of the range is less well understood (reviewed in PRBO 2011). Effects of wildfire on *D. ensatus* are unknown, though mortality and habitat degradation due to fire has been documented in other stream-breeding amphibians (e.g., Gamradt and Katz 1997, Pilliod et al. 2003). In northwestern California, vegetation communities are expected to shift from moist conifer to drier mixed evergreen forest, with reductions in Douglas fir and redwood forest in particular (Lenihan et al. 2008, PRBO 2011), which may impact the availability of *D. ensatus* habitat.

Status Determination

Dicamptodon ensatus is an endemic, ecologically specialized salamander with a small geographic range that is restricted to an area with a high human population density. These factors combine to place it at high risk of habitat loss and disturbance. However, data are not available to determine whether ongoing declines and population losses have occurred, resulting in a Priority 3 designation for this species.

Management Recommendations

We know little about the basic biology of this species, which makes it difficult to formulate management recommendations beyond minimizing disturbances to existing habitat. Habitat protection may be particularly important for small headwater streams where siltation and other stream disturbances are known to severely impact other *Dicamptodon* species. Construction and use of roads should be eliminated or minimized within *D. ensatus* habitat, particularly during the breeding season. Riparian buffer vegetation should be retained in areas that are developed or harvested, though efficacy of buffers and optimal buffer widths for this taxon are unknown.

Monitoring, Research, and Survey Needs

Distribution, abundance, habitat requirements, and life history of *Dicamptodon ensatus* all need further study. Most research to date has focused on the more widespread *D. tenebrosus* to the north and was conducted before the two species were recognized as distinct. This substantial knowledge gap needs to be addressed with basic ecological studies. Nothing is known about dispersal in this species, especially the importance of movement through terrestrial habitats. Both mark-recapture and landscape genetic studies are needed for *D. ensatus*. Studies are also needed that examine the efficacy of streamside

buffers in ameliorating the effects of disturbance on stream habitats. Such studies should be replicated both north and south of San Francisco Bay, given that these are completely isolated population segments living in different habitats. Distributional surveys are particularly needed in the Inner Coast Range portion of the northern range (N. Waters, pers. comm.).

While larvae are easy to find by searching aquatic habitats, transformed *D. ensatus* are infrequently encountered using typical amphibian survey techniques. For example, only 12 individuals were captured in 18,032 trap nights over 3 years of pitfall trapping along 840 m of drift fence in suitable habitat at Point Reyes National Seashore (G. Fellers and D. Pratt, unpublished data, in Fellers et al. 2010). In the same study, no *Dicamptodon* were detected under 84 coverboards during nearly 2000 coverboard checks. However, culvert removal using heavy equipment uncovered aggregations of >20 adults at the same study sites, suggesting that terrestrial sampling may severely underestimate abundance (Fellers et al. 2010). Another account from Santa Cruz County reported several adults and eggs getting washed out of a drill hole made 6 m into a hillside to access a subterranean spring (Dethlefsen 1948). These reports suggest that metamorphosed individuals may be largely subterranean in their habits, a possibility that needs further investigation.



SOUTHERN TORRENT SALAMANDER

Rhyacotriton variegatus Stebbins and Lowe 1951

Status Summary

Rhyacotriton variegatus is a Priority I Species of Special Concern, receiving a Total Score/Total Possible of 75% (83/110). Previously it was also considered a Species of Special Concern, although at a lower priority level. Additional research on ecology and phylogeography since Jennings and Hayes (1994a) supports this change in status.

Identification

Rhyacotriton variegatus is a small to medium-sized salamander (5 cm SVL) (Welsh and Lind 1992, Tait and Diller 2006), with a small head and a short, laterally compressed tail (Stebbins 2003). Expanded square-shaped glands lateral and posterior to the vent in adult males distinguish this genus from all other North American salamanders (Petranka 1998). *Rhyacotriton* has large bulging eyes, with eye diameter roughly equal to the distance between the anterior edge of the eye and the tip of the snout (Stebbins 2003). The dorsal ground

color is brownish to olive, and the venter is yellow to yellowish green with a sharp, abrupt demarcation between the dorsal and ventral coloration (Petranka 1998). California *R. variegatus* are heavily speckled with small dark spots

Southern Torrent Salamander: Risk Factors

Ranking Criteria (Maximum Score)	Score
i. Range size (10)	10
ii. Distribution trend (25)	20
iii. Population concentration/ migration (10)	10
iv. Endemism (10)	3
v. Ecological tolerance (10)	10
vi. Population trend (25)	10
vii. Vulnerability to climate change (10)	10
viii. Projected impacts (10)	10
Total Score	83
Total Possible	110
Total Score/Total Possible	0.75




PHOTO ON PREVIOUS PAGE: Southern torrent salamander, Mendocino County, California. Courtesy of Robert Thomson.

on the dorsum and venter (Good and Wake 1992).

Larvae are of the stream type and have morphological adaptations unique to headwater specialists (Valentine and Dennis 1964). Larvae have short stubby gills and a tail fin that does not extend anteriorly onto the trunk. The dorsum is light brown above, the venter is cream to yellow, and the body is sprinkled with dark speckling above and below except on the tail fin. The eyes are prominent and dorsally positioned (Petranka 1998).

Taxonomic Relationships

Rhyacotriton variegatus has been recognized as a species since 1992 based on protein variation (Good and Wake 1992). Miller et al. (2006) identified three mitochondrial DNA clades within *R. variegatus*. The California clade/southern Oregon clade split occurs at the Smith River in California, a common biogeographic boundary. Miller et al. (2006) concluded that the California clade constitutes an evolutionarily significant unit (sensu Moritz 1994). The California clade is endemic to the state with a ~50% smaller range than the species as a whole, and the southern Oregon clade animals in California have an extremely small range. Although Miller et al. (2006) recognized these clades as potential management units, we consider them as a single taxon here pending additional research on their geographic ranges and genetic distinctiveness using additional molecular markers.

Life History

Breeding may occur throughout much of the year. Males produce sperm year-round, with peak production from February through April (Humboldt County; Tait and Diller 2006). California females have been found carrying spermatophores from February through June (Stebbins and Lowe 1951, Tait and Diller 2006), and females from an Oregon population had cloacal spermatophores as late as October (Nussbaum and Tait 1977).

Females produce smaller clutches than most similarly sized stream-breeding salamanders

(Petranka 1998), with gravid females carrying from 4 to 16 ovarian eggs (Nussbaum and Tait 1977, Good and Wake 1992, Tait and Diller 2006). Karraker (1999) found a nest with 11 cream-colored eggs deposited singly beneath a small boulder in a first-order stream channel in Humboldt County.

Developmental times are slow, with oviposition to sexual maturity taking approximately 4.5 years (Nussbaum and Tait 1977, Tait and Diller 2006). Time from oviposition to hatching is roughly 8 months (Karraker 1999), with time from oviposition to absorption of yolk probably closer to a year (Tait and Diller 2006). Peak oviposition is in August and September in California, with peak hatching occurring in the spring (Humboldt County; Tait and Diller 2006). Larval development from hatching to metamorphosis takes 2–2.5 years (Nussbaum and Tait 1977, Tait and Diller 2006). After metamorphosis, an additional 1–1.5 years of growth is required before sexual maturity is attained (Nussbaum and Tait 1977, Tait and Diller 2006).

The extended reproductive period and overwintering of larvae result in overlapping size cohorts in streams (Welsh and Lind 1992, Tait and Diller 2006). Hatchlings are 14–16 mm SVL (Tait and Diller 2006), and size at metamorphosis is around 35 mm SVL (Nussbaum and Tait 1977, Good and Wake 1992, Tait and Diller 2006). In Humboldt County, larval growth rates were recorded as 2.3 mm/year in Six Rivers National Forest (Welsh and Lind 1992) and 8.9 mm/year in a more coastal site in the Mad River drainage (Tait and Diller 2006). Larvae and adults weighed more in the spring than fall at one site, suggesting active foraging and growth over the winter months (Welsh and Lind 1992).

Adults are active at air and water temperatures of 5–10°C, lower than those known for any other aquatic salamander (Stebbins and Lowe 1951, Stebbins 1955, Brattstrom 1963). The average critical thermal maximum for adults and larvae are also lower than reported for other salamanders (larvae: 26.7°C; adults:

27.9°C; Bury 2008b). Welsh and Lind (1996) observed signs of stress in adults at 17.2°C. Thermal tolerances of eggs are unknown (Bury 2008b).

Very few data are available on movement or diet in this species. One mark-recapture study at a single headwater stream/seep site in Humboldt County found extremely low levels of movement, with approximately 1 m/year of movement for adults and 2 m/year for larvae on average (Welsh and Lind 1992). However, unrecaptured animals may have moved longer distances (20% of originally marked animals were recaptured). The diet of *Rhyacotriton variegatus* appears to be generalized on aquatic and semiaquatic invertebrates, with amphipods and collembolans the most abundant prey (Bury and Martin 1967).

Habitat Requirements

Rhyacotriton variegatus occurs within a relatively narrow range of ecological conditions that are typical of late-seral forests. These conditions include cold, clear, flowing permanent seeps and headwater to low-order streams with coarse, rocky substrates in mesic to moist forests (Welsh and Lind 1988, Welsh 1990, Welsh and Lind 1991, Welsh and Lind 1996, Vesely and McComb 2002, Welsh et al. 2005, Ashton et al. 2006, Welsh and Hodgson 2011). Key habitat requirements are the maintenance of cold water temperatures (6.5–15°C) and presence of loose substrates composed of gravel and cobble (Diller and Wallace 1996, Welsh and Lind 1996, Stoddard and Hayes 2005, Welsh et al. 2005, Bury 2008b, Welsh and Hodgson 2008). In the Mattole Watershed, *R. variegatus* occurred primarily in undisturbed headwater channels and was never detected in streams where canopy closure was less than 91% or water temperatures were warmer than 13.5°C (Welsh and Hodgson 2011). *Rhyacotriton variegatus* is extremely desiccation intolerant (Ray 1958), although it will occasionally venture away from the stream channel and use riparian and forest habitat in the wet season (Vesely and McComb 2002; Vesely and McComb, pers. obs., in Welsh and Lind 1996).

Rhyacotriton variegatus is sensitive to fine sediment load and embeddedness (Welsh and Lind 1996, Welsh and Ollivier 1998) and has been found to be positively associated with high-gradient streams, particularly in areas with timber harvesting. This may be due to stream network processes that flush fine sediments out of high-gradient reaches (Corn and Bury 1989, Diller and Wallace 1996, Stoddard and Hayes 2005, Ashton et al. 2006). In a review of seven studies of *R. variegatus* habitat associations, Welsh and Hodgson (2008) found that the species occurred at sites where fine sediment ranged from 2% to 40%, and zero detections occurred when more than 65% of the coarse substrate was embedded with fine sediment.

Distribution (Past and Present)

Rhyacotriton variegatus occurs patchily at elevations below 1469 m throughout the Pacific Coast Ranges of Oregon and California, from the Little Nestucca River and Grande Ronde Valley in Oregon to near Alder Creek in Mendocino County in California (Good and Wake 1992). Populations also occur in the Cascade Range in Oregon (Good and Wake 1992, Miller et al. 2006). A previously reported disjunct population in the McCloud River, Siskiyou County, appears to be based on incorrectly identified museum specimens of the southern long-toed salamander (*Ambystoma macrodactylum sigillatum*) in the California State University, Chico collection.

Suitable microhabitat is patchily distributed in California, and *R. variegatus* is only found in suitable sites about half of the time. Random stratified sampling of 117 sites throughout the geographic range in California found that 45% of sites contained suitable microhabitat, but only 62% of those sites were occupied (Welsh and Lind 1992). Sampling of 38 different sites in the same region selected for the US Forest Service "Old-growth Wildlife Project" found suitable microhabitat in 79% of sites, with *R. variegatus* present in 47% of suitable sites (Welsh and Lind 1992). Systematic stratified

sampling of 53 mixed conifer–hardwood stands on public lands in northern California found *R. variegatus* at 62% of sites (Welsh and Lind 1996).

Some of the variation in distribution can be explained by forest age and timber harvest histories, with *R. variegatus* more often found in older, unharvested stands. Welsh (1990) surveyed spring and seep habitats in 34 forest stands in the Coast Ranges in California and southern Oregon ranging from 30 to 560 years old and at elevations of 150–1500 m. *Rhyacotriton variegatus* was found in 70% of old-growth stands, 50% of mature stands, and 11% of young stands. Recent surveys of the Mattole Watershed in northern California (Humboldt and Mendocino Counties) found *R. variegatus* mostly in late-seral headwater tributaries, habitats that are now rare in the watershed (Welsh et al. 2005, Welsh and Hodgson 2011). However, occupancy rates were higher in young forests along the coast where temperatures are mediated by the maritime climate: *R. variegatus* was found in 48% of 30 m sampling reaches and 80% of entire stream reaches in stands less than 80 years old (Diller and Wallace 1996).

Exact figures are difficult to come by, but most of the historical coastal old-growth habitat in California is now gone (85–96.5% gone; references in USFWS 1997). In addition to habitat modification, several investigators have hypothesized that *Dicamptodon* predation may restrict *Rhyacotriton* distribution to small headwater streams (e.g., Stebbins 1955, Nussbaum 1969, Welsh and Lind 1996, Welsh and Ollivier 1998). However, Rundio and Olson (2001) found that *R. variegatus* larvae were unpalatable to *D. tenebrosus* larvae, surviving 90% of encounters in experimental trials.

Trends in Abundance

Estimates of abundance are not available for time periods before timber harvesting became a prominent factor in landscape management, but the highest documented abundances over the last several decades have been in late-seral sites, supporting the idea that abundances are

reduced in response to disturbances such as timber harvest and road building. *Rhyacotriton variegatus* can be locally abundant, with densities of up to 22 salamanders/m² recorded in suitable streamside habitat at an old-growth site in Six Rivers National Forest, Humboldt County (Welsh and Lind 1992). However, most sites in that study yielded 1–5 captures/10 m² (Welsh and Lind 1992). By sampling across the range of *R. variegatus* in California and across stands of different ages, Welsh and Lind (1996) documented a much lower mean density of 0.68 salamanders/m². In young stands in coastal northern California (<80 years old), Diller and Wallace (1996) found that densities were 0.18–5.5 salamanders/m². Welsh et al. (2000) reanalyzed Welsh and Ollivier's (1998) data from sites in Prairie Creek Redwoods State Park in Humboldt County for comparison to encounter rate data reported by Wroble and Waters (1989) from timber company lands in the same county. *Rhyacotriton variegatus* was found at the rate of 0.72 salamanders/hour on parkland compared to 0.05 salamanders/hour on harvested lands (Welsh et al. 2000). In Oregon, densities averaged 0.29 salamanders/m² on forested lands versus 0.04 salamanders/m² on logged habitat (Corn and Bury 1989).

Nature and Degree of Threat

Major threats to this species include timber harvesting, road building, rural development, marijuana cultivation, and climate change. *Rhyacotriton variegatus* is sensitive to the impacts of timber harvesting and roadbuilding due to direct impacts of heavy equipment and indirect effects on temperature, humidity, and sediment load (Welsh et al. 2000, Welsh and Hodgson 2008). Several researchers have argued that declines and extirpations will continue due to timber harvesting and related land management practices (e.g., Welsh et al. 2000, Ashton et al. 2006, Olson et al. 2007, Welsh and Hodgson 2008). While *R. variegatus* can persist in some harvested areas, particularly in coastal forests where the effects of logging may be ameliorated by the milder climate (e.g.,

Welsh 1990, Diller and Wallace 1996; S. Barry, unpublished data), it occurs in more sites and with higher density in older stands.

Habitat loss and degradation due to rural residential development and marijuana cultivation is a growing concern for this species in California. Every new house built in forested lands requires a source of water, which is often provided by diverting headwater streams. In some cases, *R. variegatus* has been observed to occur above but not below such diversions (M. van Hattem, pers. comm.). This threat is likely to increase in the near future. For example, the Humboldt County General Plan is currently being updated, with some proposals considering a doubling or tripling of rural development. Marijuana cultivation also presents a water diversion threat to this species, as well as potential negative impacts due to grading, roadbuilding, and the application of herbicides and pesticides (e.g., Thompson et al. 2014).

Rhyacotriton variegatus has slow developmental times and low vagility, leading to potentially high susceptibility to rapidly changing environmental conditions. Expected climate changes within its range over the next 100 years include increased temperatures, changes in hydrology, changes in fire regime, and vegetation shifts. Mean annual temperatures are expected to increase throughout the range of *R. variegatus* in California (reviewed in PRBO 2011). The frequency of extremely hot days is projected to increase, with roughly 9 additional days over 32.2°C (Bell et al. 2004). Such temperatures exceed the critical thermal maxima for adults and larvae of *R. variegatus*, although water temperatures, microhabitat structure, and behavioral thermoregulation may ameliorate these effects. For coastal populations, upwelling is expected to intensify, which may increase fog development and contribute to cooler, moister conditions (Snyder et al. 2003, Lebassi et al. 2009). Coastal populations may therefore continue to provide more favorable climatic conditions than areas farther inland. Potential changes in precipitation are less clear, with some models predicting modest increases,

others modest decreases, and others reductions in rainfall of up to 28% (reviewed in PRBO 2011). Warmer temperatures will result in less precipitation stored as snow, and reductions of 30–80% are predicted for snowpack accumulation in northwestern California (Snyder et al. 2004, Cayan et al. 2008b). The timing of spring snowmelt has shifted later in the spring in this region over the last 50 years (Stewart et al. 2005), though the timing of future shifts is unknown. Reductions in water availability due to reduced snowpack and possibly reduced precipitation will affect the timing and magnitude of stream flows. This may negatively affect habitat quality and availability for all life stages of this highly aquatic salamander. How fire regime will be affected by climate change in northwestern California is not well understood. Some models predict little change in fire regime or even decreases in area burned along the northern coast (Fried et al. 2004, Lenihan et al. 2008), while increases in area burned have been predicted for the southern coast of northwestern California (Lenihan et al. 2008). Westerling et al. (2011) projected a 100% increase in area burned in northwestern California under some scenarios. How fire affects *R. variegatus* needs further study, although direct mortality and habitat degradation due to fire have been documented in other stream-breeding amphibians (e.g., Gamradt and Kats 1997, Pilliod et al. 2003). Vegetation communities are expected to shift from moist conifer to drier mixed evergreen forest, with reductions in Douglas fir and redwood forest in particular (Lenihan et al. 2008, PRBO 2011). It is unclear what effect these shifts may have on *R. variegatus* because stream conditions and forest age seem to be more important indicators of habitat quality than forest type.

Status Determination

Rhyacotriton variegatus is a Priority 1 Species of Special Concern due to its high degree of habitat specificity resulting in a patchy distribution in isolated habitat islands, high degree of genetic variation among management units,

and association with late-seral forests that are now rare and often ecologically compromised by timber harvesting (Good and Wake 1992, Welsh and Lind 1996).

Management Recommendations

Rhyacotriton variegatus populations would benefit from forest management activities that maintain cold water temperatures and low sedimentation levels such as decreasing the use and building of roads, decreasing timber harvest, and leaving riparian vegetation intact in harvested areas. Suitable microhabitats should be surveyed for *R. variegatus* presence during the wet season when salamanders are more likely to be detected before such areas are disturbed (Tait and Diller 2006, Olson et al. 2007). Monitoring activities themselves can damage sensitive microhabitats (L. Diller, pers. comm.), and personnel should be well trained in techniques to minimize such negative effects. Occupied microhabitats in particular should be protected from direct impacts of heavy equipment. In areas where timber harvest occurs, vegetation should be left intact around *R. variegatus* habitat, particularly to maintain canopy cover, though the width and configuration of such buffers is an important research need detailed below. In the absence of more detailed research, Olson et al. (2007) recommend using relatively wide buffers on the order of 40–150 m to maintain obligate riparian species. In addition to buffers along streams, habitat should be left intact around seeps (“leave islands”; reviewed in Olson et al. 2007). Marijuana cultivation appears to pose a growing threat to maintenance of high-quality habitat for this species. Enforcement and regulation of marijuana cultivation is an ongoing issue in California and we suggest that the environmental impact of such activities be considered. Little is known about use of upland habitats, but protection of large channel networks and associated seeps and springs to maintain aquatic and upland connectivity would likely help maintain populations of *R. variegatus* (Welsh and Lind 1992,

Vesely and McComb 2002, Olson et al. 2007, Welsh 2011).

Monitoring, Research, and Survey Needs

Several studies have been conducted to determine the presence/absence of *Rhyacotriton variegatus* across the landscape, and such surveys should continue. A critical research need is studies that monitor population abundance over time, particularly under different timber harvesting regimes. Given the long life span and slow development time of this species, such long-term studies might provide insights that shorter, single-season analyses would miss. When possible, population estimates in managed forests should be compared to *R. variegatus* abundance in nearby undisturbed mature forest stands (i.e., reference populations) to assess the impacts of disturbance (Welsh 2011). Additional studies on movement ecology and dispersal beyond localized movements would aid in designing management strategies to promote habitat connectivity. The extent to which upland versus aquatic habitats are used for dispersal is unknown and is crucial for determining whether buffers should be focused around continuous waterways, upland linkages between waterways, or both (Welsh and Lind 1992, Olson et al. 2007, Welsh 2011).

Experiments that test the efficacy of buffer strips for maintaining favorable habitat conditions in harvested areas would also be valuable. Buffer strips from 6 to over 90 m wide have been proposed for maintaining riparian fauna under a range of management scenarios (reviewed in Olson et al. 2007). Stoddard and Hayes (2005) recommended buffer strips >46 m wide for *Rhyacotriton*. Similarly, riparian buffer strips 40 m wide around first through third-order streams in Oregon supported similar salamander abundance (including *R. variegatus*) as unharvested stands (Vesely and McComb 2002). Welsh and Hodgson (2008) recommend stream temperatures <15°C to maintain populations. The relationship between the size and aspect of a subbasin, the amount of the surrounding area harvested, the

resulting maximum stream temperature, and how much buffer would be required to ameliorate any critical biological temperature thresholds are important research needs (Welsh et al. 2005). Temperature is not the only factor that can be influenced by management activities however, and other indicators of habitat quality such as embeddedness should be measured as well (Olson et al. 2007).

Because *R. variegatus* is patchily distributed, monitoring studies should first identify areas with suitable habitat. In surveys for *R. variegatus* in Douglas fir/hardwood forests in the Klamath region, Welsh and Lind (1992, 1996) defined minimum essential microhabitat for *R. variegatus* as an area of at least 10 m² of flowing water (e.g., a patch of spring seep or first- or second-order streams) at least 75 m away from a

forest edge. Within these sites, aquatic searches seemed most effective at detecting *R. variegatus*, as they are rarely encountered using techniques such as terrestrial pitfall trapping (e.g., Welsh 1990). Sampling should be done in the spring when *R. variegatus* are most abundant (Welsh and Lind 1992, Ashton et al. 2006, Tait and Diller 2006).

Landscape genetic studies that quantify levels of connectivity within and across stream systems would help to better delimit local management units as well as important dispersal corridors for this species. Studies similar to recent analyses on another western stream salamander, the Idaho giant salamander (*Dicamptodon aterrimus*) (Mullen et al. 2010), would be particularly instructive as a way to examine the relationship between stream connectivity and salamander gene flow.



RED-BELLIED NEWT

Taricha rivularis (Twitty 1935)

Status Summary

Taricha rivularis is a Priority 2 Species of Special Concern, receiving a Total Score/Total Possible of 81% (69/85). During the previous evaluation, *T. rivularis* was determined to not merit Species of Special Concern status (Jennings and Hayes 1994a). *Taricha rivularis* ranked high enough to warrant status in the current evaluation, although very little information is available on population distribution or abundance trends.

Identification

All species in the genus *Taricha* are stocky, medium-to-large newts with granular skin, dark dorsal coloration, and indistinct or absent costal grooves (Petranka 1998, Stebbins 2003). *Taricha rivularis* has bright, tomato red ventral coloration and reaches up to 8 cm SVL (Stebbins 2003). In all members of the genus *Taricha*, breeding males seasonally acquire smooth skin and an enlarged tail fin (Petranka 1998). Larvae have a stream-type-like morphol-

ogy where the tail fin does not extend all the way to the shoulders (Stebbins 2003). The range of *T. rivularis* overlaps with the range of the rough-skinned newt (*T. granulosa*), and the

Red-Bellied Newt: Risk Factors

Ranking Criteria (Maximum Score)	Score
i. Range size (10)	10
ii. Distribution trend (25)	15
iii. Population concentration/ migration (10)	10
iv. Endemism (10)	10
v. Ecological tolerance (10)	10
vi. Population trend (25)	Data deficient
vii. Vulnerability to climate change (10)	7
viii. Projected impacts (10)	7
Total Score	69
Total Possible	85
Total Score/Total Possible	.81




PHOTO ON PREVIOUS PAGE: Red-bellied newt, Mendocino County, California. Courtesy of Adam Clause.

southeastern edge of its range overlaps with the Coast Range newt (*T. torosa*). These species can be distinguished based on several morphological and color characteristics. In addition to distinctive red ventral coloration, *T. rivularis* has dark brown eyes, compared to the yellow or silvery irises in the other species (Twitty 1935).

Taxonomic Relationships

Taricha rivularis was described on the basis of the clear morphological differences existing between it and other California *Taricha* (Twitty 1935), and its species status has never been questioned. Gene flow among populations was previously thought to be very low because animals return to the same stream areas for breeding and show very strong homing behavior (Hedgecock and Ayala 1974, Hedgecock 1978; see the "Life History" section). Kuchta and Tan (2006a) found low levels of allozyme and mitochondrial DNA divergence among four populations in the north and south of the range, which may suggest that gene flow is higher than previously thought. Although *T. rivularis* shows a high degree of philopatry, long-distance movements are well documented, and this may explain the observed low levels of divergence (Kuchta and Tan 2006a).

Life History

Breeding coincides with the receding of streams after heavy winter rains (Twitty 1942). Adults are terrestrial, and the aquatic breeding phase lasts from February to May, with most breeding occurring between March and early April (Twitty 1955, Packer 1960, Twitty 1966, Stebbins 1985). Males typically breed annually, whereas most females breed every 2–3 years (Twitty 1961, Twitty et al. 1964). Adults have been observed returning to the same ~15 m segment of creek to breed across multiple years (Twitty 1959, Packer 1962, Packer 1963, Twitty et al. 1967a). Adults tend to use a small reach of stream during the breeding season, although movements of a couple hundred meters within a season have been observed (Packer 1962). Adults are also capable of moving several kilom-

eters across years and have excellent homing abilities (Twitty 1959, Packer 1962, Twitty et al. 1964, Twitty et al. 1967a). After breeding, adults leave streams but usually remain in the same drainage (Twitty et al. 1967b). Fall rainfall triggers movement, but heavy rainfall can inhibit overland movement (Packer 1960, Grant et al. 1968), and sustained rainfall, increased stream volume, or increased sediment load can stimulate animals to temporarily leave breeding streams (Packer 1960). Little is known about terrestrial habitat use by metamorphs. Underground retreats are used from May to October, and adults forage on the surface before and as they migrate to streams (Twitty 1966, Licht and Brown 1967, Marks and Doyle 2005).

Eggs are attached in a single layer to the bottom of stones or submerged vegetation in fast-flowing water (Twitty 1935, Twitty 1942). The average size of an egg mass is 10 eggs (range 6–16) (Twitty 1935, Riemer 1958, Twitty 1964), and as many as 70 egg masses have been observed attached to a single stone (Twitty 1935, Twitty 1942). The incubation period in the lab is 16–34 days, with faster development times at warmer temperatures (Licht and Brown 1967). Larvae hatch at a minimum of 10 mm TL (Riemer 1958, Twitty 1964) in mid to late April and metamorphose in late August (Licht and Brown 1967) at 45–55 mm TL (Stebbins 1951). There is no evidence that larvae overwinter in streams (Riemer 1958, Twitty 1964). It is unknown how far or to what habitats metamorphs travel, but they go into hiding shortly after metamorphosis (Twitty 1955, Twitty 1961, Twitty 1966, Twitty et al. 1967b). Juveniles are not captured in terrestrial habitats when adults are abundant, suggesting that they remain underground, or at least in a distinct, unknown microhabitat, for several years (Twitty et al. 1967a). It takes approximately 5 years to reach sexual maturity (Licht and Brown 1967). Hedgecock (1978) estimated life spans on the order of 20–30 years based on Twitty's (1966) data, and annual survivorship of adults is probably >90% in most years (Twitty 1961). At one site in Sonoma County, 40% of originally

marked adult animals were still being recaptured 11 years later (Twitty 1966).

Insects and other small invertebrates presumably make up the bulk of the diet of larvae and adults. In one study, adult stomach contents contained exclusively terrestrial organisms (mostly insects), and adults apparently do not feed while in the water during the breeding season (Packer 1961, Licht and Brown 1967).

Habitat Requirements

Taricha rivularis is found in redwood forests along the coast, although other forest types such as Douglas fir, tan oak, and madrone are also used (Marks and Doyle 2005). Aquatic breeding habitats are moderate to fast-flowing mountain streams with rocky bottoms (Twitty 1935, Stebbins 1951). In the Mattole Watershed (northern Mendocino and southern Humboldt Counties), *T. rivularis* was reported to use both steep headwater and 2–4% gradient step-pool reaches, but was most abundant in lower-gradient plane-bed channels (Welsh and Hodgson 2011). Other features of occupied stream habitats were water temperatures ranging between 15°C and 26°C, a mix of coarse streambed substrates, and intermediate levels of canopy closure (Welsh and Hodgson 2011). Unlike other members of the genus, *T. rivularis* rarely breed in ponds or other standing water habitats (Riemer 1958, Stebbins 1985) and seem to avoid streams used by *T. torosa* (Twitty 1942, Twitty 1955). *Taricha rivularis* will breed in the same streams as *T. granulosa* but tend to use faster-flowing reaches (Twitty 1942).

Distribution (Past and Present)

Taricha rivularis is endemic to California and has the smallest geographic distribution among its congeners (Stebbins 2003). The species occurs in coastal northern California in Sonoma, Lake, Mendocino, and southern Humboldt Counties, at elevations from 150 to 450 m (Stebbins 2003, Marks and Doyle 2005). An isolated population is known from the Stevens Creek watershed in Santa Clara County, although it is unclear if this is an introduction or a native population (Reilly et al., in press).

Some habitat has likely been lost to vineyard and other agricultural development in Sonoma and Mendocino Counties, although systematic surveys are not available (H. Welsh, pers. comm.). Some populations have been lost due to damming of creeks and rivers (e.g., Skaggs Spring, which was inundated during the formation of Lake Sonoma). Data from the Mattole Watershed in the mid-1990s documented *T. rivularis* presence in 35% of sampled streams (Welsh et al. 2005), with *T. rivularis* restricted to the forested southern portions of the watershed (Welsh and Hodgson 2011).

Trends in Abundance

Few abundance data are available for this species. Hedgecock (1978) used Twitty's (1961, 1966) census data to estimate that ~60,000 breeding adults occurred along a ~2.5 km stretch of creek in Sonoma County. In the Mattole Watershed, 300 m stretches of randomly selected stream reaches ($n = 83$ stream reaches) yielded 24 metamorphs and 104 aquatic larvae (Welsh and Hodgson 2011).

Nature and Degree of Threat

The paucity of distribution and abundance data makes it difficult to determine the status of most *Taricha rivularis* populations. However, the species has a small range in an area that has experienced high levels of habitat conversion to vineyards and subdivisions, rendering them vulnerable to habitat loss and fragmentation (Marks and Doyle 2005). *Taricha rivularis* may also be experiencing increasing mortality from vehicular traffic (Marks and Doyle 2005), especially during breeding migrations.

Climate change poses potential risks to *T. rivularis* through increased temperatures, changes in hydrology, changes in fire regime, and vegetation shifts. Mean annual temperatures are expected to increase throughout northwestern California (reviewed in PRBO 2011); however, maximum temperature tolerances of *T. rivularis* are unknown. *Taricha rivularis* populations on the coast may be less affected by temperature increases because upwelling is

expected to intensify, potentially leading to increased fog development and cooler, moister conditions (Snyder et al. 2003, Lebassi et al. 2009). Potential changes in precipitation are less clear, with some models predicting little change and others reductions in rainfall of up to 28% (reviewed in PRBO 2011). If conditions become warmer and drier, especially in inland sites, this may restrict terrestrial habitat use and overland dispersal. Changes in precipitation may affect stream hydrology, although how *T. rivularis* will respond to such changes is unknown. How fire regime will be affected by climate change in northwestern California is not well understood. Some models predict little change in fire regime or even decreases in area burned along the northern coast (Fried et al. 2004, Lenihan et al. 2008). Increases in area burned have been predicted for the southern coast of northwestern California and inland areas (Lenihan et al. 2008). Westerling et al. (2011) projected a 100% increase in area burned in northwestern California under some scenarios. How fire impacts *T. rivularis* needs more study, although direct mortality and habitat degradation due to fire has been documented in other stream-breeding amphibians in similar habitats (e.g., Gamradt and Kats 1997, Pilliod et al. 2003). Vegetation communities are expected to shift from moist conifer to drier mixed evergreen forest, with reductions in Douglas fir and redwood forest in particular (Lenihan et al. 2008, PRBO 2011). *Taricha rivularis* may not be severely negatively affected by such shifts, as they use multiple forest types.

Status Determination

Taricha rivularis has a small range in an area that has experienced increased levels of habitat loss and fragmentation in recent decades, resulting in a Priority 2 Species of Special Concern status for this endemic salamander.

Management Recommendations

Given the limited ecological information on this species outside of a handful of sites, it is difficult

to make management recommendations other than protecting known breeding habitats. Disturbances such as timber harvest, roadbuilding and use, housing development, agricultural development, and water diversions should be minimized or eliminated in *Taricha rivularis* habitat. Occupied habitat should be protected, with a focus on protecting the entire stream network (Olson et al. 2007, Welsh 2011). Retaining streamside buffers on managed lands can help mitigate the effects of logging and roadbuilding, but more research is needed to determine buffer prescriptions, particularly how to protect stream network processes (Olson et al. 2007). The ecological effects of buffer protections may vary across habitat types, and narrower buffers may be effective in more mesic coastal habitat compared to more xeric inland sites. One model recommends riparian management zones 40–150 m wide and patch reserves along headwater streams to accommodate upland habitat use and promote connectivity among drainages (Olson et al. 2007). Given the long-range movements documented in this species, large terrestrial habitat patches may be necessary to maintain connectivity among populations. Any efforts to translocate individuals should also take the strong evidence for adult homing behavior into account, as animals are likely to try and return to their original streams. Construction of new roads should be minimized or avoided in areas where protecting *T. rivularis* is a high conservation priority. To reduce the sedimentation impacts of runoff from roads, forest roads should be disconnected from stream systems (e.g., through the use of ditch-relief culverts). Use of heavy equipment should be avoided or restricted on forest roads when larvae are present in nearby aquatic habitat. Road management strategies should be applied to all forest roads, not just those used for timber harvest. In areas that are known to suffer high road mortality, migration barriers and under-road tunnels may reduce vehicular death (e.g., see review in Schmidt and Zumbach 2008), although research is needed into the design and efficacy of such interventions.

Monitoring, Research, and Survey Needs

Surveys to determine the current distribution of occupied breeding habitats are a first step to documenting potential extirpations. Resurveys of Twitty's field sites along Pepperwood Creek, a tributary along the Wheatfield Fork of the Gualala River in northwestern Sonoma County, would be useful for assessing whether population abundance has changed, as this is one of the few areas where demographic data have been collected (e.g., Twitty 1961, Twitty 1966). However, locating the original sites has proven difficult, and they may occur on private lands that are largely inaccessible (S. Kuchta, pers. comm.). Basic ecological research into habitat preferences (both terrestrial and aquatic) are needed as well as demographic data on all life stages (Petranka 1998, Marks and Doyle 2005). Additional research is needed on dispersal,

using both field and genetic techniques. Experiments that moved individuals to different streams found that animals traveled overland to return to their native streams, moving as much as 8 km through terrestrial habitat (Twitty 1959, Twitty et al. 1966). If such terrestrial movements are typical of naturally dispersing animals, then large patches of terrestrial habitat will be needed to maintain connectivity among populations. Finally, although it is assumed that introduced trout and bullfrogs are not a threat to *Taricha* due to their toxic skin secretions, this should be examined for eggs, larvae, and breeding adults. In other California newts, recent experimental research has shown that larval *T. torosa* are highly susceptible to predation by *Ambystoma* (Ryan et al. 2009), and tetrodotoxins have not been isolated from larvae or eggs of *T. granulosa* (Fuhrman 1967).



COAST RANGE NEWT, SOUTHERN POPULATIONS

Taricha torosa (Rathke 1833)

Status Summary

Populations of *Taricha torosa* from the Salinas River in Monterey County south constitute a Priority 2 Species of Special Concern, receiving a Total Score/Total Possible of 66% (73/110). During the previous evaluation, these populations were also considered Species of Special Concern (Jennings and Hayes 1994a).

Identification

Taricha are stocky, medium-to-large newts (up to 8 cm SVL) with granular skin, indistinct or absent costal grooves, and dark dorsal coloration (Petranka 1998, Stebbins 2003). *Taricha torosa* has yellowish brown to dark brown dorsal coloration and pale yellow to orange ventral coloration (Petranka 1998). Adults that enter aquatic habitats for breeding develop smooth skin and a flattened tail while they are in the aquatic habitat, and the tail fin becomes enlarged in males (Stebbins 2003). Larvae are pond type, with large gill filaments and a large fin, and have two dark, irregular longitudinal stripes running down the back (Stebbins 2003).

Taricha torosa is the only newt in southern California but may be confused with other *Taricha* species in northern California, and with the Sierra newt (*T. sierrae*), where the two overlap in Tulare County. All of the characters for distinguishing among *Taricha* can be

Coast Range Newt, Southern Populations: Risk Factors

Ranking Criteria (Maximum Score)	Score
i. Range size (10)	10
ii. Distribution trend (25)	15
iii. Population concentration/migration (10)	10
iv. Endemism (10)	10
v. Ecological tolerance (10)	3
vi. Population trend (25)	15
vii. Vulnerability to climate change (10)	7
viii. Projected impacts (10)	3
Total Score	73
Total Possible	110
Total Score/Total Possible	0.66




PHOTO ON PREVIOUS PAGE: Coast range newt, southern populations, Los Angeles County, California. Courtesy of Adam Clause.

variable, and in some individuals differentiating the species can be difficult. *Taricha torosa* resembles *T. granulosa* but can be distinguished based on the extensive light ventral coloration that reaches the underside of the eyes, eyes that extend beyond the margin of the head when viewed from above, and palatal teeth in the roof of the mouth forming a Y shape (Stebbins 2003). In *T. granulosa*, the dark dorsal coloration extends beneath the eyes, the eyes are more closely inset and do not extend to the margin of the head when viewed from above, and the teeth in the roof of the mouth are in a V-shaped configuration (Stebbins 2003). *Taricha rivularis* has dark eyes (*T. torosa* has yellow in the eyes), a tomato red venter, and dark coloration under the limbs and over the cloaca (Stebbins 2003). *Taricha sierrae* tends to be darker brown dorsally than *T. torosa* and has a burnt or reddish ventral coloration (Stebbins 2003). *Taricha sierrae* also has more of the lighter ventral coloration on its snout and upper eyelids than *T. torosa* (Twitty 1942, Riemer 1958), and these differences in color pattern are intermediate in hybrid populations (Kuchta 2007).

Taxonomic Relationships

Previously, two allopatric subspecies were recognized: *Taricha torosa sierrae* in the Sierra Nevada and *T. t. torosa* in the Coast Range (Riemer 1958). Phylogeographic work has shown that populations in the southern Sierra are *T. t. torosa* (Tan and Wake 1995), and further molecular work has supported elevation to species status for both subspecies (Kuchta and Tan 2006b, Kuchta 2007). There is a contact zone between the two species around the Kaweah River in Tulare County. Kuchta and Tan (2006b) concluded that while newts from San Diego County do not show long-term evolutionary independence, they still constitute a conservation unit due to genetic differentiation, demographic independence, and geographic isolation.

Life History

Terrestrial adults migrate to aquatic breeding habitats such as ponds, streams, and reservoirs

from December to early May, and timing varies by locality, weather, and habitat conditions (Storer 1925, Twitty 1942, Riemer 1958, Gamradt and Kats 1997). Southern populations migrate in March and April (Storer 1925, Brame 1968, Kats et al. 1992) and tend to breed in quiet stream pools (Gamradt and Kats 1996, Gamradt and Kats 1997). No other stream-breeding salamanders occur in the southern part of the range of *Taricha torosa*. Eggs are attached under rocks or to vegetation, with egg masses ranging in size from 7 to 47 eggs (Ritter 1897, Storer 1925, Twitty 1942, Brame 1956, Brame 1968, Mosher et al. 1964). Females may lay 3–6 egg masses at a time, but it is unknown if they breed every year or skip years like *T. rivularis* (Ritter 1897, Twitty 1961, Twitty et al. 1964, Brame 1968). Adults typically leave breeding habitats in early to midsummer (Kats et al. 1994).

Eggs hatch after 4–6 weeks (Kats et al. 1994), and larvae develop for several months, typically metamorphosing in summer or fall (Kuchta 2005). Overwintering has been documented in larvae from Los Angeles (Storer 1925) and Riverside (Carroll et al. 2005) Counties, but given a lack of other reports, this behavior is likely uncommon (Kuchta 2005). Average size at metamorphosis for a Berkeley, Alameda County, population was 47 mm TL, although this probably varies widely depending on local conditions (Ritter 1897). Larvae from a vernal pool in Sonoma County metamorphosed in late July and early August at an average size of 43.8 mm TL (Kuchta 2005). Metamorphosis in permanent water habitats, as are commonly used in the southern part of the range, has not been studied.

Taricha torosa appears to show similar breeding site fidelity, homing ability, and longevity as other *Taricha*, although relatively fewer data are available from *T. torosa*. Watters and Kats (2006) PIT-tagged 36 breeding adults in the Santa Monica Mountains in Los Angeles County in the early 1990s, and recaptured animals for several years. Thirty-nine percent of animals originally tagged were recaptured in subsequent years, some as long as 11 years later,

yielding minimum age estimates of 12–14 years. Animals were recaptured on average 15.5 m from the original capture locality. Terrestrial habitat use is poorly studied in juveniles and adults, although overland movements can be substantial. Trenham (1998) recaptured juveniles up to 3.5 km from their natal ponds. Once adults leave breeding sites, they use mesic microhabitats for aestivation during the dry summer (Stebbins 1951, Trenham 1998).

Larvae presumably eat small invertebrates, detritus, and possibly cannibalize conspecifics (Ritter 1897, Kuchta 2005). Aquatic adults will cannibalize eggs and larvae (Ritter 1897, Kats et al. 1992, Hanson et al. 1994). Terrestrial adults are generalist predators consuming a variety of invertebrate prey and the occasional small vertebrate (Ritter 1897, Hanson et al. 1994, Kerby and Kats 1998).

Habitat Requirements

Northern populations occur in mesic forests in hilly or mountainous terrain, while southern populations occur in drier habitats such as oak, chaparral, and grassland (Riemer 1958). Southern populations tend to use permanent streams for breeding, though recruitment may be higher in seasonal reaches that are free of nonnative predatory fish (E. Ervin, pers. comm.). *Taricha torosa* in southern California are also limited by the availability of rocky canyons with clear, cold water (S. Barry, pers. comm.; R. Fisher, pers. comm.). In the Santa Monica Mountains in Los Angeles County, *T. torosa* using a perennial stream laid 89% of their egg masses in pools and 9.5% in runs (Gamradt and Kats 1997). Rifles were rarely used for oviposition (Gamradt and Kats 1997).

Distribution (Past and Present)

Taricha torosa ranges from central Mendocino County south through the Coast Ranges to San Diego County, and also occurs in the southern Sierra Nevada north to Tulare County, from sea level to 1280 m (Stebbins 1959, Tan and Wake 1995). Species of Special Concern status extends only to those populations found in

Monterey County and farther south, excluding the southern Sierra Nevada isolate. Our map only shows these populations, though we note that it includes museum specimens from the San Bernardino Mountains that have been questioned (E. Ervin, pers. comm.). *Taricha torosa* is restricted to the Santa Ynez Mountains in Santa Barbara County (S. Sweet, pers. comm.). The southernmost populations of *T. torosa* are highly fragmented and occur in the Santa Monica, San Gabriel, and Santa Ana Mountains (Stebbins 2003). Within San Diego County, populations farthest south are geographically isolated from the rest of the range. Jennings and Hayes (1994a) reported these populations as extirpated; however, since then San Diego populations in the Cuyamaca Mountains have been reported to persist in small isolated pockets of 15–20 breeding adults in the Boulder, Ceder, and Conejos Creek systems (E. Ervin, pers. comm. in Kuchta 2005). Surveys in the 1990s of the foothills and mountains around the Central Valley found *Taricha* species (*T. torosa* and *T. granulosa*) absent from more than half of historically occupied counties (Fisher and Shaffer 1996). Jennings and Hayes (1994a) estimated that a third of localities in southern California have been extirpated. Surveys from 2000 to 2002 in the Santa Monica Mountains and Simi Hills in southern California found *T. torosa* present in 43% (15/35) of streams (Riley et al. 2005). *Taricha torosa* tended to be absent from urban streams, and Riley et al. (2005) hypothesized that this was due to effects on habitat quality from artificial flow regimes, increased presence of introduced species, and possibly also collection pressure.

Trends in Abundance

Historically, *Taricha torosa* was noted as common along the Pacific slope (Klauber 1928, Bogert 1930, Klauber 1930, Dixon 1967, Brattstrom 1988), and it may have been one of the most abundant amphibians in California (Jennings and Hayes 1994a). Populations in the upper Carmel Valley adjacent to the Hastings Reservation in Monterey County

numbered in the thousands in the early 1990s but have not been systematically resampled more recently (B. Shaffer and W. Koenig, unpublished data). Southern populations in the Santa Ynez Mountains of Santa Barbara County may have always been small (Jennings and Hayes 1994a). Population size estimates are not available, but populations in the south that used to be in the hundreds are now in the tens (R. Fisher, pers. comm.; E. Ervin, pers. comm., in Kuchta 2005), with populations in San Diego County potentially on the brink of extirpation (S. Kuchta, pers. comm.).

Nature and Degree of Threat

Major threats to *Taricha torosa* include habitat loss and degradation, wildfire, introduced species, and vehicular traffic (Jennings and Hayes 1994a). Sedimentation has caused a large amount of habitat degradation, especially in Los Angeles, Orange, Riverside, and San Diego Counties (Jennings and Hayes 1994a), and *T. torosa* is absent from previously occupied streams in heavily urbanized watersheds (Riley et al. 2005). Wildfire also contributes to habitat degradation. Surveys before and after a chaparral wildfire along a perennial Santa Monica Mountain stream in Los Angeles County documented a roughly 50% reduction in the availability of preferred pool and run habitat due to erosion (Gamradt and Kats 1997). As a result, egg mass density was reduced by two-thirds compared to prefire levels (Gamradt and Kats 1997). Terrestrial adults were observed to produce foamy skin secretions while walking through a prescribed burn area of chamise habitat in Monterey County (Stromberg 1997).

Negative effects of introduced predators on *T. torosa* have been documented. In the Santa Monica Mountains in Los Angeles County, introduced crayfish (*Procambarus clarkii*) and mosquitofish (*Gambusia affinis*) are predators on *T. torosa* and may be contributing to declines (Gamradt and Kats 1996). Stream surveys did not detect either invasive species in the 1980s. Resurveys in the 1990s of previously used breeding habitats found no evidence of breed-

ing in streams with crayfish and mosquitofish present. In one case, *T. torosa* recolonized a reach following floods that removed crayfish, supporting the hypothesis that crayfish exclude newts from breeding habitat. In field and lab trials, survivorship of eggs and larvae was less than 30% in the presence of crayfish. Mosquitofish did not affect egg survivorship but did predate heavily on larvae. Only 46% of larvae survived in the presence of mosquitofish (Gamradt and Kats 1996). Crayfish also aggressively attack and chase adult *T. torosa* out of the water (Gamradt et al. 1997). Native California tiger salamanders (*Ambystoma californiense*) will prey on *T. torosa* larvae where the two co-occur around the Central Valley. However, recruitment is even lower in the presence of hybrids between native *A. californiense* and introduced barred tiger salamanders (*Ambystoma tigrinum mavortium*) (Ryan et al. 2009).

Bd has been documented in 7% (6/90) of *T. torosa* sampled from Santa Clara County (Padgett-Flohr and Longcore 2007), but the role of *Bd* in *T. torosa* declines is unknown. The role of UV radiation in declines is also unknown. Anzalone et al. (1998) reared eggs in field enclosures in the Santa Monica Mountains and found that eggs exposed to UV radiation had 40% survivorship compared to 80% survivorship of eggs when UV was shielded out. However, given that eggs are often attached under rocks and to vegetation, UV is unlikely to be responsible for large-scale declines in the field (Palen and Schindler 2010).

Under climate change, mean annual temperatures are projected to increase throughout the southern range of *T. torosa*, with warmer winters and summers and earlier spring warming expected (reviewed in PRBO 2011). There is less certainty about future precipitation patterns, with estimates ranging from little change to roughly 30% decreases in rainfall (Snyder and Sloan 2005, PRBO 2011). Warmer and potentially drier conditions may affect availability of intermittent and ephemeral waterways used for breeding. Snowpack reductions of up to 90% are predicted in southern California

(Snyder et al. 2004), which will likely result in altered flow regimes. How *T. torosa* may respond to these changes is unknown. The probability and extent of large (>200 ha) fires is expected to increase in the northern part of the special concern range (Fried et al. 2004, Westerling and Bryant 2008). Increases and decreases in fire probability and extent have been predicted for southern California. There is little consensus on future fire dynamics in this part of the range because of the difficulty in modeling Santa Ana weather events (Westerling et al. 2004, Westerling and Bryant 2008). Increases in fire are likely to negatively impact *T. torosa*, largely through habitat degradation but possibly also through direct mortality. Predicted vegetation shifts due to climate change include decreases in chaparral, shrubland, and woodland, and increases in grassland area (Lenihan et al. 2008, PRBO 2011). *Taricha torosa* uses all of these habitat types, and the effects of shifts in their relative abundance and distribution are unknown.

Status Determination

Documented extirpations and reductions in density of remaining populations in southern California, combined with occurrence in an area of high human density, result in a Priority 2 designation for southern populations of *Taricha torosa*.

Management Recommendations

Disturbances such as roadbuilding and road use, housing development, and water diversions should be minimized or eliminated in *Taricha torosa* habitat. Known breeding habitat

should be a high priority for protection. Upland terrestrial habitat also needs to be protected, though the extent and configuration of upland habitat required to maintain population connectivity needs more study. Measures to prevent invasion or remove existing nonnative predators are high-priority activities to stabilize populations of this newt. Road mortality is a clear issue in some areas, particularly south of the Santa Monica Mountains. Road signage has been used to try to reduce road mortality in Monterey County, although its effectiveness is not known. Migration barriers and under-road tunnels may reduce vehicular death in key areas, though research is needed into the design and efficacy of such interventions (Schmidt and Zumbach 2008).

Monitoring, Research, and Survey Needs

Research into terrestrial habitat use and movement is critical for understanding habitat requirements and potential corridors of movement among populations, and these should be undertaken for both stream- and pond-breeding sites. Monitoring of sites where invasive species have been removed should be conducted to determine the long-term efficacy of removals and the recovery time and stability of populations following removal. Genetic analyses at the landscape level could be very informative with respect to both metapopulation dynamics and habitat corridor use and should be conducted in both relatively intact (e.g., Santa Monica Mountains) and more fragmented landscapes. Research is also needed into potential management strategies for dealing with wildfire and erosion control in order to protect breeding habitat.



CALIFORNIA LEGLESS LIZARD

Anniella pulchra Gray 1852

Status Summary

Anniella pulchra is a Priority 2 Species of Special Concern, receiving a Total Score/Total Possible of 55% (61/110). During the previous evaluation, it was also considered a Species of Special Concern (Jennings and Hayes 1994a).

Identification

Anniella pulchra is a medium-sized (11.1–17.8 cm SVL), elongate, legless lizard that is snake-like in body form. This species possesses several characteristics that are related to an underground burrowing lifestyle such as smooth cycloid scales, a shovel-shaped snout, counter-sunk jaw, a short blunt tail, and the absence of external ear openings (Stebbins 2003). The dorsal coloration is generally metallic light silver or golden with a black middorsal line down the length of the body and black lateral stripes. *Anniella pulchra* typically have a lemon-yellow ventral coloration. Faintly striped variants sometimes occur, and dark-brown and black forms

occur on the Monterey peninsula and around Monterey Bay, as well as from Morro Bay, Monterey County, south to Gaudalupe, Santa Barbara County (Stebbins 2003). This species is unlikely to be confused with other lizard species

California Legless Lizard: Risk Factors

Ranking Criteria (Maximum Score)	Score
i. Range size (10)	5
ii. Distribution trend (25)	15
iii. Population concentration/migration (10)	0
iv. Endemism (10)	7
v. Ecological tolerance (10)	7
vi. Population trend (25)	10
vii. Vulnerability to climate change (10)	7
viii. Projected impacts (10)	10
Total Score	61
Total Possible	110
Total Score/Total Possible	0.55




PHOTO ON PREVIOUS PAGE: California legless lizard, Kern County, California. Courtesy of Adam Clause.

in California because it is our only legless lizard. Though *A. pulchra* bears a superficial resemblance to some snake species, the presence of moveable eyelids effectively distinguishes it.

Taxonomic Relationships

Here we treat all California animals as a single species, *Anniella pulchra*. There is substantial evidence for population structure within this species in California from karyotype, allozyme, mitochondrial DNA, nuclear DNA, and morphological studies (e.g., Bezy and Wright 1971, Bezy et al. 1977, Rainey 1985, Pearse and Pogson 2000, Parham and Papenfuss 2009, Papenfuss and Parham 2013). A recent genetic study by Parham and Papenfuss (2009) identified five major lineages within California and documented more extensive genetic diversity within the species than previously reported. Papenfuss and Parham (2013) subsequently proposed that these clades be elevated to species status based on genetic information and some additional data on morphology. This revision occurred as we were finishing our evaluation of special concern status, and we retain the traditional arrangement here to allow the herpetological community time to evaluate this proposed change in taxonomy.

Life History

Breeding occurs between early spring and July in these live-bearing lizards. Oviductal eggs have been observed between July and October, and 1–4 young are born after a 4-month gestation period (Miller 1944, Goldberg and Miller 1985). Juveniles grow rapidly (2.5–4.4 mm SVL/month) and reach sexual maturity after about 2 years at ~9 cm SVL for males and after about 3 years at ~12 cm SVL for females (Miller 1944, Goldberg and Miller 1985). Life span in the field is unknown, but captive animals have survived for almost 6 years (L. Hunt, pers. comm., in Jennings and Hayes 1994a).

Anniella pulchra is rarely seen active on the surface, but they do use the soil/litter interface for feeding and mating (Miller 1944). Daily activity patterns peak in the morning and

evening, though animals have been observed active at night (Miller 1944, Stebbins 1954, Gorman 1957, Bury and Balgooyen 1976, Kuhnz 2000). Coastal and southern populations are likely active year-round, while inland populations (e.g., Sierra Nevada foothills) may enter a period of dormancy during cold months (Banta and Morafka 1968, Zeiner et al. 1988).

Little is known about movement ecology. These fossorial lizards have been found at soil depths from a few to 50 cm below the surface (Miller 1944, Hunt 1984, Kuhnz 2000). Animals have been observed burrowing to a depth of 46 cm in the laboratory (Kuhnz 2000). In one short-term study (~2 months), 10 lizards were recaptured within 10 m of their original capture points (Miller 1944). A two-year PIT tagging study documented an average home range size of 71 m² (Kuhnz 2000).

Anniella pulchra prefers lower temperatures than most other California lizards (~21–28°C in lab trials, Bury and Balgooyen 1976; critical thermal maximum 34°C, Brattstrom 1965), which is consistent with a non-basking fossorial lifestyle. Surface activity by this species is likely limited by both ambient and substrate temperature (Miller 1944).

Little is known about the feeding ecology of this species. *Anniella pulchra* is a generalist sit-and-wait insectivore (Coe and Kunkel 1906, Miller 1944) that eats larval insects (e.g., microlepidopterans and beetles), adult beetles, termites, and spiders (L. Hunt, pers. comm. in Jennings and Hayes 1994a).

Habitat Requirements

At a regional scale, *Anniella pulchra* occurs in sparsely vegetated habitat types including coastal sand dunes, chaparral, pine-oak woodland, desert scrub, open grassland, and riparian areas (Stebbins 2003; S. Sweet, pers. comm.). At local scales, this lizard is a microhabitat specialist requiring sandy or loose loamy substrates conducive to burrowing (Miller 1944, Gorman 1957, Cunningham 1959a, Banta and Morafka 1968). Soils that are not used include gravel-sized substrates and those with greater than

approximately 10% clay content, resulting in absence of this species from serpentine and shale bedrock (S. Sweet, pers. comm.).

At a Monterey County coastal sand dune site, *A. pulchra* used non-compacted, organic-rich soil preferentially and were most abundant in undisturbed soil types, although they were also found in slightly cemented clay-/silt-rich sands (Kuhnz et al. 2005). Plant community structure also contributed to microhabitat suitability, with *A. pulchra* more common around native shrubs such as silver bush lupine, mock heather, and yellow lupine and less common around nonnative grasses, forbs, and iceplant (Kuhnz et al. 2005). In the Mojave Desert, *A. pulchra* can be found in leaf litter under juniper trees (*Juniperus*) (J. Parham and T. Papenfuss, pers. obs.). Soil moisture may also be a limiting factor for this species (Burt 1931, Miller 1944, Bury and Balgooyen 1976). Kuhnz et al. (2005) found more lizards in the low areas between dunes than in other areas, which may be due to water retention.

Distribution (Past and Present)

Most of the range of *Anniella pulchra* occurs in California, from Contra Costa County south through the Coast Ranges, in parts of the San Joaquin Valley, the western edge of the Sierra Nevada Mountains, the western edge of the Mojave Desert, and northern Baja California (Hunt 1983, Jennings and Hayes 1994a). Although most commonly found within 100 km of the coast, *A. pulchra* ranges in elevation from sea level to about 1800 m (Hunt 1983).

Based on the assumption that *A. pulchra* cannot persist in habitat where soil has been disturbed (e.g., plowing, bulldozing), Jennings and Hayes (1994a) estimated that ~20% of historical habitat is no longer suitable. Parham and Papenfuss (2009) noted that several localities they sampled around Bakersfield in the early 2000s no longer existed by the time their study was published. However, some populations have persisted in developed areas, particularly around fence lines, road verges, utility corridors, and gardens (S. Sweet, pers. comm.).

For example, populations that were present in the 1970s were still extant in the 2000s in Fontana, San Bernardino County, in residential areas that were formerly the Delhi Dunes (S. Barry, pers. comm.). *Anniella pulchra* has also been observed in irrigated gardens in Contra Costa County where naturally sandy soils are available (E. Ervin, pers. obs.). The long-term viability of populations in such developed areas is an important research question.

Trends in Abundance

Very few population size estimates are available for this cryptic species. *Anniella pulchra* can be locally abundant, with the highest documented density of 1.67/m² occurring under a single yellow lupine bush in coastal dune habitat at Moss Landing, Monterey County (Kuhnz et al. 2005). Given the high degree of development within its coastal range, we suspect that some populations are declining. In particular, the black form on the Monterey Peninsula may be at great risk given the substantial development pressure in the region.

Nature and Degree of Threat

The greatest threats to *Anniella pulchra* are habitat loss and degradation, and climate change is also a potential emerging threat. Anthropogenic impacts that disturb soil moisture levels or result in soil compaction likely degrade habitat suitability for this species. While some disturbance may be tolerated, development that covers large areas (>8 ha) can potentially cause local extinctions of *A. pulchra* (S. Sweet, pers. comm.). Invasive plants may also have a negative impact on habitat suitability and abundance (Kuhnz et al. 2005). Over the next 100 years, mean annual temperature is expected to increase throughout the range of *A. pulchra* (reviewed in PRBO 2011). There is greater uncertainty in how precipitation will change, with some models predicting decreases in precipitation of up to 37% and other models predicting no change or only moderate declines (Bell et al. 2004, Snyder et al. 2004, Snyder and Sloan 2005, PRBO 2011). Warmer and

drier conditions might limit activity to deeper soil depths, although the population impacts of such a shift are unknown. Alterations in vegetation communities due to climate change may pose a larger threat to this species, as increases in grassland habitat are predicted through much of its range with concomitant decreases in preferred open habitat types such as coastal scrub, particularly in southern California (Lenihan et al. 2008, PRBO 2011). The frequency and size of fires in the Coast Ranges is expected to increase up to 50% by the end of the century (Fried et al. 2004, Lenihan et al. 2008, Westerling and Bryant 2008). Fire dynamics are more difficult to predict in southern California, partly due to the role of Santa Ana winds (reviewed in PRBO 2011, Franco et al. 2011). How fire affects *A. pulchra* is unknown. Direct mortality effects may be small due to its subterranean lifestyle, although indirect negative effects may occur through habitat shifts and changes in soil chemistry.

Status Determination

Anniella pulchra is a near-endemic, ecologically specialized lizard with much of its range occurring in heavily populated and impacted coastal areas. Little data is available on the abundance of this cryptic species, particularly in non-dune habitats, which limits our ability to quantify population trends or document extirpations.

Management Recommendations

Protection of dune areas both along the coast and in the Coast Range is critical. In occupied areas, disturbances such as development, agriculture, and off-highway vehicle use should be reduced or eliminated. Activities that compact soil, in particular, should be avoided. Given that *Anniella pulchra* appears to persist in some developed areas provided that sandy soils and native plant communities remain intact, incentivizing or requiring natural landscaping in low-density housing (as has been done in Monterey County for the federally and state endangered Santa Cruz long-toed salamander, *Ambystoma macrodactylum croceum*) may allow

lizards to coexist with some development. The spread of nonnative plant species into remaining habitat should be minimized. Eradication of invasive plants and restoration of native vegetation may help increase *A. pulchra* density and should be explored.

Monitoring, Research, and Survey Needs

With a few exceptions (e.g., Miller 1944, Kuhnz et al. 2005), little is known about *Anniella pulchra* abundance across its range. Studies of basic ecology are needed in other parts of the range and in other habitat types. Minimally, surveys summarizing habitat use, soil characteristics, and population density in coastal southern California and the southern Sierra Nevada should be conducted to complement work in Monterey County. Understanding under what conditions this species can persist in human-disturbed habitats would be valuable, particularly with respect to soil characteristics and fragmentation that occurs as a consequence of urbanization and agricultural land use. *Anniella pulchra* co-occurs with Argentine ants (*Linepithema humile*) along the coast, but it is unknown whether this introduced species has any substantial impacts on *A. pulchra*.

Presence and abundance of this cryptic species are both difficult to assess, and more research into the best sampling methods for different habitats would be useful for the development of monitoring efforts. In a comparison of survey techniques in dune habitat in Monterey County, Kuhnz et al. (2005) concluded that time-constrained searches were the most reliable method for detecting *A. pulchra* presence across a range of population densities and dune vegetation types. In time-constrained searches, surveyors searched the surface, under dried vegetation or cover objects, and up to 15 cm below the surface. Kuhnz et al. (2005) noted that all survey methods were poor at detecting lizards at low densities of $\sim 1/100 \text{ m}^2$, and even time-constrained searches greatly underestimated density compared to depletion raking (raking of substrate until one or fewer individuals were found per 40 hours of search

effort). However, these results may not apply in general across habitat types. For example, some investigators prefer to use cover objects at inland sites where *A. pulchra* is relatively rare (J. Parham, pers. comm.).

Additional genetic analyses at the population level may be the best way to efficiently determine the effective population size and genetic connectivity of apparently isolated pop-

ulations. Particularly in conjunction with intensive time-constrained surveys, genetic data can be used to measure habitat-specific gene flow, current population size, and changes in population size. We recommend that appropriate genetic markers be developed and that tissues be collected and deposited in appropriate repositories for such analyses.



COASTAL WHIPTAIL

Aspidoscelis tigris stejnegeri (Van Denburgh 1894)

Status Summary

Aspidoscelis tigris stejnegeri is a Priority 2 Species of Special Concern, receiving a Total Score/Total Possible of 54% (59/110). It was not considered a Species of Special Concern during the previous evaluation (Jennings and Hayes 1994a).

Identification

Aspidoscelis tigris stejnegeri is a member of the *A. tigris* species complex, a group of 8–13 species that are all similar in appearance (Grismer 2002, Reeder et al. 2002, Stebbins 2003). This is a large (6–12.7 cm SVL), extremely active, diurnal lizard with a slim body and a long tail. The dorsal ground color is dark, with a series of lighter tan or beige spots forming stripes down the sides. These stripes may be broken and irregular, suggesting a checkered appearance (Stebbins 2003, Lemm 2006). The ventral coloration is whitish to cream with scattered black spotting which sometimes forms longitudinal lines between the scale rows (Stebbins 2003). The dorsal scales are granular, while the ven-

tral scales are relatively large, rectangular plates (Lemm 2006). The scales on the head are also enlarged dorsally and ventrally, forming plates in front of the gular fold (Lemm 2006). In the San Diego area, juveniles develop a distinctive

Coastal Whiptail: Risk Factors

Ranking Criteria (Maximum Score)	Score
i. Range size (10)	10
ii. Distribution trend (25)	20
iii. Population concentration/ migration (10)	0
iv. Endemism (10)	3
v. Ecological tolerance (10)	3
vi. Population trend (25)	10
vii. Vulnerability to climate change (10)	3
viii. Projected impacts (10)	10
Total Score	59
Total Possible	110
Total Score/Total Possible	0.54




PHOTO ON PREVIOUS PAGE: Coastal whiptail, Los Angeles County, California. Courtesy of Robert Hess.

spotted pattern (Stebbins 2003; R. Fisher, pers. comm.).

Within its range, *A. t. stejnegeri* is only likely to be confused with its congener, the orange-throated whiptail (*A. hyperythra*). Both lizards have similar body shapes and scalation, though *A. hyperythra* is usually smaller (5–7.2 cm SVL) and is marked with well-defined light stripes and an intervening dark ground color (Stebbins 2003). In addition, the males of *A. hyperythra* develop a conspicuous bright orange coloration on the throat and underside of the body and juveniles have bright blue on the tail (Stebbins 2003, Lemm 2006).

Taxonomic Relationships

No modern studies of phylogenetics, phylogeography, or species boundaries exist within the *Aspidoscelis tigris* species complex, although the validity of this subspecies has not been questioned. Reeder et al. (2002) presented a phylogenetic analysis of whiptail lizards of the genus *Cnemidophorus* (sensu lato) and showed that the genus, as historically defined, was not monophyletic. To remedy this, they moved North American whiptails to the genus *Aspidoscelis*, an arrangement that is now widely accepted.

Some confusion surrounds the application of the name *A. t. stejnegeri* in the literature. A closely related whiptail occurs as an insular endemic on Isla Cedros, Baja California, Mexico, which most authors refer to as the subspecies *A. t. multiscutata* (previously, *Cnemidophorus tigris multiscutatus*). However, others have treated *A. t. stejnegeri* as a junior synonym of *A. t. multiscutata* and refer both the insular endemic and the coastal southern California forms to this latter name. Thus, some literature referring to the *A. tigris* subspecies in southern California uses *A. t. multiscutata*. This has sometimes led authors to consider the two names to refer to two separate biological taxa that both occur in southern California (Maslin and Secoy 1986). To clarify, there is only a single member of the *A. tigris* complex in coastal

southern California, and its currently accepted name is *A. t. stejnegeri*.

Life History

The life history of *Aspidoscelis tigris stejnegeri* is poorly studied, particularly within its California range, although it is probably similar to other subspecies within the *A. tigris* species complex. This is a diurnally active, wary lizard, which rarely stops moving during its activity period. *Aspidoscelis tigris stejnegeri* is a generalist predator that actively searches for insects, spiders, scorpions, and other small arthropods, including larvae (Grismer 2002, Lemm 2006). Some subspecies in the complex are known to prey upon small lizards, though this has not been documented in *A. t. stejnegeri* to our knowledge. *Aspidoscelis tigris stejnegeri* is a relatively high-temperature specialist that emerges to begin foraging in late morning as the air temperature rises. It can become active as early as mid-March and remain so until early October, although juveniles can remain active into November (Grismer 2002). When active, *A. t. stejnegeri* moves with a distinctive gait, taking a step, halting briefly, then moving again in rapid succession.

Reproduction takes place in spring and summer. Grismer (2002) documented gravid females and courtship behavior in mid-July in Baja California. Courtship may occur earlier in the California populations (Lemm 2006), although few data exist. Hatchlings begin to appear in late July and August in Baja California; again, this may occur earlier in California (Grismer 2002, Lemm 2006).

Habitat Requirements

Aspidoscelis tigris stejnegeri can be found in a wide variety of habitats within the California portion of its range, including coastal sage scrub, chaparral, riparian areas, woodlands, and rocky areas (Lemm 2006). Early observations of this subspecies in California, as well as data from the Baja California portion of the range, indicate that the species prefers sand-

and/or gravel-bottomed habitats and brushy areas associated with washes—habitats that have largely been destroyed by development in southern California (J. Grinnell, pers. comm. reported in Van Denburgh 1922). The species continues to persist outside of these preferred habitats, particularly in open chaparral and coastal sage with a gravelly substrate (Grismer 2002, Cooper and Matthewson 2008), although possibly at reduced densities. *Aspidoscelis tigris stejnegeri* requires large blocks of contiguous habitat and is rarely encountered where development and roads have fragmented the available habitat (Case and Fisher 2001, Brehme 2003, Cooper and Matthewson 2008).

Distribution (Past and Present)

Aspidoscelis tigris stejnegeri was formerly present in California from the southern slopes of the Transverse Ranges south to the United States–Mexico border and east to the Peninsular Ranges (Van Denburgh 1922). In Mexico, it ranges farther south between the coast and the western slopes of the Peninsular Ranges, eventually intergrading with the reddish whiptail (*A. t. rubida*) in the Vizcaino region of the central Baja California peninsula. In California the species occurs from sea level to about 1500 m (Lemm 2006).

The species is apparently extirpated, or nearly so, from large areas of the Los Angeles basin and the San Diego region due to habitat loss. By 1922, the species was already scarce in the vicinity of Pasadena, reportedly as a result of habitat loss due to development (J. Grinnell, pers. comm. reported in Van Denburgh 1922). Further declines have occurred throughout the Los Angeles basin and in coastal San Diego County (Stebbins 2003; R. Fisher, pers. comm.). Much of the inland range is still intact, though increasing wildfires may pose a threat (Rochester et al. 2010).

Trends in Abundance

Few data exist regarding historical abundance of this species, although it is susceptible to habitat fragmentation and development.

Cooper and Matthewson (2008) reported that the species is rarely encountered in small habitat patches and is an indicator species for large blocks of unfragmented coastal sage and chaparral habitat. Grinnell (1908) reported seeing “many of them” along the lower Santa Ana canyon, San Bernardino County, California, in 1905. This area is now heavily modified and does not provide ideal habitat for this taxon. By 1922, the lizard was reportedly “rare” in the vicinity of Pasadena because of habitat fragmentation and loss (J. Grinnell, pers. comm. reported in Van Denburgh 1922), although Bogert (1930) reported it as being moderately common throughout the southern foothills of the San Gabriel Mountains and most of the Santa Monica Mountains. Atsatt (1913) reported that it was frequently encountered throughout several areas of the San Jacinto Mountains, Riverside County, California. Because habitat fragmentation and loss have continued to occur throughout its range, it is reasonable to assume that declines are continuing.

Nature and Degree of Threat

The primary threat facing *Aspidoscelis tigris stejnegeri* is habitat loss and fragmentation due to development. This species occurs in some of the largest population centers within California and requires relatively large habitat blocks, making it particularly susceptible to urbanization. Further, the increasing frequency and intensity of wildfires in southern California may convert large portions of its remaining habitat to suboptimal grassland, causing further declines in range and/or abundance (Lemm 2006, Rochester et al. 2010, R. Fisher, pers. comm.). Projections from several climate models suggest that the frequency and intensity of wildfires in southern California could increase, although these results appear to be strongly dependent on the model that is employed (Cayan et al. 2008b, Franco et al. 2011, PRBO 2011). If this occurs, additional habitat destruction is likely to occur, negatively impacting this species.

Status Determination

Documented and ongoing declines in the distribution of this species, coupled with ongoing suspected declines in abundance, are the primary contributors to this status. *Aspidoscelis tigris stejnegeri* also has a relatively small range in California. Projected impacts from wildfire (which may increase with future climate change) coupled with the above impacts justify a Priority 2 status.

Management Recommendations

Conservation of remaining habitat is essential for the long-term protection of this species. Habitat protection efforts should focus on maintaining large, unfragmented blocks, and this species should be included in large-scale planning efforts like Natural Community Conservation Planning where the process permits. Establishing the minimum size of habitat blocks is a critical research need. Until these data become available, additional fragmentation and degradation should be prevented in habitat patches that currently support this taxon, and corridors of suitable habitat that con-

nect occupied patches should be identified, protected, and/or restored as necessary.

Monitoring, Research, and Survey Needs

Additional data on this taxon's home range size, habitat requirements, and movement ecology are required to determine the minimum patch sizes and maximum amount of fragmentation that can support viable populations. As for many active, wide-ranging species, the effects of road traffic on mortality would be valuable information for future management efforts. Abundance surveys should be conducted in remaining populations of *Aspidoscelis tigris stejnegeri*. Information on abundance should be correlated with the local habitat patch size to better understand the minimum patch size required for population persistence. Further research should examine the effect of moderate habitat fragmentation on existing populations if habitat corridors between patches can be maintained. Given the patchy nature of the species, a landscape genetic approach that quantified both connectivity and effective population sizes of remaining populations would be valuable.



SAN DIEGO BANDED GECKO

Coleonyx variegatus abboti Klauber 1945

Status Summary

Coleonyx variegatus abboti is a Priority 3 Species of Special Concern, receiving a Total Score/Total Possible of 54% (59/110). It was not considered for Species of Special Concern status during the previous evaluation (Jennings and Hayes 1994a).

Identification

Coleonyx variegatus abboti is a small (maximum 5.8 cm SVL) lizard with slender padless toes, moveable eyelids, vertical pupils, and soft skin covered in fine granular scales (Klauber 1945, Grismer 2002, Stebbins 2003). The dorsal ground coloration is variable and ranges from pale yellow to grayish pink. A series of contrasting darker-brown or tan lateral cross-bands extend down the length of the body and are approximately the same width or narrower than the intervening areas of ground coloration. Areas between bands occasionally contain spots of the darker coloration. The head is dark and usually unmarked or only lightly mottled

with a narrow light nuchal crescent extending backward from the eyes (Klauber 1945). The ventral surface is semi-translucent and immaculate white to faint pink. The juvenile pattern is similar to that of adults, but the

San Diego Banded Gecko: Risk Factors

Ranking Criteria (Maximum Score)	Score
i. Range size (10)	10
ii. Distribution trend (25)	15
iii. Population concentration/ migration (10)	0
iv. Endemism (10)	3
v. Ecological tolerance (10)	7
vi. Population trend (25)	10
vii. Vulnerability to climate change (10)	7
viii. Projected impacts (10)	7
Total Score	59
Total Possible	110
Total Score/Total Possible	0.54




PHOTO ON PREVIOUS PAGE: San Diego banded gecko, San Diego County, California. Courtesy of Jeff Lemm.

coloration is often more pronounced and contrasting.

In California, this subspecies is only likely to be confused with other geckos that occur nearby. The closely related desert banded gecko (*C. v. variegatus*) is parapatric with *C. v. abbotti* along the Peninsular Ranges of Southern California and adjacent Baja California, Mexico. The two taxa are best distinguished based on color pattern, locality, and size. Although color and pattern in both subspecies are variable, *C. v. variegatus* generally lacks the nuchal collar, often has wider and less well-defined dark bands, has extensive spotting on the head, and attains larger overall body sizes (up to 7.1 cm) (Klauber 1945, Grismer 2002, Stebbins 2003, Lemm 2006). The two subspecies intergrade across narrow contact zones in Baja California and probably also in southern California, with *C. v. abbotti* occurring on the coastal side of the Peninsular Range mountains and *C. v. variegatus* on the inland side (Klauber 1945; D. Leavitt, pers. comm.). In some areas, animals that are morphologically referable to *C. v. abbotti* are genetically more similar to *C. v. variegatus* (D. Leavitt, unpublished data; see the "Taxonomic Relationships" and "Distribution" sections). The barefoot banded gecko (*C. switaki*) also has a superficially similar appearance but is more rarely encountered. In California, it has only been found in a narrow area of the Peninsular Range. Other geckos in southern California have expanded toe pads and immovable eyelids and are often extreme habitat specialists (Stebbins 2003).

Taxonomic Relationships

Coleonyx variegatus abbotti is a close relative of *C. v. variegatus*. Its initial description was based primarily on coloration, pattern, and scalation (Klauber 1945). Sequence data from seven nuclear DNA markers confirm the distinctiveness of *C. v. abbotti* but also restrict the known range (see the "Distribution" section). The two taxa are not genetically isolated but have an abrupt genetic and morphological contact zone

in Baja California and possibly also in Southern California (D. Leavitt, unpublished data).

Life History

The life history of *Coleonyx variegatus abbotti* is poorly studied, although it is likely similar to that of the better-studied *C. v. variegatus* in many respects. *Coleonyx variegatus abbotti* is active from March until September or October (Lemm 2006). It is nocturnal, emerging from rock crevices and burrows usually within 2 hours following sunset. Like other geckos, it is a predator, presumably taking a variety of small invertebrates, although the diet has not been studied in detail (Kingsbury 1989, Grismer 2002).

Reproduction takes place in late spring. Females lay one or two eggs at a time (Lemm 2006). Other subspecies of *C. variegatus* are known to lay up to three clutches per year between May and September (Stebbins 2003), and this may also occur in *C. v. abbotti*. Juveniles have been found as late as September (Lemm 2006).

Habitat Requirements

Coleonyx variegatus abbotti is restricted to rocky coastal sage and chaparral habitat, usually in areas between 150 and 900 m in elevation (Lemm 2006). Klauber (1945) noted that the subspecies seems to prefer areas with granite outcrops, though it is not restricted to them and has been found in dry rocky riverbeds. Most specimens have been found under cover objects or on roads at night. It is more frequently found under large cap rocks than under the small rock flakes favored by other small lizard species such as the granite night lizard (*Xantusia henshawii*) (Klauber 1945). Extensive pitfall trapping data indicate that *C. v. abbotti* is absent from areas with a high intensity of artificial night lighting (Perry and Fisher 2006; R. Fisher, unpublished data).

Distribution (Past and Present)

Ongoing genetic analyses of the *Coleonyx variegatus* complex are revising our understanding

of *C. v. abbotti*'s distribution, and thus our current concept of its range may change as these studies are completed. Historically, all *Coleonyx* ranging from the United States–Mexico border north along coastal and cismontane Southern California were considered *C. v. abbotti*. However, genetic data indicate that the range is more limited and primarily restricted to San Diego County, with populations farther north belonging to *C. v. variegatus* (D. Leavitt, unpublished data). The extent of the potential intergrade zone between the two subspecies is not yet well understood. In Mexico, *C. v. abbotti* ranges from the border south along coastal and cismontane Baja California to the vicinity of Cataviña, then extends east across the peninsula and south, eventually intergrading with the Peninsular banded gecko (*C. v. peninsulae*) in the Vizcaíno mid-peninsula region.

Geckos have disappeared from much of coastal San Diego County, primarily in areas with high-intensity artificial night lighting (Perry and Fisher 2006). Whether night lighting itself or other habitat changes associated with artificial night lighting drove the declines is not well studied. Development and agricultural impacts have also extirpated geckos from some areas (R. Fisher, pers. comm.).

Trends in Abundance

Few quantitative data on historical or current abundances are available, although *Coleonyx variegatus abbotti* is less frequently encountered than *C. v. variegatus* farther east (Lemm 2006). This was apparently also the case historically. Klauber (1945) specifically noted that *C. v. abbotti* was less common throughout its range than *C. v. variegatus*. Bogert (1930) also reported that the geckos were rare in Los Angeles County, although genetic data suggest these might actually have been *C. v. variegatus*. Pitfall surveys indicate that the subspecies is found at a small number of sites within southern California (7 out of 21 survey areas, 15 individuals in total) compared to lizard species occupying similar habitats (Case and Fisher 2001). However, these surveys were not designed to specifi-

cally target *Coleonyx*, and no historical baseline data exist with which to compare current abundances.

Nature and Degree of Threat

The primary threat facing *Coleonyx variegatus abbotti* is apparently habitat loss due to agricultural and urban development, including deaths from automobile traffic. Some data further suggest that artificial night lighting is correlated with declines, although no causal link has been established. Climate change within its limited range is expected to increase the frequency and intensity of wildfires, which could degrade some currently suitable habitat. Finally, *C. v. abbotti* is encountered relatively rarely even in suitable habitat, which poses significant challenges in monitoring population trends and the impacts of habitat disturbance.

Status Determination

Coleonyx variegatus abbotti has a restricted range in California that falls within an area that is currently experiencing a large amount of development. Some data suggest that the subspecies has disappeared along the coast in a substantial fraction of its range. This, coupled with the ongoing habitat loss due to development and wildfire, could reduce the current distribution further and justifies a Priority 3 Species of Special Concern designation.

Management Recommendations

The most important management priority for *Coleonyx variegatus abbotti* is to protect remaining habitat. Our current understanding of habitat requirements and this taxon's sensitivity to habitat degradation is unfortunately weak, and there is a strong need for additional study before a thorough and informed management strategy can be developed.

Monitoring, Research, and Survey Needs

The relative rarity with which this subspecies is encountered makes the detection of past and ongoing declines difficult. A comparison of survey protocols for this subspecies, including

time-constrained searches and pitfall trapping should be initiated. A goal of this comparison should be to develop a survey protocol that is capable of detecting changing abundances. A mark-recapture study would help determine whether the apparently low population densities currently observed reflect detectability or true population numbers. This should include a power analysis to clarify the trapping intensity needed in order to detect changes of varying magnitude. Surveys should include relatively pristine sites, moderately disturbed habitats, and those with varying degrees of artificial night lighting. Survey data should also be utilized to inform our understanding of habitat preferences, seasonality, and life history in this taxon.

Additional genetic surveys should also be undertaken to further clarify the range limits and genetic differentiation among members of the *Coleonyx variegatus* complex. In particular, contact zones between different subspecies should be further studied in order to develop a clear understanding of the range for both taxa in southern California. Landscape genetic studies would help to inform management in terms of connectivity of remaining populations and potentially help identify habitat corridors. Information from genetics, morphology, and survey data should be integrated to develop a more comprehensive understanding of differentiation between this subspecies and other members of the *C. variegatus* complex.



PANAMINT ALLIGATOR LIZARD

Elgaria panamintina (Stebbins 1958)

Status Summary

Elgaria panamintina is a Priority 3 Species of Special Concern, receiving a Total Score/Total Possible of 44% (48/110). During the previous evaluation, it was also designated as a Species of Special Concern (Jennings and Hayes 1994a).

Identification

Elgaria panamintina is a large (9.2–15.2 cm SVL), slender, elongate lizard with a light yellow-brown or beige dorsum and a series of contrasting brown crossbands extending from the neck down the length of the body and tail (Stebbins 1958, Banta et al. 1996, Stebbins 2003). The ventral surface is light gray or cream, with small dusky markings forming continuous or broken longitudinal lines that run down the center of each scale row (Jennings and Hayes 1994a, Stebbins 2003). The iris is pale yellow (Stebbins 2003). The contrast between the dark crossbands and lighter dorsal coloration is usually more pronounced in juveniles than in

adults. The tail, when intact, is up to twice the length of the body, although shorter broken/regenerated tails are common (Stebbins 2003).

This lizard is unlikely to be confused with any other species within its range. However,

Panamint Alligator Lizard: Risk Factors

Ranking Criteria (Maximum Score)	Score
i. Range size (10)	10
ii. Distribution trend (25)	0
iii. Population concentration/ migration (10)	0
iv. Endemism (10)	10
v. Ecological tolerance (10)	10
vi. Population trend (25)	5
vii. Vulnerability to climate change (10)	3
viii. Projected impacts (10)	10
Total Score	48
Total Possible	110
Total Score/Total Possible	0.44




PHOTO ON PREVIOUS PAGE: Panamint alligator lizard, Inyo County, California. Courtesy of Adam Clause.

two similar congeners occur in much of California: the northern alligator lizard (*E. coerulea*) and the southern alligator lizard (*E. multicarinata*). Neither of these species has the pattern of broad strongly contrasting crossbands down the length of the body. The crossbands are usually interrupted by a longitudinal, middorsal stripe in *E. coerulea* and are much narrower in *E. multicarinata* (Stebbins 2003).

Taxonomic Relationships

Different studies have recovered discordant phylogenetic placements of *Elgaria panamintina*. Good (1988) recovered a sister relationship between *E. panamintina* and the Madrean alligator lizard (*E. kingii*) from Arizona, using a dataset composed of 34 allozyme loci. More recent studies find that *E. panamintina* is nested within *E. multicarinata*, a placement that was supported by both mitochondrial sequence data (Feldman and Spicer 2006) and nuclear sequence data (D. Leavitt et al., unpublished data).

Leavitt et al. (unpublished data) found low levels of variation among populations of *E. panamintina* and no evidence for recent or ongoing gene flow between this species and other *Elgaria* in western North America. The discordance of the allozyme and nuclear sequence data, and therefore the monophyly of *E. multicarinata* with respect to *E. panamintina*, awaits further investigation.

Life History

The life history of *E. panamintina* is poorly understood. The species spends a large amount of time in rock piles and deep vegetation or brush, so it is not commonly observed (Stebbins 1958, Macey and Papenfuss 1991b). We presume that many aspects of *E. panamintina*'s life history are similar to that found in the better-studied *E. multicarinata*, particularly given the recent molecular evidence of their very close relationship.

Elgaria panamintina emerges from hibernation in late winter or spring, with higher-elevation populations becoming active later in the

year. The species is generally diurnal in the spring through midsummer, when it may switch to nocturnal activity or aestivation, presumably as a response to increasing daytime temperatures (Stebbins 1958, Banta 1963, Dixon 1975, Stebbins 2003). Reproduction has not been documented in the wild, although captive animals have been observed copulating in mid-May (Banta and Leviton 1961). *Elgaria multicarinata* enters reproductive condition at this time of year as well (Goldberg 1972), so we assume that reproduction occurs in mid-spring, although the precise timing likely depends on elevation. Goldberg and Beaman (2003) examined sperm formation in museum specimens and concluded that reproduction takes place during the spring. Like *E. multicarinata* (and unlike *E. coerulea*), *E. panamintina* is oviparous. *Elgaria multicarinata* typically lays eggs in early summer, and we assume that *E. panamintina* does as well (Goldberg 1972). The timing of reproductive events in *E. multicarinata* varies among areas, with some populations producing only one clutch a year and others up to three (Burrage 1965, Goldberg 1972). No data on the number of clutches produced per year or incubation times exist for *E. panamintina*, although Goldberg and Beaman (2003) report a clutch size of four eggs from a single museum specimen.

Dietary data are lacking. We presume that *E. panamintina* is likely a generalist predator like *E. multicarinata*. The latter feeds on a wide variety of insects and other small arthropods, including spiders, centipedes, and scorpions, as well as on small vertebrates, including mice, birds, and lizards (including conspecifics) (Cunningham 1956). Observations in captivity found no obvious differences in feeding behavior between *E. panamintina*, *E. multicarinata*, and *E. kingii*, and we tentatively assume that feeding behavior is also similar in the wild (Stebbins 1958).

Elgaria species have a lower thermal tolerance than most sympatric lizards, which may allow them to maintain higher activity levels in the shaded moist habitats in which they are

most commonly found (Cunningham 1956, Stebbins 1958). Predation on *E. panamintina* has not been recorded, though we assume that they are preyed upon by co-distributed lizard-eating snakes (e.g., coachwhips [*Masticophis*] and patch-nosed snakes [*Salvadora*]) and birds (e.g., raptors and roadrunners [*Geococcyx*]).

Habitat Requirements

Elgaria panamintina are most frequently found in rocky canyons in the immediate vicinity of permanent springs and seeps that are patchily distributed across their limited range (Stebbins 1958, Macey and Papenfuss 1991b). The species usually occurs in or adjacent to narrow strips of riparian vegetation immediately below springs and in deep leaf litter and rock piles along the margins of riparian habitat (Stebbins 1958, Macey and Papenfuss 1991b, Jennings and Hayes 1994a). *Elgaria panamintina* was initially thought to be restricted to these areas, but pit-fall trapping surveys have documented their presence in arid areas well away from water (Banta 1963). Few quantitative data are available on the relative frequency of arid versus mesic habitat use, and it seems likely that populations require permanent water for persistence.

Distribution (Past and Present)

Elgaria panamintina occurs in relatively remote regions of the Great Basin in California. Given the difficulty of accessing much of its potential habitat and the limited work on the species to date, it may occur more widely than has so far been recorded. The known range encompasses many of the desert mountain ranges of Inyo and southern Mono Counties, including the Panamint, Inyo, Nelson, Argus, and Coso Mountains, as well as the western slopes of the White Mountains (Macey and Papenfuss 1991b, Banta et al. 1996, La Berteaux and Garlinger 1998). The known elevational range extends from 760 to 2290 m (Dixou 1975, Macey and Papenfuss 1991b, Stebbins 2003).

The species' present-day distribution is likely relictual, resulting from gradual drying of the Great Basin throughout the Pliocene and

Pleistocene. This general drying has presumably isolated the remaining populations around the few remaining water sources (Stebbins 1958, Good 1988).

Trends in Abundance

No data are available regarding current or historical abundance, although habitat degradation due to mining, livestock grazing, and off-highway vehicle use has likely resulted in population declines (Jennings and Hayes 1994a). Given the very sensitive nature of the remaining islands of mesic habitat in the region, surveys of both population size and connectivity via arid habitat occupancy are needed to provide baseline information on current status.

Nature and Degree of Threat

The primary threat to this species is habitat loss or alteration in its already small range. Many of the known localities occur on private land and are vulnerable to mining, livestock grazing, off-highway vehicle use, and/or diversion of the water sources. Climate change could potentially impact this species if changes in hydrology cause springs to dry up or become less regular in their flow regimes.

Status Determination

Elgaria panamintina is a California endemic with a very small range. It primarily occurs in, and is likely dependent upon, uncommon, small patches of mesic habitat that are scattered widely throughout its range. Each habitat patch is sensitive to several potential disturbances, and if local extirpations occur, natural recolonization seems unlikely. Nearly all known localities occur on unprotected land and are subject to further alteration (Jennings and Hayes 1994a). These factors all contribute to a Priority 3 designation.

Management Recommendations

Terrestrial habitat surrounding permanent springs and seeps should be protected from water diversion and destruction or alteration of

riparian vegetation. There may well be conflicts with livestock and large feral mammals since these animals may trample or otherwise disturb the vegetation and leaf litter surrounding desert springs. *Elgaria panamintina* may also occur at additional springs outside of its currently known range; therefore, riparian areas throughout the area should be preserved to the extent possible, even if *E. panamintina* has not yet specifically been documented at them.

Monitoring, Research, and Survey Needs

Surveys should be conducted at additional springs surrounding the known distribution of *Elgaria panamintina*. These surveys should involve pitfall trapping and/or drift fence arrays, in order to increase detection probabilities. A thorough understanding of *E. panamintina*'s habitat requirements would be invaluable in

determining what habitat modifications can be made to riparian areas without negatively impacting the species, as well as identifying suitable areas to focus survey efforts to look for new populations. A key question is the extent to which the species uses arid habitat away from springs, both as corridors for dispersal among springs and as upland habitat. Both drift fence surveys of this habitat and landscape genetic analyses of known spring populations may contribute to greater understanding of habitat use in this species. The lack of basic life history information on *E. panamintina* also needs to be addressed. Mark-recapture surveys would yield important information about population sizes and the extent of migration between springs. This basic information is crucial for any kind of active management and is largely lacking at the present time.



COPE'S LEOPARD LIZARD

Gambelia copeii (Yarrow 1882)

Status Summary

Gambelia copeii is designated as a Species of Special Concern, although we refrain from assigning a priority score due to a paucity of information. This taxon received a Total Score/ Total Possible of 45% (38/85). It was not designated as a Species of Special Concern during the previous evaluation (Jennings and Hayes 1994a).

Identification

Gambelia copeii is a large (maximum 14 cm SVL) lizard, with a robust head and limbs, granular body scales, and a long cylindrical tail (Grismer 2002, Stebbins 2003, Lemm 2006, Mahrtdt et al. 2010). The dorsal coloration is variable across the range, changing from dark brown in the north to light golden brown or tan in the south (Grismer 2002, Mahrtdt et al. 2010). California populations of *G. copeii* form the northern edge of the species' overall range and are dark above with pairs of large, dark paravertebral spots on the dorsal surface that usually fade anteriorly, are almost always absent

from the head, and broaden to form transverse bands on the tail (McGuire 1996, Stebbins 2003, Mahrtdt et al. 2010). In many individuals, a lighter cream-colored transverse bar separates each pair of these spots along the trunk

Cope's Leopard Lizard: Risk Factors

Ranking Criteria (Maximum Score)	Score
i. Range size (10)	10
ii. Distribution trend (25)	15
iii. Population concentration/ migration (10)	0
iv. Endemism (10)	0
v. Ecological tolerance (10)	0
vi. Population trend (25)	Data deficient
vii. Vulnerability to climate change (10)	3
viii. Projected impacts (10)	10
Total Score	38
Total Possible	85
Total Score/Total Possible	0.45




PHOTO ON PREVIOUS PAGE: Cope's leopard lizard, San Diego County, California. Courtesy of Rob Schell Photography.

(Mahrđt et al. 2010). Flecking is generally present on the sides, and females in breeding condition develop bright orange or red spots on the sides and underside of the tail (Stebbins 2003). In addition, there is pronounced sexual size dimorphism, with females averaging 6.5 mm larger in SVL and 1.3 mm in head length than males (Lappin and Swinney 1999, Goldberg et al. 2010).

In California, *G. copeii* is unlikely to be confused with other lizards within its range. However, it is found immediately adjacent to the range of the more widely distributed long-nosed leopard lizard (*G. wislizenii*), within which *G. copeii* appears to be phylogenetically nested (McGuire et al. 2007). *Gambelia wislizenii* populations that are adjacent to *G. copeii* are generally paler, with dorsal coloration ranging from off-white to tan and many moderately sized spots asymmetrically scattered along the dorsal surface (McGuire 1996, Grismer 2002). The spotting in *G. wislizenii* does not fade anteriorly, and small spots generally occur on the head (McGuire 1996, Stebbins 2003, Mahrđt et al. 2010).

Taxonomic Relationships

Though it was described over a century ago, *Gambelia copeii* was not widely recognized as a distinct species until recently. Morphologically and genetically, *G. copeii* is similar to *G. wislizenii*, which led many authors either to consider the two as conspecifics or to recognize them at the subspecific level. McGuire (1996) provided a comprehensive systematic analysis of the Crotophytidae (the family in which *Gambelia* is included) and argued for the recognition of *G. copeii* as a distinct species, based in large part on the presence of a narrow zone of sympatry between the two species in Baja California, Mexico. Following McGuire's monographic review, the species became widely accepted.

Phylogenetically, *G. copeii* appears to form a monophyletic group that is nested within *G. wislizenii* (McGuire et al. 2007), although this result is based on an analysis of mitochondrial data alone and requires further verification.

Rates of potential gene flow and/or hybridization within the zone of sympatry have not been measured.

Life History

Little is known about the natural history of *Gambelia copeii*, and the limited information that is available comes from populations that occur farther south in Baja California, Mexico. We assume that the California populations are similar in most aspects of their life history to populations from the northern regions of Baja California.

Gambelia copeii emerges from hibernation as early as mid-March in northern Baja California, with adults remaining active at least until September (Grismer 2002). The breeding season begins in March or April and lasts at least until July (Fitch 1970, McGuire 1996, Grismer 2002, Goldberg et al. 2010). Grismer (2002) reported a single female in breeding coloration in August at the southern end of the species' range near Todos Santos, Baja California Sur, suggesting that the breeding season could extend much later in the north. Gravid females have been documented in both March and June, providing some evidence that *G. copeii* may produce multiple clutches in optimal years (Fitch 1970, Goldberg et al. 2010). In a sample of 10 museum specimens, the mean clutch size was 5 and did not appear to depend on female body size (Goldberg et al. 2010).

Gambelia copeii is primarily an ambush predator that preys upon other lizards, including whiptail lizards (*Aspidoscelis*), zebra-tailed lizards (*Callisaurus*), and side-blotched lizards (*Uta*), as well as arthropods (McGuire 1996, Grismer 2002).

Habitat Requirements

Gambelia copeii occurs across a wide latitudinal gradient and tolerates a variety of ecological conditions throughout its range. Little published information exists for California populations, although the species appears to prefer open habitat in mixed chaparral and sage scrub (R. Fisher, pers. comm., C. Mahrđt, pers.

comm.). In Baja California, the species occurs across a wider variety of habitat types, although this likely reflects habitat availability throughout the Baja California peninsula rather than specialization of California populations.

In northern Baja California, *G. copeii* occurs on mesas and foothills in scattered patches of chaparral and inland sage scrub with coarse sandy soils (C. Mahrtdt, pers. comm.) and in an increasingly wide variety of habitat types farther south in Baja California (Grismer 2002). *Gambelia copeii* apparently prefers relatively open habitat throughout the diversity of plant communities in which it is found.

Distribution (Past and Present)

In California, *Gambelia copeii* is restricted to an approximately 70 km² area centered around Campo and Potrero Valleys in extreme southern San Diego County (Mahrtdt et al. 2010; C. Mahrtdt, pers. comm.). However, recent field surveys have failed to reconfirm this species at several sites in both Potrero and Campo Valleys, and the species may be locally extirpated at some of these sites particularly along the western edge of its range (R. Fisher, pers. comm.).

Outside of California, *G. copeii* occurs from the California border throughout much of the Baja California peninsula south at least as far as Todos Santos (Grismer 2002). Few data exist on changes in distribution, although agricultural expansion and development in northern Baja California are likely to cause declines (R. Fisher, pers. comm.).

Trends in Abundance

Few data exist regarding historical or present abundance in California. Unpublished pitfall trapping data collected over a 2-year period indicate that the species occurs at very low densities. Between March 1970 and December 1971, pitfall trapping at a 60 × 60 m study site 2.7 km north-east of Cameron Corners, San Diego County, California, yielded many captures of other lizard species in the area but only a single capture of

Gambelia (C. Mahrtdt, unpublished data). A second individual was captured near this site 3 years later (C. Mahrtdt, unpublished data).

Nature and Degree of Threat

The principal threat facing *Gambelia copeii* is habitat loss due to development. The species is able to persist in a wide variety of habitats farther south, so long as the habitat remains relatively open and, presumably, abundant prey (primarily arthropods and smaller lizards) remains available. However, the species occurs at the extreme northern limit of its range in California, so even minor changes in environmental conditions could have large impacts here. Development, including habitat degradation and fragmentation, and climate-change-associated increases in wildfire frequency and intensity have the potential to cause these changes. Invasion of exotic grasses may also lead to further habitat degradation by reducing the availability of open habitat that this species prefers.

Status Determination

Gambelia copeii has an extremely small range in California, which makes it inherently sensitive to any declines. Ongoing habitat loss and potential impacts from climate change may negatively impact the species, but we have relatively few data to assess risk beyond these broad measures of sensitivity, so we refrain from assigning a priority score at this time.

Management Recommendations

Within its very limited California range, remaining large blocks of habitat require protection from further development to prevent future declines. In the absence of information to the contrary, we assume that grazing, wood clearing, and activities that might negatively impact the density of prey (including the presence of feral or pet cats) are all threats to *Gambelia copeii*. Frequent high-intensity wildfire should also be prevented, to the extent possible, within the species range.

Monitoring, Research, and Survey Needs

As no population density data are available, presence/absence surveys followed by mark-recapture monitoring programs should be undertaken throughout the species' range in California to establish baseline information. Loss of habitat across the United States–Mexico border has the potential to isolate the California populations. To begin studying the potential for this to occur, field studies of migration rates and patterns through disturbed and fragmented habitats should be conducted with the


aim of identifying and protecting remaining habitat corridors, as well as characterizing this taxon's sensitivity to various sources of habitat disturbance. Such information will also be useful for developing models of the effects of future climate change scenarios on *Gambelia copeii*. Additional genetic data from nuclear markers should help confirm the species status of this taxon as well as quantify whether, and to what extent, hybridization occurs between it and *G. wislizenii*.



GILA MONSTER

Heloderma suspectum Cope 1869

PHOTOS: (top) Gila monster documented 29 May 1993 in the Kingston Mountains, San Bernardino County, California. Courtesy of Beth Behm. (bottom) Gila monster documented 7 May 2015 in the Mesquite Mountains, San Bernardino County, California. Courtesy of Barrett Scurlock.



GILA MONSTER
Heloderma suspectum



- ▲ Museum Record
- CNDS, BIOS, or Contributor

- Range
- USDA Ecoregion

2,840,000

Status Summary

Heloderma suspectum is a Species of Special Concern, though we refrain from assigning it a priority status due to lack of information. The species received a Total Score/Total Possible of 60% (30/50) and was data deficient for several metrics. During the previous evaluation, it was also considered a Species of Special Concern (Jennings and Hayes 1994a).

Identification

Among California lizards, *Heloderma suspectum* is virtually unmistakable. *Heloderma suspectum* is a large (22.8–35.5 cm SVL) stocky lizard with a dark ground color and distinctive pinkish, orange, or yellow patterning over the trunk and tail that forms bands or a reticulating network. This species possesses distinctive bead-like scales and large, strongly curved claws (Bogert and Martín del Campo 1956, Beck 2005). The ventral coloration is similar to the rest of the body, with alternating black and yellowish or pinkish bands that may form a reticulated pattern (Bogert and Martín del Campo 1956). Within its range, this species could only possibly be confused with the chuck-

walla (*Sauromalus ater*), which sometimes develops a pinkish or yellowish coloration on top of a dark ground color but lacks the banded or reticulate patterning and does not have large, bead-like scales.

Taxonomic Relationships

Heloderma suspectum is one of two extant members of the family Helodermatidae. It is a close relative of the Mexican beaded lizard (*H. horridum*). The description of this species is generally attributed to Cope (1869), although it was actually depicted in print earlier by Baird (1859) using the name *H. horridum*. Cope's (1869) description is a one-paragraph secondhand summary; a far more complete description of the taxon is given by Bogert and Martín del Campo (1956) in their monographic treatment of the family Helodermatidae. The recognition of two species in the genus has not been questioned since the initial description. More recent molecular results confirm the distinctiveness of the two taxa (Douglas et al. 2010).

Two subspecies of *H. suspectum* have been described based on the pattern of reticulation (or lack thereof) in coloration. *Heloderma suspectum suspectum* has a reticulated color pattern, whereas *H. s. cinctum* has a banded pattern that largely lacks reticulations among the bands. A recent genetic survey of intraspecific variation found little evidence supporting these groupings. Additional data are needed to more fully examine intraspecific variation within this species (Douglas et al. 2010). All specimens known from California match the *H. s. cinctum* color pattern, with the single exception of an individual photographed near Piute Springs, San Bernardino County (see the "Distribution" section) (Lovich and Beaman 2007).

Life History

The life history of *Heloderma suspectum* has not been studied in California. Here we use data from other parts of the range (primarily Utah) and cautiously assume that the life history in California is similar.

Gila Monster: Risk Factors

Ranking Criteria (Maximum Score)	Score
i. Range size (10)	10
ii. Distribution trend (25)	Data deficient
iii. Population concentration/migration (10)	0
iv. Endemism (10)	0
v. Ecological tolerance (10)	10
vi. Population trend (25)	Data deficient
vii. Vulnerability to climate change (10)	10
viii. Projected impacts (10)	Data deficient
Total Score	30
Total Possible	50
Total Score/Total Possible	0.60

Heloderma suspectum overwinters in burrows on rocky slopes adjacent to lower-elevation arroyos and bajadas (Beck 1990, Beck 2005). In California, it likely emerges in April or early May. The species spends nearly all of its time in underground burrows (>95% in Utah), emerging rarely to forage for food and to locate mates (Beck 1990). This species is a strict nest predator, preying on the nests of mammals, ground-nesting birds, and reptiles (Hensley 1949, Jones 1983, Beck 1990, reviewed by Beck 2005). *Heloderma suspectum* is venomous, although it is not known to use venom in subduing prey (Beck 2005). Rather the venom probably serves as a predator avoidance mechanism (Beck 2005).

In California, the daily activity pattern is not well characterized. Nocturnal activity has not been reported, although data are lacking. Reproduction likely occurs in April and May, with oviposition occurring shortly thereafter. Elsewhere in the range (Arizona), males leave their burrows and undertake relatively long (~1.6 km) walks to visit other burrows in search of females (Beck 2005). When males encounter each other during this period of activity, prolonged male-male combat may ensue. This behavior entails males entwining one another and attempting to pin one another to the ground (Beck 2005). The time required for eggs to hatch is poorly characterized, although young appear in the spring, which suggests that they overwinter in the burrow before dispersing. Elsewhere in the range, sexual maturity develops in 2–3 years, and adults are probably long-lived (>20 years) (Jennings 1984, Beck 2005). This species appears to be highly susceptible to water loss, which partially explains its relatively sedentary activity patterns (Beck 2005).

Habitat Requirements

Heloderma suspectum occupies a relatively wide variety of desert habitats throughout its range. In California, it is known primarily from a few desert mountain ranges in the eastern Mojave Desert. It inhabits rocky slopes, arroyos, baja-

das, and washes, and is presumably limited on a larger scale by the availability of summer rainfall in the California deserts. Areas that are known to support this species receive a moderate amount of their total annual rainfall during the summer months (24% of the total), which is similar to the pattern in adjacent areas of Arizona that also support this species (39% of total; Lovich and Beaman 2007). On a more local scale, distribution may be controlled by the availability of relatively deep burrows, the presence of food, and availability of riparian or xeroriparian habitat (Lovich and Beaman 2007). Preferences for certain burrow conditions apparently exist but are poorly understood (Beck 2005). Individuals frequently return to specific burrows while leaving others, apparently suitable ones, unoccupied (Beck 2005). Adult Gila monsters are known to return to the same burrows year after year, showing remarkable homing ability and apparent knowledge of the location of many different burrows within their home range (Beck 2005). Too few records exist from the California portion of the range to form a thorough understanding of habitat requirements, although many records are associated with large and relative high mountain ranges as well as with riparian areas (Lovich and Beaman 2007)

Distribution (Past and Present)

Heloderma suspectum ranges from extreme southwestern Utah, through southern Nevada, southwestern Arizona, and south to Sinaloa, Mexico. In California, the species is known from 30 records in the Kingston, Providence, Clark, Piute, and Chocolate Mountain ranges (Bradley and Deacon 1966, De Lisle 1979, Ford 1981, Bicket 1982, De Lisle 1983, Ford 1983, Lovich and Beaman 2007, Ruppert 2010a, Ruppert 2010b, Lovich and Haxel 2011). Lovich and Beaman (2007) reviewed 26 records in California. Four additional records are now known. On 29 May 1993, a single adult *H. suspectum* was photographed on Smith Talc/Kingston Mountain road in the Kingston Mountains, Inyo County, California, approximately 24 km

east of Tecopa (B. Behm, pers. comm.). The photographs show an animal with the banded pattern typical of other animals found in California (we include the clearest photograph here). An additional record comes from Vulcan Mine Road on the western side of the Providence Mountains on 2 May 2009. A natural history class from Cuesta College observed and photographed a single adult moving along the road (Sneed 2009, Ruppert 2010a, Ruppert 2010b). The most recent record that we are aware of from California was documented on 7 May 2015 in the Mesquite Mountains of California. A single adult animal was found resting under the partial shade of a cat's claw plant in a wash running parallel to Kingston Road (B. Scurlock, pers. comm.). Lovich and Haxel (2011) report an additional credible sighting from Black Mountain in the southern Chocolate Mountains that occurred on 30 April 1974 as well as a second record from the same vicinity that is less well substantiated but may be credible. In addition, old records from the vicinity of Blythe, the Lower Colorado River in Imperial County, Chuckwalla Valley, and the Mojave River are in the literature but are less well substantiated than the more recent records (Woodson 1949, Funk 1966, Tinkham 1971, Lovich and Beaman 2007). The species may also occur in a few additional desert mountain ranges in California where records have not yet been recorded. In particular, the New York Mountains are a likely candidate for future records. These mountains lie between the Providence and Piute Mountains, both of which have records and contain what appears to be suitable *Heloderma* habitat. Other large and potentially suitable mountain ranges in the area include the Whipple Mountains, Turtle Mountains, Chemehuevi Mountains, and the Chuckwalla Mountains (Brown and Carmony 1991, Lovich and Beaman 2007).

Trends in Abundance

No data exist on the current or historical abundance of this taxon in California. Elsewhere in the range, the species exists in low densities

(maximum recorded is ~10 individuals/km²) (Beck 1985). Given the paucity of records in California, the species is likely more rare here than in the rest of the range.

Nature and Degree of Threat

The principal threats facing *Heloderma suspectum* in California are its small and extremely patchy distribution, coupled with the probable marginal habitat found in the state and presumed sensitivity to the effects of climate change. Further, we know virtually nothing about the ecology or population status of this species in California, so declines may occur that go undetected.

Status Determination

The almost complete lack of information on this taxon in the state, coupled with a life history that is potentially sensitive to changing climate, justifies designating this taxon as a Species of Special Concern. Because we have virtually no information about the magnitude of threat in this species, we refrain from assigning it a priority at this time.

Management Recommendations

Management recommendations are extremely difficult to formulate other than to protect habitat known to support this species from modification. Activities that might collapse or otherwise destroy burrows, including intense livestock grazing and mining activities, should be avoided in areas suspected of harboring *Heloderma suspectum* populations. Sightings of this infrequently encountered species should be submitted to the California Natural Diversity Database or other natural history databases (e.g., the LACM RASCals project, <http://www.nhm.org/site/activities-programs/citizen-science/rascals>).

Monitoring, Research, and Survey Needs

It may be impossible to study this species in the field in California because it is so rarely encountered. However, opportunities to do so should be pursued. Telemetric data, in particular,

would be difficult to gather because this species is encountered so infrequently, but would also be an important step in enabling the collection of additional information about California populations. We recommend modeling the climate envelope capable of supporting *Heloderma suspectum* to help focus efforts for future surveys. After potential habitat patches have been identified, dawn and dusk surveys during the spring and following summer rain events probably have the best

chance at identifying additional populations. A key priority for future sightings of this species is to collect nonlethal genetic samples that can then be compared to those collected from elsewhere in the range. These tissues will help to clarify intraspecific variation in the species and, if enough samples can eventually be collected, have the potential to supply information about distinctiveness and isolation of populations inhabiting different mountain ranges in the state.



COAST HORNED LIZARD

Phrynosoma blainvillii Gray 1839

Status Summary

Phrynosoma blainvillii is a Priority 2 Species of Special Concern, receiving a Total Score/Total Possible of 49% (54/110). During the previous evaluation, it was also considered a Species of Special Concern under the name *P. coronatum* (see the "Taxonomic Relationships" section) (Jennings and Hayes 1994a).

Identification

Phrynosoma blainvillii has the typical oval, flattened body form of a horned lizard and reaches a maximum SVL of 11.4 cm (Stebbins 2003). It has a row of large horns behind the head, with the two central horns usually longer than the rest and separated at their base. Two rows of large pointed fringe scales run down each side of the body. Large pointed scales also occur on the throat in two or three rows on each side. The dorsum of the body and tail have randomly scattered large, pointed, keeled scales. The general dorsal coloration is tan, yellowish, brown, reddish, or gray, with large dark blotches. Col-

oration can vary within and between populations and with respect to substrate color. Ventral coloration is cream, beige, or yellow, with dusky spotting (Stebbins 2003).

Coast Horned Lizard: Risk Factors

Ranking Criteria (Maximum Score)	Score
i. Range size (10)	0
ii. Distribution trend (25)	20
iii. Population concentration/ migration (10)	0
iv. Endemism (10)	7
v. Ecological tolerance (10)	7
vi. Population trend (25)	10
vii. Vulnerability to climate change (10)	3
viii. Projected impacts (10)	7
Total Score	54
Total Possible	110
Total Score/Total Possible	0.49




PHOTO ON PREVIOUS PAGE: Coast horned lizard, Kern County, California. Courtesy of Nicholas Hess.

Phrynosoma blainvillii may be confused with the desert horned lizard (*P. platyrhinos*) where the ranges of the two species meet in a small region of the southern and eastern part of the range of *P. blainvillii* in California. *Phrynosoma platyrhinos* is easily distinguishable based on a single row of fringe scales down each side of the body, a single row of pointed scales on either side of the throat, and smaller keeled scales on the dorsum.

Taxonomic Relationships

Phrynosoma blainvillii is a member of a species complex that has had a tumultuous taxonomic history, with several species and subspecies recognized by different researchers over time (Klauber 1936, Reeve 1952, Brattstrom 1997). During the previous Species of Special Concern evaluation (Jennings and Hayes 1994a), a single species, *P. coronatum*, was recognized, and California populations were considered as two subspecies: the California coast horned lizard (*P. c. frontale*) and the San Diego coast horned lizard (*P. c. blainvillii*). Recent studies on morphological, ecological, and genetic variation among populations support the recognition of only a single taxon in California, *P. blainvillii*, leading to a revised species-level taxonomy that restricts the species name *P. coronatum* to populations in Baja California Sur, Mexico (Montanucci 2004, Leaché et al. 2009). Three clades have been identified in California based on mitochondrial DNA: northern Baja California, southern California, and northern California (Leaché et al. 2009; see "Distribution" trend). However, two nuclear loci did not distinguish among the clades in California, and ecological and morphological data show substantial overlap among the clades (Montanucci 2004, Leaché et al. 2009). Therefore, we do not recognize any of these clades as conservation units at this time.

Life History

Phrynosoma blainvillii adults are typically active in California from February to November, with peak activity between April and July (Banta and

Morafka 1968, Hager and Brattstrom 1997, Fisher et al. 2002, Alberts et al. 2004, Gerson 2011). Hatchlings are active from mid to late summer into November (Banta and Morafka 1968, Hager 1996, Hager and Brattstrom 1997, Fisher et al. 2002, Alberts et al. 2004). Diurnal activity switches from midday peaks in the spring to more crepuscular activity in summer and early fall (Heath 1965, Hager and Brattstrom 1997).

Most information on reproduction has been collected in the southern part of the range in California. Goldberg (1983) looked at reproductive condition in 164 specimens collected mostly from March to September in Los Angeles, Riverside, San Bernardino, San Diego, Ventura, and Riverside Counties. Reproductive activity occurred from March to June, with females commonly ovipositing in May. Clutch sizes usually average around 11–12 eggs (Stebbins 1954, Howard 1974, Pianka and Parker 1975, Goldberg 1983). Goldberg (1983) reported that a single female appeared to be yolking a second clutch, suggesting the possibility for multiple clutches per year in this species, though how common this may be is unknown. In northern Baja California and southern California, males have spermatozoa present from April until early June (Howard 1974), and oviposition occurs from late May to July with an incubation period of about 60 days (Howard 1974, Pianka and Parker 1975). Montanucci (1968) observed mating in the field as late as May in Merced County. Howard (1974) observed 25 mm SVL hatchlings in late July and early August in northern Baja California. These animals had attained sizes averaging 42 mm SVL by October. First-year males emerged from winter dormancy at ~51 mm SVL. Animals in this population were sexually mature around 75 mm SVL (Howard 1974). Pianka and Parker (1975) reported minimum female SVL at maturity as 73 mm in Baja California and southern California. Goldberg (1983) reported that the smallest mature males were 62 mm SVL, and the smallest females were 73 mm SVL in southern California.

Annual adult survival estimates from radio-tracked animals in Riverside County were roughly twice as high for males as females: males 62% (95% CI 42–81%) and females 34% (95% CI 15–53%) (estimates assume animals of unknown fate are dead; Alberts et al. 2004). Most deaths were due to predation (31% birds, 23% snakes), followed by road mortality (15%), with the rest due to unknown causes (Alberts et al. 2004). Average home range size varied from 1.9 to 4.0 ha across habitat types, with smaller ranges and lower activity levels observed during a drought year (Alberts et al. 2004).

Surface activity is determined partly by temperature. Adults in a Riverside County population had field active body temperatures ranging from 13.3°C to 39.4°C (mean 34.5°C), and hatchlings had a narrower range of temperatures ranging from 21.1°C to 41.1°C (mean 34.4°C) (Alberts et al. 2004). Animals were not active when ground surface temperatures were below 19.4°C or above 57.3°C (Alberts et al. 2004). Gerson (2011) reported capturing lizards when surface temperatures were up to 63°C in a Merced County population. Pianka and Parker (1975) reported a mean field active body temperature for 15 animals of 36.7°C. The critical thermal minima and maxima are –3°C and 46.7°C, respectively (Brattstrom 1965).

Ants can make up 90% of prey items and 45% of prey volume in stomach contents ($n = 214$; Pianka and Parker 1975), although many other insect prey are also consumed depending on availability (Stebbins 1954, Miller and Stebbins 1964, Alberts et al. 2004). About half of the prey found in scat was *Pogonomyrmex* ants (*P. rugosus* and *P. californicus*) (Riverside County; Alberts et al. 2004). Other ant prey and non-ant insects were taken as well. In Merced County, every scat examined contained beetles, but not every scat contained ants, suggesting less reliance on ant prey in this area (M. Gerson, unpublished data). See the “Nature and Degree of Threat” section for effects of nonnative ants.

Habitat Requirements

Phrynosoma blainvillii is found in a variety of habitat types, including sage scrub, dunes, alluvial scrub, annual grassland, chaparral, oak woodland, riparian woodland, Joshua tree woodland, coniferous forest, and saltbush scrub (Grinnell and Grinnell 1907, Klauber 1939, Stebbins 1954, Banta and Morafka 1968, Montanucci 1968, Tollestrup 1981, Hager and Brattstrom 1997). However, microhabitat preferences are much narrower. *Phrynosoma blainvillii* needs loose, fine soils for burrowing, open areas for thermoregulation, and shrub cover for refugia (Jennings and Hayes 1994a). In undisturbed sage scrub habitat in Riverside County, animals preferred leafy plant species with relatively dense foliage for cover, overwintering, and aestivation (Alberts et al. 2004). In the absence of shrubs, *P. blainvillii* may rely instead upon California kangaroo rat (*Dipodomys californicus*) burrows for refugia (Shedd et al. 2011). In a mark-recapture study in San Bernardino and Riverside Counties, Hager and Brattstrom (1997) observed *P. blainvillii* in the open 64% of the time, in the shade of vegetation 14% of the time, next to vegetation 7% of the time, and in rodent burrows 5% of the time.

Pitfall trapping at 21 sites in 4 counties in southern California revealed that within sites, *P. blainvillii* abundance was positively correlated with the presence of organic soils and chaparral vegetation and negatively associated with non-native Argentine ant (*Linepithema humile*) presence (Fisher et al. 2002). At a larger scale, the abundance of *P. blainvillii* between sites was positively associated with the presence of native ants and chaparral vegetation and negatively associated with canopy height. Similar to patterns in abundance, *P. blainvillii* presence was positively associated with sandy soils and chaparral vegetation and negatively associated with Argentine ant presence.

Distribution (Past and Present)

Phrynosoma blainvillii occurs from northern Baja California north along the coast, continuing into

the Central Valley and Coast Range, and east to the Sierra Nevada foothills and the western edge of the Mojave Desert (Leaché et al. 2009). The southern and northern California clades (see the "Taxonomic Relationships" section) roughly correspond in range to the previously recognized subspecies *Phrynosoma coronatum blainvillii* and *P. c. frontale*, respectively. The northern Baja California clade extends from Ensenada, Mexico, north into San Diego County. The southern California clade slightly overlaps with the northern Baja California clade in San Diego County and continues north to the Los Angeles Basin and east to the San Gabriel Mountains and the edge of the Mojave Desert. A third group, the northern California clade, comprises the rest of the range in California, from the Los Angeles basin north through the Central Valley and Coast Ranges.

Historically, this species occurred in California from an isolated record in Shasta County in the north, south along the edges of the Sacramento Valley, through much of the south Coast Ranges, the San Joaquin Valley, the Sierra Nevada foothills, south along the coast to the Mexican border, and throughout the Transverse and Peninsular Ranges, ending along the western edge of the desert slope (Jennings 1988c). Recent field observations in the NAFHA database document this species at Kennedy Meadows in Tulare County; further information about the status here is needed. Jennings and Hayes (1994a) estimated that *P. blainvillii* has disappeared from 35% of its historical range in northern California and from 45% of its historical range in southern California. Remaining populations in the northern end of its range in the Coast Range and in the Sierra Nevada foothills from Butte County to Fresno County are highly disjunct (Jennings and Hayes 1994a; J. Shedd, pers. comm.).

Trends in Abundance

Declines in the early decades of the twentieth century were partly due to collecting for the curio trade in the Los Angeles basin. Jennings (1987) estimated that at least 115,000 *Phryno-*

soma blainvillii were harvested over a 45-year period, with substantial collecting ending around the 1930s. Due to collecting, lizards were noted as being scarce or absent in many areas where they had formerly been abundant (Grinnell and Grinnell 1907, Bryant 1911, Van Denburgh 1922). Agriculture and development has led to declines in more recent decades (see the "Nature and Degree of Threat" section).

It is very difficult to estimate population sizes for horned lizards because their cryptic coloration and behavior make them difficult to detect. In sage scrub habitat in Riverside County, *P. blainvillii* density was estimated as 3–4 adults/km of road transect traveled and 1.7–4.2 adults/ha, with a total of 402 lizards (adults and juveniles) captured over 5 years (Alberts et al. 2004). In Merced County, Gerson (2011) captured 145 individuals (adults and juveniles) on 2.4 ha of transect over an 8-month period, roughly 60 lizards/ha. Lizards were patchily distributed at this site, and transects were purposefully placed in areas with high lizard abundance (M. Gerson, pers. comm.). The sites in both studies experienced controlled burns and grazing and supported a mix of native and introduced plants (Alberts et al. 2004, Gerson 2011).

Nature and Degree of Threat

Major threats to *Phrynosoma blainvillii* include urbanization, agriculture, off-highway vehicles, flood control structures, energy development, and nonnative Argentine ants (Grinnell and Grinnell 1907, Montanucci 1968, Jennings 1987, Jennings and Hayes 1994a; J. Shedd, pers. comm.). These threats may be more pronounced in the southern part of the range (S. Sweet, pers. comm.). Leatherman (1996) observed a single *P. blainvillii* that had apparently died from getting its horns stuck in an erosion control blanket. Introduced Argentine ants have displaced native ant prey over parts of central and southern California and appear to be spreading largely as a commensal with human development (Ward 1987, Holway 1995, Holway 1998). In choice tests, lizards preferred

native ants to Argentine ants, and Argentine ants were not detected in field-collected scat, suggesting that they are not commonly taken as prey (Suarez et al. 2000). In the laboratory, growth rates were lower for animals raised experimentally on Argentine ant diets relative to native diets (Suarez and Case 2002). However, lizards will shift their diets to include more non-ant prey in Argentine ant-invaded areas (Suarez et al. 2000).

The effects of wildfire on *P. blainvillii* are complex and only beginning to be studied. In southern California, capture rates increased by about 30% in chaparral habitat a few years post-fire compared to unburned reference plots (Rochester et al. 2010). No changes were detected in coastal sage scrub habitat, though both habitat types lost substantial vegetative cover. The positive response to fire in chaparral was likely due to the creation of open habitat and the fact that ant prey communities appeared to be unaffected (Rochester et al. 2010). However, the proportion of plots occupied in chaparral habitat decreased in response to fire, possibly due to direct mortality effects of fire. Population increases in burned areas were hypothesized to be due to recolonization from unburned refugia. If so, then the timing and distribution of fire across the landscape would affect how lizards are able to respond and whether the net effect of fire on populations is positive or negative. Additionally, monitoring for this study detected very few *P. blainvillii* in grassland habitats. Because repeated or high-intensity fires can lead to conversion of shrubland to grassland, this also represents a potential threat.

Under climate change, the probability of large (>200 ha) fires and area burned is expected to increase in the northern coastal part of the range and the Sierran foothills, and be largely unchanged in the Central Valley (Fried et al. 2004, Lenihan et al. 2008, Westerling and Bryant 2008). In the southern part of the range where wildfire is common, there is little consensus on future fire dynamics because of the difficulty in modeling Santa Ana weather

events (Westerling et al. 2004, Westerling and Bryant 2008). Land use in the Central Valley is predominantly agricultural; thus, habitat availability is likely to remain low in this area. Elsewhere in the range, large decreases are expected in shrubland with concomitant increases in grassland (Lenihan et al. 2008, PRBO 2011).

Status Determination

Documented extirpations and declines in this species, coupled with a moderate ecological sensitivity, justify a Priority 2 Species of Special Concern status.

Management Recommendations

Protecting remaining populations from further habitat loss and disturbance is the most important management strategy for this species. The presence/absence and abundance of *Phrynosoma blainvillii* appears to be determined by local, rather than regional-scale factors, so management strategies should focus on protecting local populations (Fisher et al. 2002). Because they tend to rely on crypsis rather than speed for protection, they may be particularly sensitive to land uses that increase the likelihood of animals being crushed or killed, including off-highway vehicle use and grazing. Preventing the spread of Argentine ants into *P. blainvillii* habitat is difficult but also important for the persistence of the species. Given that Argentine ants prefer moist microhabitats, xeric landscaping and reducing artificial surface water may be beneficial for native ants and horned lizards in developed areas.

Monitoring, Research, and Survey Needs

Existing populations should be monitored to determine trends in population abundance. An important research question is the extent to which small habitat fragments, on the order of a few hectares or less, can support viable populations of this lizard. Given the high human population density in much of its range, the effects of human commensal predators, including raccoons, skunks, ravens, and domestic cats should be studied, with control measures

implemented as feasible. Continued work on the effects of Argentine ants, including follow-up studies on shifts in lizard diet after their long-term establishment, would provide valuable information on whether *Phrynosoma blainvillii* can adjust to this widespread invasive ant. More research is needed on the effects of introduced plants, which may increase cover, affect native ant prey, and influence thermoregulation and locomotion (Germano et al. 2001, Alberts et al. 2004, Newbold 2005, Rieder et al. 2010). Grazing and fire can have positive effects by maintaining open habitat and negative effects by facilitating the spread of inva-

sives or through direct mortality (Kimball and Schiffman 2003, Alberts et al. 2004, HilleRis-Lambers et al. 2010). The effects of cattle grazing on *P. blainvillii* need more study. Cattle and other grazers may help maintain open habitats that are favorable to *P. blainvillii* but also may degrade habitat through soil compaction. The net effect of grazing and fire as management strategies requires more study and likely needs to be determined at the site scale. The effects of wildfire on *P. blainvillii* should continue to be studied, particularly given the uncertainty concerning future fire dynamics in the southern part of the range.



FLAT-TAILED HORNED LIZARD

Phrynosoma mcallii (Hallowell 1852)

Status Summary

Phrynosoma mcallii is a Priority 2 Species of Special Concern, receiving a Total Score/Total Possible of 57% (63/110). During the previous evaluation, it was also considered a Species of Special Concern (Jennings and Hayes 1994a).

Identification

Like other horned lizards, *Phrynosoma mcallii* has a round body and is dorsoventrally flattened. It is readily distinguished from other horned lizards by a dark middorsal stripe (Smith 1946). The two largest horns behind the head are long and thin, the tail is broad and flat, and two rows of lateral spines run down each side of the body. The limbs are long and thin relative to other horned lizards. The dorsum is cryptically colored, ranging from pale cream to a light rusty brown, and the ventral surface is white and unmarked. Adults can be as large as 8.7 cm SVL (Boundy and Balgooyen 1988, McGrann et al. 2006), but 6.5–8.0 cm SVL is more typical.

Phrynosoma mcallii co-occurs in narrow sympatry with the desert horned lizard (*P. platyrhinos*) along the Salton Trough in California (Stebbins 2003). The two species are easily distinguished because *P. platyrhinos* has a single

Flat-Tailed Horned Lizard: Risk Factors

Ranking Criteria (Maximum Score)	Score
i. Range size (10)	10
ii. Distribution trend (25)	20
iii. Population concentration/ migration (10)	0
iv. Endemism (10)	3
v. Ecological tolerance (10)	10
vi. Population trend (25)	10
vii. Vulnerability to climate change (10)	3
viii. Projected impacts (10)	7
Total Score	63
Total Possible	110
Total Score/Total Possible	0.57




PHOTO ON PREVIOUS PAGE: Flat-tailed horned lizard, Sonora, Mexico. Courtesy of Rob Lovich.

row of lateral spines, shorter horns on the head, and lacks a dark middorsal stripe (Smith 1946, Stebbins 2003). Morphologically intermediate animals thought to be hybrids have been observed near Ocotillo, California (Stebbins 2003), and near Yuma, Arizona (Young 2010).

Taxonomic Relationships

Mulcahy et al. (2006) conducted a phylogeographic study of *Phrynosoma mcallii* and *P. platyrhinos*. They identified management units for *P. mcallii* on either side of the Colorado River. Populations west of the Imperial Valley were historically connected but are now fragmented by human development. The Coachella Valley population, in particular, appears to be highly isolated (Mulcahy et al. 2006).

Life History

Phrynosoma mcallii is generally most active in the summer and inactive during the winter, although there is some flexibility in their winter dormancy behavior. Adult activity in the Coachella Valley in Riverside County peaked from June to August, with little or no activity observed from November to February (Barrows and Allen 2009). At sites in San Diego and Imperial Counties, adults entered hibernation burrows from early October to late December, and smaller animals entered dormancy later than larger animals (Grant and Doherty 2006). The average onset of winter dormancy occurred in mid-November in Imperial County and lasted for an average of 89 days (range 14–138 days), with most animals emerging in mid-February (Muth and Fisher 1992). Radiotelemetry studies have shown that not all individuals enter this distinct period of dormancy (Muth and Fisher 1992, Wone and Beauchamp 2003, Grant and Doherty 2006). Juveniles have been observed surface-active on warm days in December, suggesting that winter dormancy behavior may be more flexible in juveniles compared to adults (Grant and Doherty 2006). Burrows at sites in Imperial and San Diego Counties were 6 cm deep on average (range 2–17; Muth and Fisher 1992, Grant and Doherty 2006). Summer bur-

rows in Yuma, Arizona, were 25–30 cm deep and 70–80 cm long (Young and Young 2000).

Daily activity patterns shift seasonally (Mayhew 1968, Wone and Beauchamp 2003). At Ocotillo Wells State Vehicular Recreation Area (Imperial and San Diego Counties), *P. mcallii* was active throughout the day in spring and fall but showed a bimodal daily activity pattern in the summer (Wone and Beauchamp 2003). *Phrynosoma mcallii* was out in the open during the early morning but retreated under shrub cover by 10:00 a.m. (Wone and Beauchamp 2003). When substrate temperatures exceeded 49°C, lizards entered burrows and reemerged in the evening when substrate temperatures dropped below 47°C (Wone and Beauchamp 2003). Norris (1949) also reported animals retreating between 10:00 and 11:00 a.m. in Riverside County during July. In outdoor enclosures, Heath (1965) observed shade-seeking behavior when body temperatures averaged 40°C and emergence from shade when mean body temperatures were 34.9°C. Brattstrom (1965) recorded lizards at temperatures ranging from 29.3°C to 41.0°C.

Breeding activity has been observed in the field from early May through the end of August (Setser 2004, Barrows and Allen 2009, Young 2010). Adults emerge from winter dormancy in reproductive condition, with testes at maximum size in males and enlarged yolked follicles present in females (Howard 1974). Eggs are laid in burrows dug by the lizards (Setser 2004) and can be deposited from 14 to 90 cm deep, depending on soil moisture (Setser 2004, Young 2010). Clutch sizes range from 2 to 10 eggs, with the average typically around 5 (Norris 1949, Stebbins 1954, Howard 1974, Pianka and Parker 1975, Setser 2004, Young 2010).

Under good conditions, *P. mcallii* can breed early in the season, young can attain adult size rapidly and breed in their first year, and two clutches per season are possible (Howard 1974, Turner and Medica 1982, Muth and Fisher 1992, Barrows and Allen 2009, Young 2010). In multiple-clutch years, the first cohort emerges in late July or early August at 35–37

mm SVL (Howard 1974, Turner and Medica 1982, Muth and Fisher 1992). These hatchlings may be capable of reproducing in their first spring because they can reach near adult sizes before entering winter dormancy (Howard 1974, Muth and Fisher 1992). The second cohort emerges in late August or early September (Howard 1974, Turner and Medica 1982). However, these animals are only ~38 mm SVL in October and may not reach sexual maturity until another season of growth has occurred (Howard 1974, Muth and Fisher 1992). Working in Yuma, Arizona, Young (2010) observed that hatchlings and yearlings did not attain adult size by the following summer under drought conditions, but in wet years animals attained adult size within six months.

Males usually have larger home ranges than females, and home ranges tend to be larger in wet compared to dry years (Wone and Beauchamp 2003, Setser 2004, Young 2010). Radiotelemetry studies at the Ocotillo Wells State Vehicular Recreation Area found average male home range sizes of 1.8–2.4 ha and female home ranges of 0.9–1.3 ha (Wone and Beauchamp 2003, Setser 2004). Setser (2004) observed high site fidelity, with few lizards shifting their range centers outside of the home range used in the previous year. Near Yuma, Arizona, average male home range size varied from 2.5 ha (males) and 1.3 ha (females) in a very dry year to 10.5 ha (males) and 1.9 ha (females) in a very wet year (Young and Young 2000). In wet years at the Yuma, Arizona, site, maximum mean daily movements were 200–700 m, compared to only 50–100 m in drier years (Young 2010).

Survivorship has been measured in a few populations using radiotelemetry and mark-recapture methods. At Ocotillo Wells State Vehicular Recreation Area, adult yearly survivorship was estimated as approximately 50% over a 2-year study period (Setser 2004). This high survivorship rate was attributed to the scarcity of ground squirrel predators, with only 5–8% of radio-tagged lizards lost to predation (Setser 2004). Similarly, adult survivorship over 2 years in Imperial County was approxi-

mately 50%, with half of known mortalities due to ground squirrel predation (Muth and Fisher 1992). In contrast, 39% (21/54) of radio-tagged *P. mcallii* succumbed to predation at the Yuma, Arizona, site in 1 year, with most deaths attributable to ground squirrels (Young 2010). Predation rates were only 10% in another year at this site (Young 2010). Survivorship of hatchlings over their first year was greater than 50% across multiple cohorts in Coachella Valley (Barrows and Allen 2009). Survivorship declined in subsequent years to less than 20% for 2-year-olds and less than 5% for 3-year-olds (Barrows and Allen 2009). Such low survivorship beyond the first year suggests that early maturity and multiple clutches may be key to positive growth of populations in the Coachella Valley (Barrows and Allen 2009).

Phrynosoma mcallii is a dietary specialist on ants, particularly native harvester ant species. Ants typically make up over 90% of prey items in stomach content and scat analyses (Pianka and Parker 1975, Turner and Medica 1982, Young 2010). In 106 specimens examined by Pianka and Parker (1975), 97% of prey items were ants. While at least 11 species of ants have been identified from scats, ants from the genera *Pogonomyrmex* and *Messor* are most commonly taken (Turner and Medica 1982). Near Yuma, Arizona, ants (mostly genus *Pogonomyrmex*) constituted 99% of prey items, with a few beetles taken as well (Young 2010).

Habitat Requirements

In California, *Phrynosoma mcallii* occurs in several Sonoran Desert habitat types, including sandy areas (flats, hills, and valleys), salt flats, badlands, and gravelly areas (Stebbins 2003, Turner and Medica 1982). While they may prefer areas with a layer of fine, wind-blown sand, *P. mcallii* also occur on substrates ranging from hard-packed soils to sand dunes and mud hills (e.g., Beauchamp et al. 1998, Muth and Fisher 1992). For example, at Ocotillo Wells State Vehicular Recreation Area, a site where sandy habitats are highly disturbed by off-highway vehicle use, *P. mcallii* abundance was highest in

sparsely vegetated gravel and mud hills in less-disturbed areas (Beauchamp et al. 1998).

In the Coachella Valley, *P. mcallii* were 2–6 times more abundant on stabilized sand fields than on active dunes (Barrows and Allen 2009) and were not observed in ephemeral sand fields or stable dune habitats (Barrows and Allen 2010). A reduction in windblown sand over the last few decades due to climatic factors and disturbance is thought to be responsible for the apparent absence of *P. mcallii* from ephemeral sand fields, habitats that still support *P. platyrhinos* populations (Barrows and Allen 2010). Lizards selected moderately compacted sands in both stabilized sand fields and active dunes, and this habitat feature may be important for maintaining the integrity of burrows while still being loose enough for digging (Barrows and Allen 2009).

Distribution (Past and Present)

Phrynosoma mcallii is a desert animal with the smallest range of any *Phrynosoma* species that occurs in the United States (Stebbins 2003). It is found from the Coachella Valley in Riverside County south into extreme northeast Baja California and northwest Sonora, Mexico, and east to the extreme southwest corner of Arizona (Stebbins 2003). The species typically occurs below 230 m elevation, but has been found as high as 520 m (FTHL ICC 2003, Rorabaugh and Young 2009, Turner et al. 1980). The currently occupied range is patchily distributed within the historical range. In California, these areas are the Coachella Valley, west of the Salton Sea and the Imperial Valley, and east of the Salton Sea and the Imperial Valley on the west side of the Colorado River (Mulcahy et al. 2006).

The Flat-tailed Horned Lizard Interagency Coordinating Committee estimated that nearly half of the entire range of *P. mcallii* has been altered by human activities, with 39–43% of the historical habitat in the United States converted to agriculture, urban areas, or other uses (reviewed in FTHL ICC 2003). The historical range of *P. mcallii* in California has been esti-

ated at 700,000–900,000 ha, mostly in Imperial County but including parts of eastern San Diego and central Riverside Counties (reviewed in FTHL ICC 2003). Of this historical range, the Flat-tailed Horned Lizard Interagency Coordinating Committee further estimated that 400,000 ha of habitat remain in California (FTHL ICC 2003).

Within the Coachella Valley, Barrows et al. (2008) used niche models to estimate that 83–92% of historically occupied habitat has been lost to development, agriculture, fragmentation, or disruption of windblown sand transport processes. Of the estimated 33,500 ha of historically available suitable habitat, 2600 ha of potential habitat remain in the valley, of which only 1400 ha is currently occupied (Barrows et al. 2008).

Trends in Abundance

Phrynosoma mcallii has long been regarded as a relatively rare species (e.g., Klauber 1939). Density is very difficult to estimate for this cryptic species, and earlier estimates were based on scat counting methods that are no longer thought to be reliable (see the “Monitoring, Research, and Survey Needs” section). Despite these difficulties, dramatic declines have been documented in some areas (Turner and Medina 1982).

Populations of *P. mcallii* appear to naturally fluctuate in abundance, and the drivers of these dynamics are beginning to be explored. In the Coachella Valley, the population declined by about 50% per year during 2002–2005, resulting in an overall decline of 90% (Barrows and Allen 2009). However, in the following 2 years, *P. mcallii* abundance rebounded to half of the 2002 levels (Barrows and Allen 2009). Unlike some other desert species, abundance was not correlated with year-to-year variation in rainfall (Barrows and Allen 2010). Instead, increased rainfall was negatively correlated with the abundance of ant prey and positively associated with increased soil compaction (Barrows and Allen 2009). Other studies have also found associations between *P. mcallii* abundance and

ant abundance (e.g., Turner and Medica 1982, Rorabaugh et al. 1987).

Mark-recapture studies have generated minimum density estimates of approximately 1.1/ha at Ocotillo Wells State Vehicular Recreation Area (Setser 2004) to 6.1/ha in Imperial County (Turner and Medica 1982). Increasingly sophisticated statistical methods for estimating abundance have been employed to compensate for the low abundance and cryptic nature of *P. mcallii*. Grant and Doherty (2007) working in Imperial County estimated densities from 0.41 to 1.55 lizards/ha at different sites, using methods that explicitly account for detection probability (see the "Monitoring, Research, and Survey Needs" section).

Nature and Degree of Threat

Habitat loss and fragmentation due to urban development and agriculture have been the major threats faced by *Phrynosoma mcallii* populations in California, with future threats anticipated due to renewable energy development. *Phrynosoma mcallii* are particularly sensitive to such disturbances because they are ecological specialists, and their ability to recover from population declines through reproductive responses is highly dependent upon favorable environmental conditions.

Phrynosoma mcallii is negatively impacted by fragmentation, and edge effects can extend several hundred meters into undisturbed habitat (Young and Young 2005, Barrows et al. 2006). Based on surveys of lizard tracks, Barrows et al. (2006) found that *P. mcallii* in the Thousand Palms Oasis Preserve in Coachella Valley, Riverside County, experienced negative edge effects along the desert/suburban boundary of the preserve. *Phrynosoma mcallii* were at low abundance within 150 m of the edge compared to farther into the preserve. The mechanism behind the negative effect was hypothesized to be mortality due to roads and subsidized predators such as shrikes and kestrels. Bird predators were positively associated with suburban edge habitats because of increased availability of trees and poles for

perching compared to desert habitat. There was no edge effect on native harvester ant abundance and nonnative ants were not detected, so the edge effect is probably not due to impacts on prey availability (Barrows et al. 2006). In another study in Yuma, Arizona, 90% of shrike-killed *P. mcallii* were within 10 m of a road (Young 2010). Nonnative ants and plants may also pose a threat to *P. mcallii* (see the "Monitoring, Research, and Survey Needs" section). Wind and solar development may be of particular concern in western Imperial County and east of the Imperial Sand Dunes, while geothermal development may threaten populations inside of the Ocotillo Wells State Vehicular Recreation Area (J. Weigand, pers. comm.). Such development may lead to habitat degradation and loss, as well as increased fragmentation.

Off-highway vehicle use has long been suspected of negatively impacting *P. mcallii* populations through direct effects such as mortality and indirect effects on habitat quality. Within protected Management Areas (see the "Management Recommendations" section), off-highway vehicle use is restricted to designated areas (Grant and Doherty 2009). Outside of these protected areas, approximately 100,000 ha of remaining habitat may be subject to off-highway vehicle activity, an area encompassing more than a quarter of remaining habitat in California (Grant and Doherty 2009). Grant and Doherty (2009) experimentally tested the hypothesis that off-highway vehicles crush dormant *P. mcallii* by controlled rides over radio-tagged animals in burrows. None of the animals in their study died or were injured, suggesting that direct effects on animals in burrows may be weak. Direct mortality of surface-active *P. mcallii* due to off-highway vehicle activity has been reported anecdotally from some sites (e.g., Turner and Medica 1982, Muth and Fisher 1992). McGrann et al. (2006) found that lizard body mass, but not density, was higher on sites with low off-highway vehicle impact compared to high-impact areas. The density of ant mounds (i.e., prey) was also

higher in low-impact sites, supporting the possibility of indirect effects of off-highway vehicle use on *P. mcallii* (McGrann et al. 2006).

Aside from increases in temperature, there is little consensus as to how climate change will affect the Sonoran Desert region of California where *P. mcallii* occurs. Mean annual temperatures are expected to increase, with 22 additional extremely hot days per year (where temperatures exceed the long-term 95th percentile) and 10 fewer days below 0°C predicted (Bell et al. 2004). High temperatures may limit surface activity, whereas warmer, shorter winters may increase opportunities for growth and reproduction. Estimates of changes in rainfall range from modest increases in mean annual rainfall up to 45% decreases (reviewed in PRBO 2011). This uncertainty in how precipitation will change makes it difficult to predict how *P. mcallii* will be affected. The effect of rainfall timing and magnitude on *P. mcallii* populations is likely complex, as drought reduces juvenile growth rate and adult movement, but wet years reduce prey abundance (see the “Life History” section). How fire dynamics will change in this area is also highly uncertain (Westerling and Bryant 2008). Little change is expected in vegetation communities (Lenihan et al. 2008, Stralberg et al. 2009).

Status Determination

The specialized diet of *Phrynosoma mcallii*, its low reproductive rates, and small geographic range in a highly fragmented region of California contribute to a Priority 2 Species of Special Concern status.

At the federal level, efforts to secure range-wide protection for *P. mcallii* have been underway for several years, with the species first identified as a candidate for listing under the federal Endangered Species Act in 1982 (reviewed in USFWS 2011a). In 1989, this lizard was rejected for listing under the California Endangered Species Act. Following these efforts, several state and federal agencies comprising the Flat-tailed Horned Lizard Interagency Coordinating Committee signed a voluntary conservation agreement, which resulted in the protec-

tion of management and research areas and a plan for monitoring the species (Foreman 1997). In 2003, the range-wide management strategy was updated, providing reviews of biology, threats, and management recommendations for *P. mcallii* (FTHL ICC 2003). In 2011, *P. mcallii* was again denied federal protection under the Endangered Species Act (USFWS 2011a). In broad terms, the USFWS concluded that the threats to *P. mcallii* that initiated consideration for listing have been largely addressed by management efforts (USFWS 2011a). However, *P. mcallii* populations continue to face a variety of threats throughout their range in California.

Management Recommendations

The main management actions that can support *Phrynosoma mcallii* populations are those that limit habitat disturbance and destruction. Development that leads to habitat conversion or fragmentation should be avoided or limited in *P. mcallii* habitat. Renewable energy projects should consider potential negative impacts on *P. mcallii*. Limiting off-highway vehicle use to the overwintering season when animals are less likely to be surface-active may help limit direct mortality impacts. Roadside barriers and crossing structures should be investigated to reduce road mortality in areas where roads may be barriers to population connectivity. The use of pesticides in or near *P. mcallii* habitat should consider potential negative impacts on native ant prey that are an important determinant of habitat quality for this species. Habitat corridors should be established or maintained to promote connectivity among remaining populations, particularly across the United States–Mexico border. Assisted migration may be important for ensuring gene flow across obstacles such as fences along the United States–Mexico border. Restoration of degraded habitats could include activities such as manipulating soil properties, removing or controlling nonnative plants, and replanting of native plant species that provide food for harvester ants and open habitat for *P. mcallii*.

The Flat-tailed Horned Lizard Interagency Coordinating Committee has implemented a management strategy for *P. mcallii* (FTHL ICC 2003). In California, this strategy includes the establishment of three Management Areas and one Research Area encompassing roughly 170,000 ha in regions of California deemed especially important to the species including the Borrego Badlands, West Mesa, East Mesa, and Ocotillo Wells. The conservation and management of these areas is described in the FTHL ICC (2003) document, and we refer the reader there for additional details. There is currently no management area in the northwestern portion of the range. However, the Coachella Valley Multiple Species Habitat Conservation Plan and Natural Communities Conservation Plan will protect approximately 44% of remaining habitat in Coachella Valley (FTHL ICC 2003).

Monitoring, Research, and Survey Needs

Monitoring *Phrynosoma mcallii* is difficult because this species is cryptic, population abundance fluctuates, and densities are often low. This results in low detection probabilities overall and a high degree of variation in detection probability with respect to different observers, habitats, substrates, and seasons. *Phrynosoma mcallii* is cryptically colored and also exhibits cryptic behavior, tending to freeze and/or bury itself in the sand instead of fleeing (Bryant 1911). Young (2010) observed that radio-tagged individuals in Arizona were almost always motionless when approached, but tracks showed that the animals ran 1–2 m to reach the cover of twigs or vegetation, then froze to avoid detection. Over 25% of the time, fleeing individuals also shuffled into the sand (Young 2010). Such crypsis results in a strong effect of observer experience on survey success (Grant and Doherty 2007).

To deal with these challenges, researchers have tried to use statistical methods to explicitly incorporate detection probability (the probability of seeing lizards if they are present) into mark–recapture estimates of population abun-

dance (e.g., Grant and Doherty 2007, Royle and Young 2008). Detection probabilities ranged from 0.06 to 0.15 (Young 2010) to as high as 0.52 on sandy plots intensively searched by experienced observers (Young and Royle 2005). Range-wide monitoring by members of the FTHL ICC from 2005 to 2012 yielded detection probabilities ranging from 0.15 in the Borrego Badlands to 0.79 in the Yuma Desert (R. Lovich, pers. comm.), and these monitoring efforts are ongoing. Even with increasingly sophisticated mark–recapture analyses, data collection requires substantial effort, and abundance estimates will always be plagued by low detection probabilities. Because of these challenges, distinguishing population declines from natural fluctuations in abundance is difficult, unless declines are severe. As an alternative, Young (2010) recommended monitoring presence/absence over large areas using scat surveys.

Scat counts were commonly used into the 1990s to estimate abundance, but their reliability for measuring density has subsequently been questioned (e.g., Muth and Fisher 1992, Beauchamp et al. 1998). However, scats have been shown to be a good indicator of *P. mcallii* presence, at least in areas where congeners are absent (Young and Royle 2005). If scats are present on a 0.75 ha plot, there is a >99% probability of an observer detecting them within an hour (Young and Royle 2005). Young (2010) proposed that such scat surveys could be useful for delineating occupied habitat across large areas and that monitoring changes in site occupancy over time might be a more viable monitoring strategy than trying to estimate abundance.

In addition to improved monitoring strategies, other research needs include determining the effects of introduced species, the design and efficacy of road-crossing structures, and landscape genetic studies of population connectivity. An additional important research problem is to identify and monitor processes that reduce the abundance of ant prey and/or affect sand compaction (Barrows and Allen 2009). Monitoring for the spread of Argentine

ants, which have been shown to negatively impact *P. blainvillii*, may also be warranted, especially along suburban–desert boundaries. Argentine ants have invaded the Coachella Valley but to date are not known to have moved into *P. mcallii* habitat (Barrows et al. 2006). Fire ants may also pose a threat to *P. mcallii* (J. Weigand, pers. comm.), and their spread and potential impacts should be studied. Nonnative plant species are suspected to negatively impact horned lizards by reducing the availability of open habitat and seed-producing plants and by impacting locomotion (Germano et al. 2001, Newbold 2005, Barrows et al. 2009, Rieder et al. 2010). Introduced plants such as tall-

growing or Sahara mustard (*Brassica tournefortii*) now occur in *P. mcallii* habitat, and the effects of these species require further study (J. Shedd, pers. comm., Barrows 2012). Barrier fences that prevent lizard access to roads have been successful in Yuma, Arizona (e.g., Gardner et al. 2004), and may be beneficial in targeted areas in California. However, more research is needed into crossing structure design and siting to prevent further fragmentation of populations. Finally, a clearer understanding of the extent of habitat fragmentation (using both genetic and mark-recapture methods) and how it affects population viability is an important research need.



COLORADO DESERT FRINGE-TOED LIZARD

Uma notata Baird 1858

Status Summary

Uma notata is a Priority 2 Species of Special Concern, receiving a Total Score/Total Possible of 58% (64/110). During the previous evaluation, it was also considered a Species of Special Concern (Jennings and Hayes 1994a).

Identification

Uma notata is a medium-sized lizard (7.0–12.2 cm SVL) with a moderately flattened body, a countersunk lower jaw, keeled labial scales, projecting row of pointed scales on the toes, eyelids, and ear openings that form a fringe (Cope 1894, Heifetz 1941, Stebbins 1954, Stebbins 2003). The dorsal color pattern consists of light pale yellow to cream ocelli, with dark or reddish centers over a dark ground color (Van Denburgh 1922, Stebbins 1954, Stebbins 2003). These ocelli tend to form broken lengthwise lines at the shoulders (Heifetz 1941). The dark dorsal coloration fades to reddish brown on the head and legs (Van Denburgh 1922). The undersurface is white, with prominent dark ventrolateral spots or bars on

the underside of the tail and narrow diagonal lines on the underside of the throat (Stebbins 2003). An orange or pinkish stripe occurs along the lower flanks and becomes more prominent during the breeding season (Stebbins 1954).

Colorado Desert Fringe-Toed Lizard: Risk Factors

Ranking Criteria (Maximum Score)	Score
i. Range size (10)	10
ii. Distribution trend (25)	15
iii. Population concentration/ migration (10)	0
iv. Endemism (10)	0
v. Ecological tolerance (10)	7
vi. Population trend (25)	15
vii. Vulnerability to climate change (10)	7
viii. Projected impacts (10)	10
Total Score	64
Total Possible	110
Total Score/Total Possible	0.58




PHOTO ON PREVIOUS PAGE: Colorado Desert fringe-toed lizard, Imperial County, California. Courtesy of Adam Clause.

Orange coloration may also be present around the eye.

Uma notata can easily be confused with its congeners in California, the Coachella Valley fringe-toed lizard (*U. inornata*) and the Mojave fringe-toed lizard (*U. scoparia*), although none of these species have overlapping ranges. *Uma inornata* lacks the large and prominent blotches on the ventral surface, although small black spots may be present (Stebbins 2003). *Uma scoparia* usually has narrow lines on the throat that form chevrons and has dorsal ocelli that do not form broken lines on the shoulders (Stebbins 2003). The sympatric zebra-tailed lizard (*Callisaurus draconoides*) also has black bars on the tail, although these form bands that encircle the tail rather than being present only on the underside. *Callisaurus* also lacks fringes on both the toes and the ear openings and has an overall slimmer body shape (Stebbins 2003).

Taxonomic Relationships

The taxonomy of the fringe-toed lizards has been confusing since their original description and remains somewhat controversial. *Uma notata* was initially described from a single preserved juvenile specimen in poor condition (Baird 1858). The initial description of morphology was inadequate to diagnose the taxon and provided details on coloration specific to the poorly preserved specimen ("light pea green, spotted with darker green") and an inaccurate type locality ("Mojave Desert"). An expanded description was later provided by Cope (1894, 1895b), which helped clarify the distinctiveness of the taxon. Heifetz (1941) provided a thorough morphological analysis of the genus and concluded that *U. notata* should be treated as a species separate from the other two California species (*U. inornata* and *U. scoparia*). However, these three species are closely related and their treatment in the literature has shifted between subspecies (of *U. notata*) and full species (Stebbins 1954, Norris 1958, Mayhew 1964a, Mayhew 1964b, Adest 1977, Zalusky et al. 1980).

In addition, some authors recognize two subspecies within *U. notata*. *Uma notata rufop-*

unctata (Cope 1895b) ranges through Arizona and northwestern mainland Mexico, while *U. n. notata* is present only in California. Analyses of mitochondrial data suggest that these two subspecies do not form a monophyletic group. Rather, *U. n. notata* is sister to *U. inornata* to the exclusion of *U. n. rufopunctata* (Wilgenbusch and De Queiroz 2000, Trépanier and Murphy 2001). Trépanier and Murphy (2001) noted that the mitochondrial DNA implied either that *U. inornata* should be considered part of *U. notata* or that *U. n. notata* should be elevated to a full species and that *U. n. rufopunctata* contains two species (one of which is cryptic and had not previously been recognized). They preferred this latter arrangement, although this has not been formally presented to date. Here, we treat *U. notata* as a full species, separate from *U. n. rufopunctata*. Further genetic analyses using multiple independent sequence markers are needed to clarify these species boundaries, as well as the phylogenetic relationships among species and subspecies.

Life History

To the extent that it has been studied, the life history of *U. notata* is essentially identical to that of *U. scoparia*. This species specializes on fine windblown sand habitats and possesses several behavioral, morphological, and physiological adaptations allowing it to do so (see account for *U. scoparia*). This species has a yearly activity cycle that is similar to *U. scoparia*, becoming surface-active as early as February, breeding between April and July, with egg laying in May–July and young appearing in September (Stebbins 1954, Grismer 2002, Stebbins 2003). The two species also exhibit similar daily activity patterns and behavior. They are known to differ in the pattern and cadence of "pushups" used in territorial displays, which may have served as a behavioral isolating mechanism (Carpenter 1963). *Uma notata* has a generalized diet composed of leaves, flowers, seeds, and a variety of small arthropods that is similar to the diet of

U. scoparia (Stebbins 1944). See the account for *U. scoparia* for additional details.

Habitat Requirements

To the extent that they have been studied, habitat requirements are identical to those of *Uma scoparia* and are described in that species' account.

Distribution (Past and Present)

Uma notata ranges from the southeastern corner of California north and west to the Salton Sea and the northeastern corner of San Diego County. Outside of California, it ranges farther south into Baja California, Mexico, to a latitude roughly parallel with the mouth of the Colorado River (Jennings and Hayes 1994a, Grismer 2002). The species' known elevational range extends from 74 m below to 180 m above sea level (Jennings and Hayes 1994a, Stebbins 2003).

Few distributional declines have been documented, although we presume that they have occurred in some areas that have been heavily impacted by off-highway vehicular use, as well as in areas that have experienced heavy development (see the "Trends in Abundance" section). In particular, agricultural development has eliminated habitat in extensive areas around the Salton Sea (Jennings and Hayes 1994a).

Trends in Abundance

Few data regarding historical *Uma notata* population densities exist, although survey data strongly suggest that ongoing declines are occurring in areas that experience off-highway vehicle use. Luckenbach and Bury (1983) conducted surveys in paired plots at the Algodones Dunes (Imperial County, California) that had or had not experienced off-highway vehicle disturbance. *Uma notata* abundance on off-highway vehicle-impacted plots was significantly lower than nonimpacted areas.

Nature and Degree of Threat

Uma notata is experiencing many of the same threats as *U. scoparia*. Habitat loss due to off-highway vehicle damage and habitat destruc-

tion due to human activities is the greatest immediate concern. Luckenbach and Bury (1983) demonstrated major decreases in abundance from off-highway vehicle use due to direct mortality and decreasing vegetation density and quality. Off-highway vehicle use in *Uma* habitat also causes increased rates of tail loss and hearing loss, neither of which are fatal but both of which decrease individual fitness (Brattstrom and Bondello 1983, Luckenbach and Bury 1983). Climate change models for this region predict relatively sharp increases in mean temperature of up to 2°C. The impact of such increases on *U. notata* is not known but should be a high priority for future research. Other threats include increasing predation associated with human commensals and the more general problems associated with reduced population size and fragmentation. See the *U. scoparia* account for additional discussion.

Status Determination

Uma notata specializes on a habitat which is uncommon, patchy, and undergoing significant degradation, and this is the primary justification for this Priority 2 designation. Several populations of this species appear to be stable, and some of the habitat occurs on protected land; thus, a higher-priority designation is not currently justified.

Management Recommendations

The primary management need for *Uma notata* is habitat protection. Protecting sand dune habitat from the impact of off-highway vehicle use alone will significantly increase the probability of long-term survival of this species in California. Habitat conversion for housing, agriculture, and solar/wind energy may all have strongly detrimental effects on *U. notata*, and the limited distribution of the species requires that impacts be reviewed on a project-by-project basis. Over the longer term, increasing temperature and potentially decreased precipitation due to climate change (PREO 2011) could also lead to habitat loss, which may require the development of additional management actions.

Given their strong association with windblown sand habitats, all species of *Uma* may be subject to local extirpations with limited opportunities for natural recolonization, and human-mediated gene flow may be necessary to maintain such populations.

Monitoring, Research, and Survey Needs

The monitoring needs for *Uma notata* are essentially identical to those of *U. scoparia*. Overall, less of *U. notata*'s range occurs on protected land, so these monitoring efforts (and accompanying habitat protection) are needed more urgently for this taxon than for *U. scoparia*. The impact that habitat modification may have on *U. notata* populations is an area in need

of additional study. Two genetic needs are critical. First, the species boundaries of *Uma*, including the distinctiveness of the subspecies of *U. n. rufopunctata* and the resolution of the number and identity of species contained within the genus, require a multi-locus nuclear dataset to complement initial work using mitochondrial DNA (Trépanier and Murphy 2001). Second, landscape genetic analyses quantifying the extent of past and current gene flow among isolated or semi-isolated populations are needed to better understand how to manage landscapes and have the least possible impact on metapopulation dynamics and future population viability.



MOJAVE FRINGE-TOED LIZARD

Uma scoparia Cope 1894

Status Summary

Uma scoparia is a Priority 3 Species of Special Concern, receiving a Total Score/Total Possible of 55% (61/110). During the previous evaluation, it was also considered a Species of Special Concern (Jennings and Hayes 1994a).

Identification

Uma scoparia is a medium-sized lizard (7.0–11.4 cm SVL) with a moderately flattened body, a countersunk lower jaw, keeled labial scales, a projecting row of pointed scales on the toes, eyelids, and ear openings that form a fringe (Cope 1894, Stebbins 1954). The dorsal ground coloration is black and is heavily covered, with a pattern of white or tan ocelli with blackish to reddish centers that do not form lines over the shoulders (Cope 1894, Heifetz 1941, Jennings and Hayes 1994a, Stebbins 2003). This dark coloration fades to brown or tan on the head, limbs, and tail. The light dorsal coloration tends to vary among populations and usually matches the color of the sand in the vicinity (Miller and

Stebbins 1964). The ventral surface is white, with two prominent black spots on either side of the body (some populations have an additional set of preanal spots) and black bars along the underside of the tail (Heifetz 1941). The

Mojave Fringe-Toed Lizard: Risk Factors

Ranking Criteria (Maximum Score)	Score
i. Range size (10)	10
ii. Distribution trend (25)	10
iii. Population concentration/migration (10)	0
iv. Endemism (10)	7
v. Ecological tolerance (10)	7
vi. Population trend (25)	10
vii. Vulnerability to climate change (10)	7
viii. Projected impacts (10)	10
Total Score	61
Total Possible	110
Total Score/Total Possible	0.55




PHOTO ON PREVIOUS PAGE: Mojave fringe-toed lizard, San Bernardino County, California. Courtesy of Luke Mahler.

throat is marked with narrow crescent-shaped black bars (Cope 1895b, Heifetz 1941, Stebbins 2003). During the breeding season, a yellow-green wash may develop on the ventral surface and fade into pink on the sides (Stebbins 2003).

This species could be confused with its congeners, the Coachella Valley fringe-toed lizard (*U. inornata*) and the Colorado Desert fringe-toed lizard (*U. notata*). *Uma inornata* has greatly reduced, or lacks altogether, the conspicuous black spots on the sides of the belly and has ocelli that tend to form lines over the shoulders. *Uma notata* usually has diagonal lines on the throat rather than crescent-shaped lines and has ocelli that tend to form lines over the shoulders (Stebbins 2003). These three species do not overlap in range, although *U. scoparia* is broadly sympatric with the zebra-tailed lizard (*C. draconoides*), with which it also might be confused. *Callisaurus draconoides* lacks fringe scales on the ear openings and toes, has an overall slimmer body shape, and has black bands that form rings around the distal portion of the tail rather than only being on the tail underside (Stebbins 2003).

Taxonomic Relationships

Uma scoparia was initially described on the basis of femoral pore counts and several scalation characters (Cope 1894, Cope 1895b). It was later placed in synonymy with *U. notata* when several of Cope's diagnostic characters were reinterpreted as representing individual variation rather than species differences (Camp 1916b, Van Denburgh 1922). The taxon was later resurrected to full species status based on a larger series of specimens that identified diagnostic morphological differences among the taxa (Heifetz 1941). Several different authors have noted external morphological, osteological, and genetic similarity among members of the genus and have variously treated *U. scoparia* as a full species or subspecies of *U. notata* (Stebbins 1954, Norris 1958, Mayhew 1964a, Mayhew 1964b, Adest 1977, Zalusky et al. 1980). Carpenter (1963) showed

that the pattern of push-up behavior used in territorial displays was distinct in *U. scoparia*, compared to *U. inornata* and *U. notata*, and suggested that this may serve as an isolating mechanism.

Phylogenetic analyses of mitochondrial data suggested that *U. scoparia* is monophyletic (Trépanier and Murphy 2001, Murphy et al. 2006) and forms a clade with the other Mojave and Sonoran Desert taxa (*U. inornata* and *U. notata*) (Wilgenbusch and De Queiroz 2000). Mitochondrial data also suggest that some haplotype diversity occurs within the *U. scoparia* (Murphy et al. 2006), although divergences are low and additional, multigene nuclear data are needed to clarify intraspecific variation. Populations occurring in the northern part of the range have been proposed as a distinct population segment based on mitochondrial phylogeography and presumed isolation (Murphy et al. 2006).

Life History

Uma scoparia is an active, wary, diurnal lizard that specializes on fine windblown sand habitat. It is extremely similar in most aspects of life history to other species in the genus (Stebbins 1944), and here we make use of life history information from these other species when it is not available for *U. scoparia*. Species in the genus *Uma* all possess a number of morphological, behavioral, and physiological adaptations that allow them to persist in arid habitats. Specifically, a countersunk lower jaw, nasal valves, and fringes on the eyes and ear openings allow *U. scoparia* to prevent sand from entering the body (Norris 1958). The nasal passages have a complex convoluted shape that reduces moisture loss and excludes sand from inhalation (Stebbins 1943, Stebbins 1948). Enlarged fringes on the toes have been experimentally shown to increase both maximum velocity and acceleration on fine sand, particularly on steeply sloped landscapes such as are often found in sand dunes (Carothers 1986). The flattened body form, wedge-shaped head, enlarged, keeled scales on the head, limbs and

toes, and the smooth granular scales over the rest of the body aid in burrowing and "sand-swimming" behavior (Stebbins 1944). *Uma scoparia* employs this behavior both to escape from predators and to take refuge from extremely hot surface conditions (typically when surface temperature exceeds 43°C; Norris 1958). *Uma scoparia* possesses both acute vision and hearing, which aid in predator avoidance and prey capture (Stebbins 1944).

Adult *U. scoparia* overwinter in the sand between November and February, then become surface-active throughout the day as temperatures allow. The species maintains a mean body temperature of 36–37.5°C, often becoming inactive during the hottest part of the day during midsummer (Mayhew 1964b, Miller and Stebbins 1964). Breeding occurs throughout the spring and summer between April and July, and females lay clutches of 1–5 eggs (usually 2 or 3); more than one clutch may be produced in optimal years (Stebbins 1954, Mayhew 1966, Fromer et al. 1983, Stebbins 2003). Young begin to appear on the surface in September (Miller and Stebbins 1964).

Uma scoparia has a generalized diet that includes a variety of beetles, ants, wasps, flies, and other small arthropods, as well as plant leaves and seeds (Stebbins 1944). At Dale Dry Lake, San Bernardino County, the diet of adult *U. scoparia* consisted of approximately 60% plant material (mainly in the form of small seeds) and 40% small arthropods (Minnich and Shoemaker 1972). The juvenile diet, conversely, was composed of over 90% arthropods (Minnich and Shoemaker 1972). In low rainfall years, adults may be forced to switch to a diet composed mostly of arthropods due to lack of vegetation, and this may be suboptimal (Barrows 2006). The quality of available food is probably dependent on the local rainfall, which varies widely from year to year throughout the species' range. Barrows (2006) found that a regression model including rainfall and diet explained 92% of the variation in *U. inornata* density and that population sizes could

approach zero during multiyear droughts and then quickly rebound when average rainfall resumed.

Habitat Requirements

Uma scoparia lives exclusively on fine wind-blown sand (Stebbins 1944). Habitat where lizards are found in the highest abundances generally consists of relatively sparse creosote scrub on loose sand dunes. The diameter of individual sand grains in these areas is usually <0.5 mm. Areas with large sand grains (>2 mm in diameter) appear to be avoided, presumably because this impedes sand swimming and burying behavior (Stebbins 1944, Norris 1958, Fromer et al. 1983). Within appropriate habitat, individuals select areas with the finest sand available (often the downwind side of vegetation and slopes) (Stebbins 1944, Norris 1958). Some vegetation is probably required for food and shade (Miller and Stebbins 1964). The species is not present in areas where the sand becomes too firmly packed to allow for sand swimming, and washes and desert flats are generally unsuitable (Miller and Stebbins 1964). No evidence exists that *Uma* will enter these areas to migrate between adjacent areas of suitable habitat, although additional study of this question would be valuable.

Uma scoparia may require relatively large habitat patches for long-term persistence. Population modeling in the ecologically similar *U. inornata* suggests that plot sizes smaller than 100–200 ha are unlikely to allow long-term persistence of isolated populations (Chen et al. 2006).

Distribution (Past and Present)

Uma scoparia is patchily distributed throughout much of the Mojave Desert in California. The range extends from near the southern end of Death Valley at the Inyo San Bernardino County line south through San Bernardino and Riverside Counties, extending west narrowly into Los Angeles County (Van Denburgh 1922, Norris 1958, Miller and Stebbins 1964, Pough

1974, Jennings and Hayes 1994a, Stebbins 2003). Norris (1958) reports a record from Inyo County, which has often been repeated in the literature. However, the stated locality “one and one-half miles southeast of Saratoga Springs” places this record in San Bernardino County, and we know of no other confirmed records from Inyo County. This species is nearly endemic to California, extending into Arizona in one small area near Parker, Yuma County (Pough 1974). A single report of possible *Uma* tracks reported from the Eureka Sand Dunes, Inyo County, California, would extend the known range ~175 km to the northwest and requires verification (Bolster et al. 2000). The known elevational range extends from below sea level to nearly 1000 m (Jennings and Hayes 1994a).

Extirpations have been documented at El Mirage and Harper Dry Lakes, San Bernardino County, and at Lovejoy Buttes and Piute Butte, Los Angeles County (Murphy et al. 2006). Additional extirpations may have occurred at Rogers Dry Lake, Kern County, California, and Saddleback Butte, Los Angeles County, California (CBD 2006).

Trends in Abundance

No quantitative data are available regarding historical abundance, though the lizard was, and is, common at many isolated localities. Some data suggest that this species has become uncommon in areas where habitat degradation due to off-highway vehicle use has occurred (Bolster et al. 2000, CBD 2006).

Nature and Degree of Threat

The most important threats facing *Uma scoparia* are habitat loss and fragmentation due to human activities and off-highway vehicle use, which negatively impacts loose sand habitat. Other activities, including the development of renewable energy facilities, may also negatively impact the structure of essential windblown sand habitat patches. The species is only found in loose sand areas, and experimental work in

the closely related and ecologically similar species *U. inornata* suggests that these lizards are highly sensitive to stabilization of their sand habitat (Turner et al. 1984). Habitat fragmentation is also an important threat. Even where patches of intact habitat remain, fragmentation and small patch sizes have been shown to be associated with declines and extirpations in *U. inornata* (Barrows and Allen 2007). In addition, surveys for the ecologically similar *U. notata* that compared lizard abundances in areas that experienced off-highway vehicle use to areas that do not, found much higher densities in the less-impacted habitat (Luckenbach and Bury 1983). Off-highway vehicles impact this species through direct mortality, destruction of vegetation (which is correlated with lizard abundance), and increased rates of tail loss (Luckenbach and Bury 1983, Ouren et al. 2007). Further, *U. scoparia* has sensitive hearing that is easily damaged by even moderate and short duration off-highway vehicle activity (Brattstrom and Bondello 1983). Hearing loss likely harms this lizard's efficiency at capturing prey and its ability to avoid predation (Brattstrom and Bondello 1983). Increasing predator densities (e.g., common ravens) in certain areas, often in association with human development and the presence of garbage dumps, may also be causing declines in lizard abundance in localized areas (Bolster et al. 2000).

Uma scoparia is likely sensitive to the effects of climate change. Climate change models for this region predict relatively sharp increases in mean temperature of up to 2°C (PRBO 2011). The impact of such increases on *U. scoparia* and on critical plant species is not known but could be large and should be a high priority for future research. The distribution of *U. inornata* is associated with an east-to-west drought gradient in the Coachella Valley (Barrows and Allen 2007). Like off-highway vehicle use, drought decreases the amount and quality of vegetation present, which limits both food and cover for this species (Barrows et al. 2010).

Because *Uma* specializes on relatively isolated patches of habitat, it is probably unable to track available habitat with changing climatic conditions. Climate change modeling studies on other *Uma* species (*U. inornata*; the Coahuila fringe-toed lizard, *U. exsul*; and the Chihuahuan fringe-toed lizard, *U. paraphygas*) predict significant habitat loss under a relatively wide range of climate change scenarios (Ballesteros-Barrera et al. 2007, Barrows et al. 2010) and these results are also likely to apply to *U. scoparia*.

Status Determination

Uma's specialized habitat is relatively uncommon and undergoing significant degradation, and this is the primary justification for Priority 3 designation. While some populations have been extirpated, several populations of this species are still common, and some habitat occurs on protected land that is not subject to off-highway vehicle use, precluding the need for a higher-priority designation.

On 10 April 2006, the Center for Biological Diversity and Sylvia Papadakos-Morafka petitioned the US Department of the Interior to list the northern population segment identified by Murphy et al. (2006) under the Federal Endangered Species Act (CBD 2006). The USFWS issued a 90-day finding that substantial evidence for listing need had been presented and initiated a 12-month status review for the taxon (USFWS 2008). This review concluded that the Amargosa River populations of *U. scoparia* do not constitute a distinct population segment and are therefore ineligible for listing under the US Endangered Species Act (USFWS 2011).

Management Recommendations

Effective management of this taxon over the short term can likely be accomplished by protecting habitat from development and degradation from off-highway vehicles and other human impacts. Over longer time periods, climate change could begin to have a larger impact, and this may require additional management efforts. Such efforts could range from

human-assisted translocation to planting drought-resistant vegetation, depending on local conditions and the extent of temperature and precipitation changes. If restoration occurs in areas where extirpation has occurred or if development activities further isolate occupied habitat patches, human-assisted translocation, potentially in association with captive breeding programs, may be a key strategy for this species.

Monitoring, Research and Survey Needs

Two key research efforts for *Uma scoparia* should focus on the effects of human activities (including off-highway vehicles, solar and wind energy development, and roads) and the genetic effects of both natural and anthropogenic habitat fragmentation. The effects of off-highway vehicles are particularly important, and monitoring efforts should be initiated in areas that experience off-highway vehicle use compared to more pristine, adjacent areas. In particular, these efforts should focus on comparing the effect of varying intensity of anthropogenic disturbance on populations, with the aim of establishing what intensity of off-highway vehicle use can be tolerated. These efforts should also attempt to disentangle the effects of habitat destruction, noise pollution, and direct mortality on populations, since each can in principle be managed independently. For example, if off-highway vehicle use primarily affects these lizards through reductions in vegetation, habitat restoration coupled with restricting off-highway vehicles to certain trails or corridors could constitute a reasonable management strategy. Alternatively, noise pollution effects may require eliminating off-highway vehicle access in areas where the lizards are present. Because population sizes naturally fluctuate with rainfall in this species (Barrows 2006), and in some cases can approach zero before rebounding, monitoring this species is inherently difficult, and multiyear surveys spanning several drought and non-drought years are essential. The frequency of lizard detection and the accuracy of population size estimates can be

increased with repeated sampling and specific detection methods (Turner et al. 1984, Bolster et al. 2000), and these should form the foundation of monitoring protocols.

Genetic studies are critical at two distinct levels of resolution that require different datasets and analytical approaches. Species boundaries across *Uma* and large-scale phylogeographic patterns within *U. scoparia* remain poorly resolved (see also the species account for *U. notata*), and both are critical for effective management. For species boundary work, the resolution of the number and identity of species contained within the genus requires a multi-locus nuclear dataset to complement initial work using mitochondrial DNA (Trépanier and Murphy 2001). Within *U. scoparia*, phylogeographic studies using multiple nuclear markers

are also needed in order to quantify the intraspecific diversity present within the species. At a finer scale, landscape and population genetic studies are also badly needed to establish natural levels of gene flow, including movement across seemingly inhospitable habitat patches, for this windblown sand habitat specialist. These data can advise and guide plans for habitat acquisition both now and in the face of climate change, and may be a critical element in establishing appropriate habitat corridors and supplementing ecological survey data to guide potential human-assisted translocation. Finally, these multi-locus microsatellite or SNP-based studies can help clarify the amount of migration (if any) between adjacent populations and effective population sizes of existing local populations.



SANDSTONE NIGHT LIZARD

Xantusia gracilis Grismer and Galvan 1986

Status Summary

Xantusia gracilis is a Priority 3 Species of Special Concern, receiving a Total Score/Total Possible of 38% (42/110). During the previous evaluation, it was also designated as a Species of Special Concern (as *Xantusia henshawi gracilis*; Jennings and Hayes 1994a).

Identification

Xantusia gracilis is a medium-sized (5.1–7 cm SVL) lizard with soft skin and granular scales on the dorsal surface, enlarged plates on the ventral surface, and a prominent gular fold (Grismer and Galvan 1986, Lovich and Grismer 2001, Stebbins 2003, Lovich 2009b). The dorsal coloration is pale tan/brown, with many round dark-brown spots, while the ventral surface is clean white or white, with a very small amount of black speckling on the front limbs and throat (Grismer and Galvan 1986). The head is flattened, and the eyes have vertically oriented pupils (Stebbins 2003). The overall body shape is relatively slender compared to its closest (and

most similar) relative the granite night lizard (*X. henshawi*) (Grismer and Galvan 1986).

Within its range, *X. gracilis* is only likely to be confused with its sister species *X. henshawi*.

Sandstone Night Lizard: Risk Factors

Ranking Criteria (Maximum Score)	Score
i. Range size (10)	10
ii. Distribution trend (25)	5
iii. Population concentration/migration (10)	0
iv. Endemism (10)	10
v. Ecological tolerance (10)	10
vi. Population trend (25)	0
vii. Vulnerability to climate change (10)	7
viii. Projected impacts (10)	0
Total Score	42
Total Possible	110
Total Score/Total Possible	0.38




PHOTO ON PREVIOUS PAGE: Sandstone night lizard, San Diego County, California. Courtesy of Jeff Lemm.

The two species do not overlap in range but occur within 32 km of each other. *Xantusia henschawi* has larger dark spots on the dorsal surface, more extensive speckling on the ventral surface, and an overall more robust body shape (Grismer and Galvan 1986). *Xantusia gracilis* also has an enlarged temporal scale (about half the size of the postparietal) compared to *X. henschawi* (typically less than one-quarter the size of the postparietal; Grismer and Galvan 1986). The peninsular leaf-toed gecko (*Phyllodactylus nocticolus*) also occurs in the vicinity of *X. gracilis*, but this lizard lacks the dark-brown dorsal spots and has prominent, expanded toe tips.

Taxonomic Relationships

Xantusia gracilis was initially described as a subspecies of *X. henschawi* on the basis of color, scalation, allozyme variation, and behavior (Grismer and Galvan 1986). The taxon was elevated to species status because it is diagnosable, geographically isolated, and forms a monophyletic clade nested within *X. henschawi* for a single mitochondrial locus (Lovich 2001). This arrangement is now widely accepted.

Life History

The life history of *Xantusia gracilis* is poorly studied, particularly so in wild populations. Given the species' overall similarity in most respects to *X. henschawi*, we expect that life history information from *X. henschawi* is a reasonably good predictor for *X. gracilis* (Lee 1975). However, the two taxa live in distinct habitats and show some behavioral differences in captivity, so some life history differences probably exist in the wild. *Xantusia gracilis* is likely active from spring through fall (Lemm 2006). In captivity, it has been shown to be more strongly nocturnal than *X. henschawi*, more frequently found on the sandy substrate on the bottom of the enclosure and does not seem to be limited to rock faces (Lee 1975, Grismer and Galvan 1986). Based on what is known about *X. henschawi*, we expect that *X. gracilis* has a low metabolic rate and is quite sedentary, feeding primarily ants, beetles, and spiders (Brattstrom 1952,

Lee 1975, Mautz 1979). In captivity, *X. gracilis* are also known to feed on the eggs of *Phyllodactylus nocticolus*, a behavior that captive *X. henschawi* in the same enclosure did not exhibit (Grismer and Galvan 1986). In *X. henschawi*, mating occurs in June and July, with one or two live young born in September or October (Brattstrom 1951, Lee 1975), and this may also be the case for *X. gracilis*. Individuals probably do not become reproductively mature until 2.5–3.5 years of age and are likely long-lived, although field data are lacking (Lee 1975).

Habitat Requirements

Xantusia gracilis lives in eroding sandstone and mudstone habitat where it utilizes crevices, rodent burrows, and the undersides of exfoliating rock flakes as shelter (Grismer and Galvan 1986). At night, it emerges from its shelters and can be found moving about on the surface (Grismer and Galvan 1986). This species is less dependent on exfoliating rock habitat than *X. henschawi* (Grismer and Galvan 1986).

Distribution (Past and Present)

Xantusia gracilis is restricted to one small area, approximately 3.9 km² in total area, on the southeastern flank of the Santa Rosa Mountains, entirely within Anza Borrego Desert State Park (Grismer and Galvan 1986). The known elevational range extends from approximately 240 to 305 m. Within this small region the species is patchily distributed, common in some areas and apparently absent in others (Grismer and Galvan 1986). *Xantusia henschawi* occurs approximately 32 km to the north and west, and no xantusiid lizards are known from the intervening area. No historical distribution data are available for this taxon, although we have no reason to think that the distribution has declined recently.

Trends in Abundance

No data on historical or current abundance have been published, although some have suggested that habitat quality has declined due to collection activity (R. Lovich, pers. comm.).

The extent and severity of such impact has not been quantified (R. Fisher, pers. comm.). Some amount of illegal collection occurs for this species, which may be driving small declines (M. Jorgensen, pers. comm.).

Nature and Degree of Threat

Xantusia gracilis lives in a fragile habitat in an extremely localized area. Damage to this small patch of habitat, be it from habitat destruction, invasive species, collecting, or climate change, is the largest risk facing the species (Lovich 2009b). It is also likely long-lived and late maturing with a low reproductive potential, and populations are likely to be slow to recover from declines. Some amount of illegal collecting occurs, which could be contributing to such declines, particularly in areas that are most easily accessible by road.

Status Determination

The extremely localized range and relative fragility of *Xantusia gracilis*' habitat are significant risk factors. The species' life history also predisposes it to decline in the face of any increased adult mortality. Although data are almost entirely lacking, *X. gracilis* appears to be relatively stable at the present time; thus, we designate it as a Priority 3 Species of Special Concern.

Management Recommendations

Limiting access and minimizing disturbance to *Xantusia gracilis*' habitat is currently the most important component of effective conservation. This management strategy should be reviewed as needed depending on the results of the surveys outlined below. All collecting should be restricted or eliminated unless it is absolutely

necessary for scientific purposes that further conservation of this species.

Monitoring, Research, and Survey Needs

As published historical or current abundances of *Xantusia gracilis* are lacking, publication of any existing data is a priority. Formal monitoring should be initiated to establish and publish baseline population data. These surveys should be performed at night, and it is essential not to disturb the fragile microhabitat (e.g., moving rocks or rock flakes, excavating rodent burrows). Aside from estimating population size, these surveys should also quantify and document any observed habitat disturbance. Year-to-year fluctuations in population size occur in other xantusiid lizards (Lee 1975) and are to be expected in *X. gracilis* as well. Establishing a long-term monitoring program is a critical objective. Additional surveys to establish the precise limits of the range of *X. gracilis* will help determine best practices for managing its fragile habitat in the heavily used Anza-Borrego Desert State Park.

The life history of this species has not been studied and an autecological study is badly needed to provide basic information on habitat suitability and reproduction. These data will be urgently needed should more extensive management efforts become necessary.

Finally, multi-locus microsatellite or SNP data should be collected to provide genetic estimates of effective population size, and potentially levels of gene flow, even for this restricted species. A key issue for this species is to sample individuals without invasive tissue-removal techniques, and it would probably be best to work out such protocols on *X. henshawi* before applying them to *X. gracilis*.



SIERRA NIGHT LIZARD

Xantusia vigilis sierrae Bezy 1967

Status Summary

Xantusia vigilis sierrae is a Priority 3 Species of Special Concern, receiving a Total Score/Total Possible of 47% (52/110). During the previous evaluation, it was also considered a Species of Special Concern (Jennings and Hayes 1994a).

Identification

Xantusia vigilis sierrae is a small (4–5.1 cm SVL), somewhat flattened lizard with granular dorsal scales, enlarged square ventral scales, soft skin, and a prominent gular fold (Bezy 1967, Stebbins 2003). The head is covered with enlarged plates, the eyes are lidless, and the pupils are vertical (Stebbins 2003). Most specimens are olive or grayish brown above, with a pattern of interconnected dark markings that form a network, which may give the animal a mottled appearance (Bezy 1967, Stebbins 2003). The ventral surface is light bluish pink and generally unmarked (Bezy 1967). A prominent light stripe extends from the rear of the

eye posteriorly to the neck or just beyond the neck (Bezy 1967).

Within its range, *X. v. sierrae* is unlikely to be confused with other species, although it is similar in appearance to the Yucca night lizard

Sierra Night Lizard: Risk Factors

Ranking Criteria (Maximum Score)	Score
i. Range size (10)	10
ii. Distribution trend (25)	0
iii. Population concentration/ migration (10)	0
iv. Endemism (10)	10
v. Ecological tolerance (10)	10
vi. Population trend (25)	5
vii. Vulnerability to climate change (10)	10
viii. Projected impacts (10)	7
Total Score	52
Total Possible	110
Total Score/Total Possible	0.47




PHOTO ON PREVIOUS PAGE: Sierra night lizard, Kern County, California. Courtesy of Jackson Shedd.

(*X. v. vigilis*), which occurs nearby. *Xantusia vigilis vigilis* has fairly distinct dark spots on the dorsal surface that do not form a network, though they form narrow longitudinal stripes in some populations (Stebbins 2003). Several aspects of the scalation also differentiate these two subspecies (Bezy 1967).

Taxonomic Relationships

Xantusia vigilis sierrae is a member of the *X. vigilis* species complex. It was initially recognized on the basis of habitat type, coloration, scalation, and femoral pore count (Bezy 1967). Since its initial recognition, genetic analyses have shown that *X. v. sierrae* forms a monophyletic group embedded within *X. vigilis* for mitochondrial and nuclear DNA sequence data (Sinclair et al. 2004, Leavitt et al. 2007). Allozyme data also suggest that it is distinct, but a close relative of *X. v. vigilis* (Bezy and Sites 1987). One population of *X. v. vigilis* that occurs within 60 km of *X. v. sierrae* is suspected to contain intergrades based on femoral pore counts (Bezy 1967), although geographically more proximate populations (~20 km apart) show no evidence of this intermediate condition (Leavitt et al. 2007). Sinclair et al. (2004) considered *X. v. sierrae* a "candidate species" whose status required further testing with additional data. Some recent taxonomic lists have elevated it to species status without additional justification (de Queiroz and Reeder 2008, Collins and Taggart 2009). The weight of current evidence suggests that species status is probably warranted, and a population genetic analysis of *X. v. sierrae* and nearby *X. v. vigilis* populations is needed help clarify this issue. In particular, such a study could quantify whether, and to what extent, migration and intergradation occur along the eastern edge of the taxon's range.

Life History

The life history of *Xantusia vigilis sierrae* has not been studied. However, among xantusiid species that have been examined, life history

features are largely conserved across southwestern United States, and we assume that the life history of *X. v. vigilis* may be a good predictor for *X. v. sierrae* in many respects. *Xantusia vigilis sierrae* is primarily a rock-dwelling species, whereas *X. v. vigilis* is more of a habitat generalist with some preference for fallen vegetation. Some aspects of the life history may therefore be more similar to other rock-specialist night lizards (e.g., *X. henshawi* or *X. gracilis*).

Based on information from other species, *X. v. sierrae* is probably a generalist predator that consumes a variety of small invertebrate prey (Brattstrom 1952, Stebbins 2003, Bezy 2009). Its diet is probably dominated by ants and other insects that occur within crevices (Brattstrom 1952, Bezy 2009). *Xantusia vigilis sierrae* is probably long-lived and takes 2.5–3.5 years to reach sexual maturity (Lee 1975), eventually producing 1 or 2 live young/year (Brattstrom 1951). This species likely has a low metabolic rate relative to other lizards and grows slowly (Mautz 1979). Daily activity cycles are unknown. Some rock-dwelling night lizards are largely diurnal and/or crepuscular (*X. henshawi*; Mautz and Case 1974), while others appear to be nocturnal (*X. gracilis*; Grismer and Galavan 1986).

Habitat Requirements

Xantusia vigilis sierrae is known primarily from exfoliating granite outcrops (Bezy 1967), though it can also be found under tree bark that has fallen on the ground or is loosely attached to trees (D. Leavitt, pers. comm.). Within its rocky habitat type, this species is more frequently found under large horizontal cap rocks than the more numerous, vertically oriented smaller flakes (Bezy 1967). *Xantusia vigilis sierrae* is also more frequently found in small clusters of one or a few boulders than in larger rock piles on rocky slopes and canyons (Bezy 1967). Some authors have speculated that this may reflect varying abundances associated with differences in predator access

(Jennings and Hayes 1994a), although it is also possible that it reflects differences in detectability. The dominant vegetation of its preferred habitat is foothill grassland with interspersed shrubs and woody vegetation (Bezy 1967).

Distribution (Past and Present)

Xantusia vigilis sierrae is known only from rocky hillsides on the western edge of the Greenhorn Mountains near Granite Station, Kern County, California (Bezy 1967, Stebbins 2003). The known elevational range extends from 450 to 500 m (Bezy 1967). No significant changes in distribution are known, although the development of small ranches may impact populations in the area.

Trends in Abundance

No historical or current abundance data are available for this taxon, although these lizards do not currently appear to be rare (D. Leavitt, pers. comm.). Moderate habitat degradation from previous collecting efforts as well as moderate amounts of landscape modification may be causing declines (R. Fisher, pers. comm.), although this has not been confirmed.

Nature and Degree of Threat

The primary threat facing *Xantusia vigilis sierrae* is its exceedingly small range that occurs on unprotected land. Development in the region is taking place and could have catastrophic effects on the existing populations, as could any fragmentation of the habitat that isolates granite outcrops in which this lizard lives. The rock cap and crevice habitat that this species prefers is also susceptible to degradation by humans (Jennings and Hayes 1994a, Stebbins 2003; D. Leavitt, pers. comm.).

Status Determination

Xantusia vigilis sierrae is a narrowly distributed habitat specialist that is endemic to a small region of the Sierra Nevada. However, no distributional declines have been documented, and

only small declines in abundance are suspected, resulting in a Priority 3 designation.

Management Recommendations

To protect this species, habitat loss and degradation need to be avoided. Effective protection of this species can likely be accomplished by protecting rocky habitats from most human interference, including intensive collecting efforts and protecting the surrounding area from development. Housing development in the form of ranchettes and other rural development projects should be closely managed to avoid impacting *Xantusia vigilis sierrae* populations, including provisions for habitat corridors to prevent fragmentation. It is unknown whether grazing adversely affects the species.

Monitoring, Research, and Survey Needs

Given the almost complete dearth of ecological work on this species, several research and monitoring needs are required for its future management and protection. Until recently, this species was known only to inhabit exfoliating granite, although its actual habitat utilization now seems to be somewhat broader than this. Further study of habitat use and preferences in *Xantusia vigilis sierrae* is essential to establish an effective management program.

A long-term population monitoring program needs to be initiated for this species, ideally across all utilized habitat types. These monitoring programs need not be extensive, but at minimum should document population size in disturbed and pristine habitats at regular intervals. Such monitoring can provide both critical data on natural population fluctuations and an early warning of declines in their initial stages.

Finally, genetic analyses using multiple nuclear markers are needed to address two important conservation issues. First, additional work at the phylogeographic/species boundary level is needed to determine whether *X. v. sierrae* is best considered a species or subspecies

within the *X. vigilis* complex. An important aspect of this work should be to examine populations in close proximity to *X. v. vigilis* to determine the degree and extent of admixture between these taxa. Second, landscape genetic work across its limited range is needed to quantify the degree of population isolation and substructure among habitat patches, migration

corridors that are most heavily used by the lizards, and effective population sizes of populations in ecologically diverse habitat patches. Ideally, tissue samples in the form of small tail clips should be collected each year from study populations to allow for genetic as well as demographic estimation of population size fluctuations over time.



CALIFORNIA GLOSSY SNAKE

Arizona elegans occidentalis Blanchard 1924

Status Summary

Arizona elegans occidentalis is a Priority 1 Species of Special Concern, receiving a Total Score/Total Possible of 67% (74/110). It was not on the list of candidates considered for Species of Special Concern designation during the previous evaluation (Jennings and Hayes 1994a).

Identification

Arizona elegans occidentalis is a medium-sized colubrid (64–99 cm SVL) with tan or brown dorsal coloration. It has dark-brown blotches edged in black running down the back and a series of similar, though smaller, blotches running down the sides (Klauber 1946, Stebbins 2003, Lemm 2006). The dorsal coloration is often lighter middorsally and darkens to a deeper brown on the sides. The lateral blotching sometimes touches the edges of the ventral belly scales, but otherwise the underside is unmarked (Klauber 1946). Scales are unkeeled, smooth and glossy, and only one pair of pre-

frontals are present (Stebbins 2003). A dark stripe runs from the corner of the mouth to the eye on each side of the face, and a third stripe connects the eyes across the posterior edge of the prefrontals (Blanchard 1924). An additional

California Glossy Snake: Risk Factors

Ranking Criteria (Maximum Score)	Score
i. Range size (10)	5
ii. Distribution trend (25)	25
iii. Population concentration/ migration (10)	0
iv. Endemism (10)	3
v. Ecological tolerance (10)	3
vi. Population trend (25)	25
vii. Vulnerability to climate change (10)	3
viii. Projected impacts (10)	10
Total Score	74
Total Possible	110
Total Score/Total Possible	0.67




PHOTO ON PREVIOUS PAGE: California glossy snake, San Diego County, California. Courtesy of Jeff Lemm.

dark spot is usually present below each eye (Klauber 1946).

In California, this taxon could be confused with other subspecies of *A. elegans*, with the gopher snake (*Pituophis catenifer*), or the night snakes (*Hypsiglena* spp.). This subspecies is generally darker than other subspecies of *A. elegans* in California, though intergrades are common along the desert slopes of the coastal mountains (Klauber 1946). Generally, *A. e. occidentalis* is best distinguished from other subspecies based on range. *Pituophis catenifer* has keeled scales and (usually) two pairs of prefrontals, while *Hypsiglena* is smaller (up to 66 cm), has strongly elliptical pupils, and an extensive dark blotch on the neck (Stebbins 2003).

Taxonomic Relationships

Arizona elegans occidentalis was initially described on the basis of scale counts and dorsal blotching and included all snakes in this genus ranging from California through southeastern Arizona (Blanchard 1924). Klauber (1946) later restricted this taxon and described two new subspecies occurring in eastern California (the Mojave glossy snake, *A. e. candida*, and the desert glossy snake, *A. e. eburnata*), which differ from *A. e. occidentalis* primarily in body color. Intraspecific (or intrageneric) variation has not yet been assessed genetically, although at the generic level, *Arizona* appears to be a relatively distant sister taxon to the long-nosed snake (*Rhinocheilus lecontei*) (Pyron and Burbrink 2009).

Life History

Arizona elegans is a nocturnal snake that is generally active from late February until November, depending on local weather conditions (Klauber 1946, Grismer 2002). In California, *A. e. occidentalis* reaches peak activity during May (Klauber 1946; S. Sweet, pers. comm.), with few specimens being collected throughout the remainder of the summer (Klauber 1939, Goldberg 2000). The species feeds primarily on diurnal lizards, which it captures while they

sleep, and small nocturnal mammals, which it ambushes (Klauber 1946, Rodríguez-Robles et al. 1999a). In a sample of 107 prey specimens, 50% were lizards (primarily *Sceloporus* and *Uta*) and 44% were mammals (primarily small rodents). Larger specimens are also known to take small birds and other snakes (Rodríguez-Robles et al. 1999a).

Arizona elegans retreats to burrows during the day, using either existing mammal burrows, excavations under rocks, or creating burrows for itself (Klauber 1946, Degenhardt et al. 1996). This species can be nocturnally active at relatively low temperatures (as low as 14°C, though typically 19–20°C; Cowles and Bogert 1944).

Reproduction is poorly studied in the wild, but museum specimens indicate that ovulation begins in June, and spermiogenesis occurs in late summer (Goldberg 2000). In *A. elegans* from New Mexico, ovulation also begins in June with oviposition occurring in July (Aldridge 1979). Clutch size is poorly documented in this subspecies, though two individuals contained three and seven eggs, respectively (Reynolds 1943, Klauber 1946). Across *A. elegans*, clutch size varies widely from 3 to 23 eggs, with a mean of 8.5 (Fitch 1970). Recent hatchlings are typically found in September (S. Sweet, pers. comm.).

Habitat Requirements

Arizona elegans is found in a wide variety of habitat types, including open desert, grasslands, shrublands, chaparral, and woodlands. However, only a subset of these habitat types occurs within *A. e. occidentalis*' range, primarily grasslands, fields, coastal sage scrub, and chaparral (Klauber 1946). No studies of habitat requirements exist, although this subspecies appears to prefer open microhabitats. The majority of records occur in relatively open patches in a surrounding matrix of denser vegetation (Klauber 1946). This subspecies can be patchy within its range, with certain areas consistently producing more records than others that have seemingly identical habitat (Klauber

1946). *Arizona elegans* appears to prefer areas where the soil is loose, which allows for burrowing (Grismer 2002, Stebbins 2003). Unpublished survey data indicate that *A. e. occidentalis* may prefer sandy soil habitats such as coastal sand dunes, alluvial creek beds, and ancient dunes on the marine terraces (R. Fisher, pers. comm.).

Distribution (Past and Present)

Range-wide, *Arizona elegans* occurs throughout much of southwestern North America, extending east as far as central Texas, Oklahoma, and Kansas, and south to central Mexico. Klauber (1946) restricted *A. e. occidentalis* to the central San Joaquin Valley south to the Tehachapi Mountains and along the base of the Coast Range mountains farther south to San Quintin, Baja California. This subspecies is known to occur from sea level to ~1800 m (Lemm 2006).

Arizona elegans occidentalis has apparently declined throughout much of its range. In San Diego County, survey data are available for Torrey Pines State Reserve, Point Loma, and the Tijuana Estuary. The subspecies was formerly present in these areas but now appears to be extirpated (Wells 1998, Case and Fisher 2001, Fisher 2004). Extensive agricultural development and habitat modification throughout the San Joaquin Valley and urban development within the Los Angeles basin have likely led to declines and/or extirpations in these areas as well (Stebbins 2003; R. Fisher, pers. comm.).

Trends in Abundance

Few abundance data exist for this subspecies. However, extensive early surveys of snakes in San Diego County failed to find the species, suggesting that they were uncommon (Klauber 1924). Bogert (1930) was aware of only two records for Los Angeles County. Klauber (1946) observed that *Arizona elegans occidentalis* existed in lower densities, relative to the total snake population, than either *A. e. candida* or *A. e. eburnata*, and that *A. e. occidentalis* was patchily distributed. Pitfall trapping data col-

lected by the US Geological Survey (USGS) over 17 years in San Diego, Orange, and Los Angeles Counties have resulted in only a single capture of this taxon (C. Rochester, pers. comm.). Presently, the subspecies is found less commonly than it once was throughout the San Diego region (Case and Fisher 2001, Lemm 2006). Both low densities and patchiness could make this taxon particularly susceptible to declines and may explain why the species has seemingly disappeared from some areas, while several other colubrid snakes remain present. Development continues within the species' range and thus ongoing declines in abundance are likely.

Nature and Degree of Threat

The greatest threat to this subspecies is habitat modification due to agricultural, commercial, and residential development. However, the specific mechanisms that cause declines are not well understood. Abundant prey and small habitat blocks that appear suitable remain in some developed areas, although the species may be sensitive to the light pollution arising from this development (Perry and Fisher 2006, Perry et al. 2008). This species' response to wildfire is not well understood, but increasing frequency and intensity of wildfires due to climate change may plausibly lead to habitat modification that impacts this taxon. The projected changes in wildfire regime in this area are mixed (PRBO 2011), so the degree of this threat is still unknown. Wildfires that are small in scale and intensity may have a beneficial impact by temporarily clearing patches of chaparral habitat, which then recover over a period of a few years, creating the patchwork of open and densely vegetated habitat that this species appear to prefer. Large and intense wildfires, conversely, kill chaparral and convert large habitat patches to grassland for longer periods of time. This process would likely have a detrimental impact on this species.

Status Determination

A moderately small range and moderate degree of ecological specialization and endemism,

coupled with documented declines within this species range and projected impacts from ongoing development, contribute to a Priority 1 designation for this subspecies.

Management Recommendations

Habitat protection is currently the most important management priority for *Arizona elegans occidentalis*. The studies outlined below will help to characterize habitat usage, home range size, distribution, and abundance. Once these data become available, a more specific management program can be developed that targets specific remaining populations and protects appropriately sized habitat blocks for the species' home range size and movement patterns.

Monitoring, Research, and Survey Needs

This is a poorly studied component of California's herpetofauna. Two immediate research priorities exist for this taxon. First, ecological studies need to be initiated to enhance our currently poor understanding of the life history and existing population sizes in this subspecies. Without this basic information, designing a coherent management strategy is impossible. These studies should take place in concert with survey efforts to more precisely quantify the

subspecies' present distribution. These surveys should employ a variety of techniques, likely including night driving, snake trapping, and artificial cover object transects in order to increase capture success. If reasonably high capture rates can be obtained, individually marking snakes for mark-recapture population size estimates should also be performed. Radiotelemetry studies may be a fruitful means for determining home range size and more thoroughly characterizing habitat usage, particularly given the indications that this species might have specific microhabitat preferences. Second, a species-wide phylogeographic study should be performed in order to elucidate intraspecific variation and identify appropriate units for conservation. Phylogenetic and phylogeographic studies of other wide-ranging snakes have frequently led to changes in the understanding of species boundaries and diversity, including the genetic diversity that exists within a species and its concordance with morphological subspecies boundaries. Finer-scale landscape ecological studies, particularly in concert with radiotelemetry on the same landscapes, would also provide important information for conservation strategies. These important data are entirely lacking for this taxon at present.



RED DIAMOND RATTLESNAKE

Crotalus ruber Cope 1892

Status Summary

Crotalus ruber is a Priority 3 Species of Special Concern, receiving a Total Score/Total Possible of 44% (48/110). During the previous evaluation, it was also considered a Species of Special Concern (Jennings and Hayes 1994a).

Identification

Crotalus ruber is a large (165 cm TL), heavy-bodied, tan, brick-red, reddish- or pinkish-brown rattlesnake (Stebbins 2003). As is typical of pit vipers, *C. ruber* has a large triangular head, a thin neck, and a heat-sensing pit on each side of the head between the eyes and nostrils. An average of 35 light-edged or indistinct diamonds run down the back (Ernst and Ernst 2003). The tail is ringed with alternating bands of black and white or gray, ending in a rattle. Two light stripes occur on the sides of the head, and the venter is light colored and unmarked (Ernst and Ernst 2003). The dorsal body scales are keeled.

Neonates of *C. ruber* are similar in appearance to the western diamond-backed rattle-

snake (*C. atrox*). Adults can be distinguished by coloration and behavior, with *C. ruber* much redder and less aggressive than *C. atrox*. In California, the ranges of these two species barely meet (Stebbins 2003).

Red Diamond Rattlesnake: Risk Factors

Ranking Criteria (Maximum Score)	Score
i. Range size (10)	10
ii. Distribution trend (25)	15
iii. Population concentration/ migration (10)	0
iv. Endemism (10)	0
v. Ecological tolerance (10)	3
vi. Population trend (25)	10
vii. Vulnerability to climate change (10)	3
viii. Projected impacts (10)	7
Total Score	48
Total Possible	110
Total Score/Total Possible	0.44




PHOTO ON PREVIOUS PAGE: Red diamond rattlesnake, San Diego County, California. Courtesy of Jeff Lemm.

Taxonomic Relationships

Based on analyses of morphology and mitochondrial DNA, Murphy et al. (1995) proposed synonymizing *Crotalus ruber* with an island species, the Cedros Island diamond rattlesnake (*C. exsul* Garman 1884). Because *C. exsul* was named first, Murphy et al. (1995) suggested changing the name of *C. ruber*. However, this has been opposed in favor of stability of the nomenclature (Smith et al. 1998, ICZN 2000). Here, we use *C. ruber* to refer to all California animals.

Life History

Crotalus ruber is generally most active between March and June (Ernst and Ernst 2003). In one study from San Diego County, snakes typically emerged from overwintering locations in late February, but some individuals were inactive until mid-April (Brown et al. 2008). Most movement occurred in late spring and summer, dens were populated in November, and no movement was recorded in December or January (Brown et al. 2008). In Riverside County, desert animals were active from early March to late November (Greenberg 2002).

During the cold winter months, *C. ruber* spends most of its time underground in dens located in rock crevices, animal burrows, or under shrubs or cacti. Several individuals may aggregate in these dens, but denning behavior is variable across sites (Klauber 1956, Ernst and Ernst 2003). In one study in San Diego County, 7 out of 11 radio-tracked snakes overwintered in communal dens located in rock crevices of granite boulders with up to 7 other individuals (Brown et al. 2008). Most snakes reused den sites over multiple years and moved ~300 m away from den sites during the active period the following year (Brown et al. 2008). In contrast, in sites where large rocks were rare, animals were observed to overwinter singly under prickly pears (*Opuntia* sp.), did not show consistent site fidelity to overwintering sites, and moved farther from overwintering sites after emergence (Greenberg 2002, Dugan et al. 2008).

Home range area is also variable in this species, and male home ranges are larger than those of females (Tracey 2000, Greenberg 2002, Brown et al. 2008). The few available radiotelemetry studies suggest that home ranges may be larger in the desert than in coastal habitats. In a reserve in San Diego County, average home range sizes were 2.8 ha for males ($n = 5$) and 0.9 ha for females ($n = 6$; Brown et al. 2008). At another relatively coastal site in Chino Hills State Park in southwestern San Bernardino County, Dugan et al. (2008) found that male home range size varied from 0.3 to 4.5 ha ($n = 7$). In contrast, average home range sizes for desert animals from Riverside County were 25.7 ha for males ($n = 5$) and 5.9 ha for females ($n = 4$; Greenberg and McClintock 2008).

Courtship and mating have been observed in the field in California from February to May (Brown et al. 2008, Dugan et al. 2008). In San Diego County, Brown et al. (2008) witnessed females mating from April to May (sometimes with den mates), and births occurred in September. Goldberg (1999) examined the reproductive condition of 43 specimens, 41 of which were from desert habitat in Riverside County and 2 from coastal Orange County. Reproductively active males were observed in August (Goldberg 1999). Although specimens were unavailable from later in the year, Goldberg (1999) speculated that sperm production continued through the early fall. Sperm was found in the vas deferens for all animals (collected February through August), suggesting the use of sperm stored overwinter for spring mating (Goldberg 1999). Females contained enlarged ovarian follicles (>10 mm) from March through September. Females may reproduce every other year, given that only 7 of 15 females showed evidence of reproductive activity (Goldberg 1999).

An average of eight young (range 3–20, $n = 40$; Klauber 1956) are live-born after a gestation period of 141–173 days ($n = 3$, data from captive animals; Klauber 1956). Goldberg (1999) estimated similar average litter sizes from counts of enlarged ovarian follicles (range 4–8, mean

6.3, $n = 7$). Klauber (1956) examined 249 specimens from San Diego County to estimate growth curves and found that young are 30 cm TL at birth on average and roughly double in length during their first year. Estimates for size at reproductive maturity range from 60 to 75 cm TL (Klauber 1956, Wright and Wright 1957, Goldberg 1999).

Crotalus ruber mostly feeds on small mammals but will also eat lizards, birds, and other snakes (Tevis 1943, Klauber 1956, Cunningham 1959b, Patten and Banta 1980). Dugan and Hayes (2012) compiled range-wide dietary data from museum specimens, live animals, road kills, existing literature, and other observations. Roughly 92% of all prey items were mammals, with lizards (8%) and birds (1%) taken less frequently. Prey items were found in snakes collected year-round, suggesting that *C. ruber* occasionally feeds during the winter (Dugan and Hayes 2012).

Habitat Requirements

Crotalus ruber occurs in several habitat types, including coastal sage scrub, chamise chaparral, redshank, desert slope scrub, desert washes, grassy fields, orchards, cactus patches, and rocky areas (Klauber 1956, Jennings and Hayes 1994a, Tracey 2000, Dugan et al. 2008). Klauber (1956) noted that 44% (30/68) of animals were found near heavy shrub and chaparral, and 21% (14/68) were found near rocks and boulders in road surveys. On a reserve in San Diego County, snakes were found in association with rock outcrops 57% of the time and in shrubby vegetated habitats without rocks 28% of the time (Brown et al. 2008). There are several accounts of *C. ruber* climbing in bushes and trees up to 2 m off the ground (Klauber 1956 and pers. comm. therein) and *C. ruber* has also been observed swimming in reservoirs (Klauber 1956).

In one radio-tracking study from San Diego County, habitat use was nonrandom with respect to available vegetation. Snakes preferred scrub vegetation less than 1.5 m tall and avoided human development (Tracey 2000). For ani-

mals that were radio-tracked in fragmented habitats, none were observed to cross a developed edge or road over a 2-year period. For example, one adult male in a naturally vegetated fragment actively avoided a road edge, and turning movements away from this edge were detectable up to 50 m from the road (Tracey et al. 2005).

Dugan et al. (2008) radio-tracked adult males at a site that lacked large rocks but had cactus, coastal sage scrub, nonnative grassland, riparian areas, and oak woodland habitats. The preferred habitat was cactus patches of prickly pear (*Opuntia* sp.) followed by chaparral, and none of the tracked snakes used oak woodland. Several individuals spent most of their time within a single cactus patch during the year (Dugan et al. 2008).

Distribution (Past and Present)

Crotalus ruber has a small range in California, occupying the southwestern corner of the state. It occurs in southeastern Los Angeles and Orange Counties, the Morongo area of southwestern San Bernardino County, western Riverside County, San Diego County, and extreme southwestern Imperial County (Klauber 1956). *Crotalus ruber* occurs in areas with rainfall ranging from 8 to 80 cm/year, usually in areas below 1200 m in elevation (Klauber 1956). The geographic range of *C. ruber* extends out onto the desert floor from the eastern slope of the Peninsular Ranges (Klauber 1956). Outside of California its range extends south through Baja California and several nearshore islands (Klauber 1956).

Much of the range in California is in close proximity to areas of high human density. Jennings and Hayes (1994a) estimated that *C. ruber* was extirpated from roughly 20% of historical sites and attributed extirpations to habitat loss from urbanization and agriculture. Coastal populations are the most reduced, particularly in southern San Diego County (S. Barry, pers. comm.). Case and Fisher (2001) conducted pitfall trapping surveys in southern California and did not capture or observe

animals at several localities where Klauber (1939 and unpublished data) had previously noted them as common. Halama et al. (2008) noted that many native habitat localities where snakes were collected in the 1990s in western Riverside County have now been developed.

Trends in Abundance

While population estimates are not available, population declines are suspected due to habitat loss and fragmentation. Current declines of existing populations may be occurring particularly in the Morongo Valley in the northern end of the range due to development (S. Barry, pers. comm.). In one San Diego County site, minimum density was estimated as 0.63 *Crotalus ruber* per hectare, although the actual density was likely higher (41 individuals observed haphazardly in a 65 ha area over ~5 years; Brown et al. 2008).

Nature and Degree of Threat

Crotalus ruber is mainly threatened by development, which causes habitat loss and fragmentation. This species may also suffer from persecution and road mortality. Climate change may affect *C. ruber* through changes in fire regime and vegetation shifts. However, both increases and decreases in fire have been predicted, and there is little consensus because of the difficulty in modeling Santa Ana weather events in southern California (Westerling et al. 2004, Westerling and Bryant 2008). How *C. ruber* may respond to changes in fire regime is unknown. Climate change is predicted to decrease the availability of chaparral and shrubland by up to 44%, while grassland is predicted to increase by up to 390% in southern California (Lenihan et al. 2008, PRBO 2011). Though *C. ruber* has been documented in grassy areas, large losses in shrub habitat may negatively affect this species.

Status Determination

Crotalus ruber has a small range in California that includes areas of high human population

density and development, resulting in a Priority 3 Species of Special Concern designation.

Management Recommendations

Remaining populations of *Crotalus ruber* in California often occur in habitats that are fragmented by roads and urban development. Existing large habitat fragments should be identified and protected. For example, a proposed Habitat Conservation Plan for western Riverside County represents a 3.5-fold increase in the amount of snake habitat protected (Halama et al. 2008). However, Halama et al. (2008) estimated from habitat suitability models that roughly 100,000 ha of predicted highly suitable habitat in the area would still be unprotected and at risk of development. It may be possible to reduce road mortality with wildlife tunnels and associated drift fences installed beneath high-traffic roads.

Monitoring, Research, and Survey Needs

Additional research into *Crotalus ruber* ecology and population dynamics in developed and fragmented landscapes would be useful for developing management strategies, particularly with regard to maintaining connectivity among populations. Creating habitat buffers around large remaining fragments and habitat corridors between fragments may help populations persist in these landscapes, but more research on habitat use and corridor placement is needed. Radiotelemetry data to date suggest high site fidelity among adults, at least in some coastal populations within years. Juveniles may show different dispersal behavior and benefit more from management strategies like habitat corridors (Tracey 2000). Current snake telemetry techniques that rely on surgically implanted transmitters have a lower size limit (e.g., animals needed to be >500 g in one study; Brown et al. 2008), making it difficult to study movement in small individuals. In these cases, landscape genetic data could provide important data to complement more detailed telemetry studies. The role of hibernacula in population viability

and movement patterns is also an important research need, particularly for juveniles.

Monitoring is needed to estimate abundances in addition to ongoing work on presence/absence to document local extirpations. Pitfall trapping has been used to success-

fully document presence and absence of this species in southern California (e.g., Case and Fisher 2001), and pitfall arrays that specifically compare habitats with different levels of human disturbance would provide valuable monitoring information.



REGAL RING-NECKED SNAKE

Diadophis punctatus regalis Baird and Girard 1853a

Status Summary

Diadophis punctatus regalis is a Species of Special Concern, although we refrain from assigning it a priority status at this time due to limited information. This taxon received a Total Score/ Total Possible of 68% (27/40) and was not previously considered a Species of Special Concern (Jennings and Hayes 1994a).

Identification

Diadophis punctatus regalis is the largest of the ring-necked snakes, reaching up to 85.7 cm TL, while most subspecies are less than 50 cm TL (Ernst and Ernst 2003). A slender snake with smooth scales, *D. p. regalis* is light gray, olive gray, or olive above with orange or red ventral coloration. The venter is speckled with irregular black spots. An orange or red neckband is generally present behind the head, though it can be faint or absent in some populations of this subspecies, particularly in New Mexico and Utah (Ernst and Ernst 2003, Stebbins 2003). Recent specimens from California and

Nevada have lacked neck rings (Emmerich and Cunningham 2003, Wood and Richmond 2003).

Regal Ring-Necked Snake: Risk Factors

Ranking Criteria (Maximum Score)	Score
i. Range size (10)	10
ii. Distribution trend (25)	Data deficient
iii. Population concentration/ migration (10)	Data deficient
iv. Endemism (10)	0
v. Ecological tolerance (10)	10
vi. Population trend (25)	Data deficient
vii. Vulnerability to climate change (10)	7
viii. Projected impacts (10)	Data deficient
Total Score	27
Total Possible	40
Total Score/Total Possible	0.68




PHOTO ON PREVIOUS PAGE: Regal ring-necked snake, Santa Cruz County, Arizona. Courtesy of Jackson Shedd.

Taxonomic Relationships

Twelve subspecies of *D. punctatus* have traditionally been recognized, largely on the basis of morphology (Ernst and Ernst 2003). *Diadophis punctatus regalis* is one of seven subspecies that occur in California (Ernst and Ernst 2003). Recent molecular work has called this traditional view of the subspecies into question (Feldman and Spicer 2006, Fontanella et al. 2008), and a taxonomic revision is likely in the near future. Feldman and Spicer (2006) sampled mitochondrial DNA from 39 animals throughout the range of *D. punctatus* in California. *Diadophis punctatus regalis* was recovered as sister to a clade containing all other California samples, although only two *D. p. regalis* individuals were included in the analysis (one from California and one from Arizona). Fontanella et al. (2008) conducted a more comprehensive phylogeographic analysis of *D. punctatus*, sampling across the known range of the species in the United States. The previously recognized seven subspecies in California were found to fall into four lineages, with *D. p. regalis* as a part of a Great Basin clade. Fontanella et al. (2008) concluded that species-level diversity is currently underestimated, warranting a full taxonomic review requiring further sampling (particularly throughout Mexico) and the addition of nuclear markers.

Life History

Very little natural history information is available for *Diadophis punctatus regalis*, especially for California populations. Being such a widespread species, life history characteristics vary greatly across the species' range. It is reasonable to presume that *D. p. regalis* are ecologically distinct from other California *D. punctatus* populations based on their much larger size and unique restriction to desert spring habitats. Unless stated explicitly, life history information here is from other subspecies of *D. punctatus* and caution should be used in generalizing to *D. p. regalis*.

Diadophis punctatus is most active in the spring and early fall, and is primarily nocturnal (Ernst and Ernst 2003). Males aggregate for

mating in the spring and fall (Noble and Clausen 1936, Dundee and Miller 1968). Females are thought to reproduce annually and may produce more than one clutch per year (Ernst and Ernst 2003). Oviposition occurs from May to September but is concentrated in June and July, and hatching occurs from July to September (Ernst and Ernst 2003). Clutches from multiple females may be laid together in communal nest sites (Blanchard 1942, Gilhen 1970). *Diadophis punctatus* eggs are 16–44 mm long (mean 25 mm, $n = 108$) and hatchlings are 7.6–18.8 cm TL (mean 12.4 cm, $n = 120$; Ernst and Ernst 2003). *Diadophis punctatus regalis* eggs and hatchlings are likely at the larger end of the spectrum. A field-collected 60 cm SVL female *D. p. regalis* from Arizona contained three large eggs (mean length 44 mm, mean width 11.3 mm) that hatched after 52 days of incubation (Vitt 1975). The neonates were 16.9–18.8 cm long (mean 18 cm). Gehlbach (1965) reported one female *D. p. regalis* carrying 18 eggs. Estimates for size at maturity for *D. punctatus* range from 17.8 to 18 cm (Wright and Wright 1957, Myers 1965), but given that hatchling *D. p. regalis* can be this large, they likely mature at a larger size. Development times to maturity in *D. punctatus* can take 1–3 years depending on locality (Fitch 1975, Degenhardt et al. 1996).

Diadophis punctatus can often be found in aggregations under cover objects (Ernst and Ernst 2003), and some populations make spring and fall migrations to and from hibernacula. *Diadophis punctatus regalis* from the Rocky Mountains in Utah at 1580 m elevation showed communal denning and repeated use of the same hibernacula in multiple years (Parker and Brown 1974). It is unknown whether California populations of *D. p. regalis* also show this behavior. Field-active body temperatures across several populations of *D. punctatus* range from 2.0°C to 34.4°C (Clarke 1958, Brattstrom 1965, Fitch 1975, Mitchell 1994).

Diadophis punctatus regalis is a mildly venomous rear-fanged colubrid snake, using enlarged posterior teeth to deliver venom to

prey such as snakes and lizards (Gehlbach 1974, Anton 1994, Hill and Mackessy 2000, O'Donnell et al. 2007). In addition to subduing prey, copious salivation has been observed as a defensive response in *D. p. regalis* (Blanchard 1942). While *D. punctatus* is a generalized predator, southwestern populations, including *D. p. regalis*, have a diet composed of proportionately more reptiles (Gehlbach 1974) than other populations, which tend to consume a greater fraction of amphibians and earthworms (Ernst and Ernst 2003).

Habitat Requirements

In California, *Diadophis punctatus regalis* appears to be restricted to riparian areas surrounding desert springs. Snakes have been found in Death Valley in Inyo County in heavy riparian vegetation within 5 m of surface water (Emmerich and Cunningham 2003) and at Pachalka Spring, Clark Mountain, San Bernardino County, near the spring head (Wood and Richmond 2003). Outside of California, *D. p. regalis* have been found in evergreen woodland, deciduous woodland, desert grassland, oak-juniper, and succulent desert habitats such as sotol-agave and juniper-agave (Gehlbach 1974).

Distribution (Past and Present)

In California, the documented range of *Diadophis punctatus regalis* is extremely small. It is known from only a few isolated populations in the Clark, Grapevine, Mute, and Providence Mountains in the Mojave Desert. However, there is a strong possibility that undetected populations exist, particularly at additional springs in the mountain ranges where this taxon occurs. Outside of California, the subspecies occurs in parts of Idaho, Utah, Nevada, Arizona, New Mexico, Texas, and Mexico (Ernst and Ernst 2003). The Great Basin lineage defined by Fontanella et al. (2008) ranges from southern New Mexico north to southern Idaho and between roughly the Sierra Nevada Mountains in the west and the Guadalupe Mountains in the east. Much of the Great Basin clade is

restricted to patches of suitable mesic environments surrounded by less hospitable xeric habitats (Fontanella et al. 2008).

Trends in Abundance

No population estimates are available for California populations. Declines may have occurred near Fort Piute in the Mute Mountain Range (R. Fisher, pers. comm.).

Nature and Degree of Threat

The small and patchy distribution of *Diadophis punctatus regalis* in California makes it at risk of extirpation. Its dependence on rare desert spring habitats is an additional risk factor, particularly because these spring habitats are threatened by overexploitation of groundwater resources. Because of their reliance on mesic habitats in an arid matrix, *D. p. regalis* may also be sensitive to climate changes that affect the timing and amount of precipitation. While there is a large degree of uncertainty in how rainfall patterns will change within its range, most studies predict decreases in mean annual rainfall of up to 40% (reviewed in PRBO 2011). In addition to decreases in mean annual rainfall, the number of extremely hot days where temperatures exceed the long-term 95th percentile is expected to increase by roughly 30 days a year (Bell et al. 2004). If conditions become warmer and drier, this could negatively impact *D. p. regalis* habitat.

Status Determination

Diadophis punctatus regalis has an extremely small range in California and is dependent upon a rare habitat type that is sensitive to human use of groundwater in the desert. However, extirpations have not been well documented, and we have virtually no information about the number or status of populations occurring in California. Because of this, we choose not to define a priority at this time.

Management Recommendations

Protecting desert springs and associated mesic habitat patches is a key requirement for the

continued existence of this species. Minimizing use of water from desert spring sites will help maintain habitat for *Diadophis punctatus regalis*. It is difficult to make further management recommendations given the lack of information on this taxon's ecology in California or similar habitats.

Monitoring, Research, and Survey Needs

Given the inhospitable nature of habitat between occupied patches, long-distance dispersal events probably do not occur, and it is unlikely that populations in California are demographically connected. The extremely iso-

lated nature of most of their desert habitat and their relatively short surface activity period also make increased surveys an important priority for this taxon. Increased genetic sampling would help determine patterns of connectivity between fragmented southwestern populations, and we strongly encourage all field surveys to take nonlethal tissue samples of any specimens that are encountered. Given that no population study has been conducted within California and the unique habitat requirements of the taxon, additional work quantifying the basic ecology and life history of *Diadophis punctatus regalis* is badly needed.



SAN JOAQUIN COACHWHIP

Masticophis flagellum ruddocki Brattstrom and Warren 1953

Status Summary

Masticophis flagellum ruddocki is a Priority 2 Species of Special Concern, receiving a Total Score/Total Possible of 53% (58/110). It was previously considered a Species of Special Concern (Jennings and Hayes 1994a).

Identification

Masticophis flagellum is a large (91–260 cm TL) slender colubrid snake with smooth scales and a large head and eyes (Stebbins 2003). The species is distributed across the southern portion of the United States from Florida to California, with western subspecies tending to be smaller than eastern animals. Brattstrom and Warren (1953) reported that their largest specimen of *M. f. ruddocki* was 170 cm TL. Coloration is highly variable within *M. flagellum*. The subspecies *M. f. ruddocki* has a tan, olive-brown, or yellowish-brown dorsal color and lacks the dark head and neckbands characteristic of other subspecies. The ventral coloration is light tan or yellow, with a pink or orange cast under the tail

(Jennings and Hayes 1994a, Stebbins 2003). The scales on the tail are often described as having a “braided” appearance (Stebbins 2003).

San Joaquin Coachwhip: Risk Factors

Ranking Metric (Maximum Score)	Score
i. Range size (10)	5
ii. Distribution trend (25)	20
iii. Population concentration/ migration (10)	0
iv. Endemism (10)	10
v. Ecological tolerance (10)	3
vi. Population trend (25)	10
vii. Vulnerability to climate change (10)	3
viii. Projected impacts (10)	7
Total Score	58
Total Possible	110
Total Score/Total Possible	0.53




PHOTO ON PREVIOUS PAGE: San Joaquin coachwhip, Kern County, California. Courtesy of Jeff Lemm.

Taxonomic Relationships

Brattstrom and Warren (1953) described *Masticophis flagellum ruddocki* as a subspecies based on morphological characters including dorsal coloration, the lack of dark neckbands, and a relatively low number of subcaudal scales. Analysis of mitochondrial DNA from California populations supported the uniqueness of *M. f. ruddocki*, corroborating the morphological data (Mitrovich 2006).

Life History

Very little is known about the life history of *Masticophis flagellum ruddocki*. In general, *M. flagellum* is an extremely active diurnal snake that prefers warm temperatures (Brattstrom 1965, Hammerson 1977). Home ranges are suspected to be large, but no movement data are available for this subspecies (R. Hansen, pers. comm. in Jennings and Hayes 1994a). Preference for warm temperatures results in late-season emergence (April–May), and daily surface activity corresponds to the warmest parts of the day (Hammerson 1977). Data from red coachwhips (*M. f. piceus*) in the Mojave Desert found body temperatures as low as 13.9°C when inactive in burrows to a high of 40.8°C while actively moving (Secor 1995). Mating is thought to take place in May, with oviposition occurring in June or July (Jennings and Hayes 1994a). Adults may cease surface activity and retreat to mammal burrows as early as August (pers. obs. in Jennings and Hayes 1994a).

Like other members of the *M. flagellum* complex, the diet of *M. f. ruddocki* is presumably generalized on vertebrates, including large prey like antelope squirrels (*Ammospermophilus nelsoni*), blunt-nosed leopard lizards (*Gambelia sila*), and whiptails (*Aspidoscelis tigris*) (Montanucci 1965, Tollestrup 1979; S. Barry, pers. comm.; R. Hansen, pers. comm.; S. Sweet, pers. comm. in Jennings and Hayes 1994a).

Habitat Requirements

Masticophis flagellum ruddocki occurs in open, dry areas with little or no tree cover (Morafka and Banta 1976). Valley grassland and saltbush

scrub habitats are used in the western San Joaquin Valley (Montanucci 1965, Banta and Morafka 1968, Tollestrup 1979, Sullivan 1981; pers. obs. in Jennings and Hayes 1994a). Spring road cruising surveys from 1972 to 1979 in eastern Alameda and western San Joaquin Counties found *M. f. ruddocki* in grassland and transitional habitat but not in mixed oak chaparral woodland (Sullivan 1981). *Masticophis flagellum ruddocki* will climb into bushes, apparently to scan for predators and prey or to seek cover (Cunningham 1955, Stebbins 2003). Mammal burrows are used for overwintering and possibly also for oviposition (Jennings and Hayes 1994a).

Distribution (Past and Present)

Masticophis flagellum ruddocki is endemic to California, with a small range extending from Arbuckle, Colusa County, in the Sacramento Valley south to the Kern County portion of the San Joaquin Valley and west into the inner South Coast Ranges (Brattstrom and Warren 1953, Jennings and Hayes 1994a). A disjunct population occurs in the Sutter Buttes (Hayes and Cliff 1982).

Much of this subspecies' historic range has undergone dramatic land use changes from grassland to intensive agriculture in the Central Valley. *Masticophis flagellum ruddocki* is thought to be sensitive to disturbance and does not persist in cultivated areas (Ernst and Ernst 2003; S. Barry, pers. comm.). It has therefore suffered a severe range contraction in its Central Valley range.

Trends in Abundance

Though neither historical nor current abundance estimates are available, we suspect that the conversion of historical habitat to row crop agriculture and urban development has resulted in lower abundances than in preagricultural times.

Nature and Degree of Threat

Habitat loss and fragmentation due to agriculture and urbanization are the major threats to

Masticophis flagellum ruddocki. As with other diurnally active, highly mobile snakes, road mortality is probably a significant source of mortality, although its overall impact requires more study. The greatest potential threats from climate change are due to changes in fire regime. In the more coastal parts of the range, the area burned is expected to increase by up to 50% (Fried et al. 2004, Lenihan et al. 2008), and the probability of large (>200 ha) fires is predicted to increase (Westerling and Bryant 2008). Modest decreases in the probability of large wildfires are expected in the San Joaquin Valley. How *M. f. ruddocki* may respond to increased fire needs more study. Fire may have direct mortality effects on snakes and negative effects on prey populations but may also benefit *M. f. ruddocki* by increasing or maintaining the availability of open habitat. Under climate change projections, grassland habitat is expected to increase by up to 140% in the coastal part of the range, with little change in vegetation expected in the Central Valley (Lenihan et al. 2008, PRBO 2011). These vegetation shifts may result in additional potential habitat for *M. f. ruddocki*.

Status Determination

Masticophis flagellum ruddocki is a California endemic with a small range and is restricted to a heavily disturbed part of the state, resulting in a Priority 2 designation.

Management Recommendations

The lack of basic ecological information on this subspecies needs to be addressed before any meaningful management can be accomplished. At a minimum, remaining large habitat fragments and connectivity among fragments must be protected if the species is to persist.

Monitoring, Research, and Survey Needs

Although additional work on all aspects of its ecology, demography, and population genetic differentiation would be useful, information on reproductive biology, movement ecology, population sizes, and fragmentation is key priority for future work. Some large habitat fragments are currently protected from some kinds of human disturbance (e.g., the Carrizo Plain National Monument) and provide suitable areas to begin studying basic ecology and habitat requirements in this taxon.



BAJA CALIFORNIA COACHWHIP
Masticophis fuliginosus (Cope 1895a)

Status Summary

Masticophis fuliginosus is a Priority 3 Species of Special Concern, receiving a Total Score/ Total Possible of 45% (50/110). This species has not previously been considered a Species of Special Concern (Jennings and Hayes 1994a).

Identification

Masticophis fuliginosus is a large slender colubrid snake with smooth scales, reaching up to 170 cm in TL (Grismer 2002). Two color phases exist in the species, a light morph and a dark morph. The light morph is yellowish, tan, or gray above with dark zigzagging bands on the body and dark neckbands. The dark morph has a dark gray brown, golden brown, or black dorsal ground color, and sometimes has distinguishable dark neckbands (Wilson 1971, Grismer 2002). Dark morph animals can be uniformly dark above, or the scales on the sides of the body can have pale edges, giving the appearance of narrow lines (Wilson 1971). The

venter is light colored with brown spots (Wilson 1971, Grismer 2002).

Masticophis fuliginosus can be distinguished from congeneric southern California snakes by geographic range, as there is little overlap with

Baja California Coachwhip: Risk Factors

Ranking Criteria (Maximum Score)	Score
i. Range size (10)	10
ii. Distribution trend (25)	20
iii. Population concentration/ migration (10)	0
iv. Endemism (10)	0
v. Ecological tolerance (10)	0
vi. Population trend (25)	10
vii. Vulnerability to climate change (10)	3
viii. Projected impacts (10)	7
Total Score	50
Total Possible	110
Total Score/Total Possible	0.45




PHOTO ON PREVIOUS PAGE: Baja California coachwhip, Baja California, Mexico. Courtesy of Brad Shaffer.

other species, and by color. *Masticophis flagellum piceus* is reddish to pinkish above, with dark bands at the neck. California whipsnakes (*M. lateralis*) have a conspicuous light stripe on either side. Racers (*Coluber constrictor*) are pale green or dark above (brown, olive, or bluish) but have unmarked white to yellow ventral surfaces (Stebbins 2003).

Taxonomic Relationships

We follow Grismer's (2002) proposal that *Masticophis fuliginosus* is a full species, rather than a subspecies of *M. flagellum*. This arrangement is based on a lack of intergradation with neighboring *M. f. piceus* (Wilson 1971, Grismer 1994). Analysis of a single mitochondrial DNA gene from 229 *M. flagellum* individuals (including 30 *M. fuliginosus*) from 30 localities in southern California supported the genetic distinctiveness of *M. fuliginosus* (Mitrovich 2006). However, 4 out of 30 snakes identified in the field as *M. fuliginosus* had mitochondrial DNA sequences that were most closely related to *M. f. piceus* (Mitrovich 2006). This could be due to hybridization or incorrect identification in the field, as the study was conducted on tissues without voucher specimens. Further resolution of this problem with multiple nuclear DNA markers would likely help to clarify the taxonomic status of this species.

Life History

Very little is known about the life history of *Masticophis fuliginosus* in California. In general, *Masticophis* are extremely active diurnal snakes that prefer warm temperatures (Brattstrom 1965, Hammerson 1977). In southern Baja California, *M. fuliginosus* can be active year-round, but in the northern part of the range, they tend to be inactive in winter and emerge in mid-March (Grismer 2002). Activity in San Diego was observed to be greatest in spring and summer and greatly reduced in the fall (Mitrovich et al. 2009). Mating has been observed in northern Baja California in late April, and hatchlings have been seen in early August (Grismer 2002).

Radiotelemetry of 24 snakes in two reserves in San Diego County found large variation in home range size, from roughly 11 to 130 ha (Mitrovich et al. 2009). Variation in home range size was largely due to habitat availability, with smaller home ranges in smaller habitat fragments. No differences in potential prey were detected among sites where snakes had different home range sizes. The diet of *M. fuliginosus*, like its close relative *M. flagellum*, is broad and includes a variety of vertebrate prey such as lizards, snakes, birds, and mammals (Cliff 1954, Grismer 2002).

Habitat Requirements

Masticophis fuliginosus is a habitat generalist throughout Baja California, Mexico, and is common in marshlands, coastal sand dunes, rocky arroyos and hillsides, thorn forests, sandy flats, and scrub vegetation (Linsdale 1932, Cliff 1954, Leviton and Banta 1964, Bostic 1971, Welsh 1988, Grismer 2002). In California, *M. fuliginosus* occurs mainly in coastal sand dunes, shrubland, and grassland, and is most commonly observed foraging in bushes and shrubs (Linsdale 1932, Bostic 1971, Welsh 1988, Grismer 2002). Hollow stumps of plants such as agave and yucca are used as retreats (Bostic 1971, Grismer 2002).

Distribution (Past and Present)

Masticophis fuliginosus has a very small geographic range in California, occurring in a small area of San Diego County near the United States–Mexico border (Wilson 1973). The range of the species extends over most of the Baja California peninsula, including some small offshore islands (Wilson 1973, Grismer 2002).

A resurvey of Klauber's (1939) sites in southern California found that *M. fuliginosus* was absent from some previously occupied sites (Fisher and Case 2000, Case and Fisher 2001; R. Fisher, pers. comm.), suggesting that the species has declined in the last seven decades. However, the full extent of local extirpations is unknown.

Trends in Abundance

While data on abundance across the range are not available, some reductions in abundance are likely to have occurred due to development, road mortality, and fragmentation. This species may be particularly prone to death from automobiles given its large home range size and high level of diurnal activity (Mitrovich et al. 2009).

Nature and Degree of Threat

Masticophis fuliginosus in California are mainly threatened by habitat loss, fragmentation, and road mortality due to development, as well as the inherent demographic threats associated with a very small geographic range. Climate change may affect *M. fuliginosus* through changes in fire regime and vegetation shifts. However, both increases and decreases in fire have been predicted, and there is little consensus because of the difficulty in modeling Santa Ana weather events in southern California (Westerling et al. 2004, Westerling and Bryant 2008). How *M. fuliginosus* may respond to changes in fire regime is unknown. Climate change is predicted to decrease the availability of chaparral and shrubland by up to 44%, while grassland is predicted to increase by up to 390% in southern California (Lenihan et al. 2008, PRBO 2011). Though *M. fuliginosus* also uses grassland habitat, large losses in shrub habitat may negatively affect this species. Finally, development along the border may effectively isolate the population that occurs in California, making it more susceptible to decline than it otherwise would be.

Status Determination

Masticophis fuliginosus has an extremely small range in California that occurs entirely in an area with substantial urban, military, and agricultural development. However, the extent of extirpation and population decline is poorly documented, resulting in a Priority 3 Species of Special Concern designation.

Management Recommendations

Protection of remaining habitat in San Diego County is necessary to prevent further declines or extirpations. Minimizing urban edge effects by creating habitat buffers may benefit populations, particularly those living in small habitat fragments (Mitrovich et al. 2009). Given the very small range of the species, it may be possible to reduce road mortality with wildlife tunnels and associated drift fences installed beneath high-traffic roads in key areas important for population connectivity.

Monitoring, Research, and Survey Needs

Drift fence arrays with funnel traps have been successfully used to document the presence/absence of *Masticophis fuliginosus* in California (Fisher and Case 2000), and mark-recapture data to establish population sizes are essential for future management. Additional genetic data would complement existing mitochondrial DNA data and radiotelemetry research, respectively. Specifically, such data are needed to further resolve the taxonomic status of this snake and to provide information on landscape-level population structure. The efficacy of road-crossing structures should be investigated for this species.



COAST PATCH-NOSED SNAKE

Salvadora hexalepis virgulata Bogert 1935

Status Summary

Salvadora hexalepis virgulata is a Priority 2 Species of Special Concern, receiving a Total Score/Total Possible of 54% (46/85). During the previous evaluation, it was also designated as a Species of Special Concern (Jennings and Hayes 1994a).

Identification

Salvadora hexalepis virgulata is a medium-sized (to 115 cm TL) snake, with an enlarged rostral scale, large eyes, and a light middorsal stripe 1.5–2 scale rows in width (Jennings and Hayes 1994a, Stebbins 2003). The dorsal stripe is yellowish to grayish and extends from the tail to the rear of the neck region (Bogert 1935, Stebbins 2003). The sides of the body are dark brown, and the head is olive or brown (Bogert 1935, Perkins 1938). Ventral coloration is generally a dull white, often with an orange wash that is more prominent toward the underside of the tail (Stebbins 2003, Lemm 2006). Usually only a single supralabial contacts the eye (Bogert 1935).

The presence of both a conspicuously enlarged rostral scale and a middorsal stripe easily differentiate this species from all other snakes within its range. Along the eastern edge

Coast Patch-Nosed Snake: Risk Factors

Ranking Criteria (Maximum Score)	Score
i. Range size (10)	5
ii. Distribution trend (25)	15
iii. Population concentration/ migration (10)	0
iv. Endemism (10)	3
v. Ecological tolerance (10)	3
vi. Population trend (25)	Data deficient
vii. Vulnerability to climate change (10)	10
viii. Projected impacts (10)	10
Total Score	46
Total Possible	85
Total Score/Total Possible	0.54




PHOTO ON PREVIOUS PAGE: Coast patch-nosed snake, San Diego County, California. Courtesy of John Andermann.

of its range, *S. h. virgultea* is parapatric with the Mojave patch-nosed snake (*S. h. mojavensis*) and the desert patch-nosed snake (*S. h. hexalepis*), and it intergrades with both (Bogert 1945). In *S. h. mojavensis*, the supralabials usually do not reach the eye, and the dark lateral coloration is sometimes less pronounced and may be somewhat discontinuous. *Salvadora hexalepis hexalepis* usually has one supralabial reaching the eye but has a wider dorsal stripe (usually three scale rows wide) than *S. h. virgultea*. The spotted leaf-nosed snake (*Phyllorhynchus decurtatus*) occurs along the eastern edge of *S. h. virgultea*'s range and also has an enlarged rostral scale. This species differs in having a light-brown spotted pattern on the dorsal surface and attaining much smaller adult sizes (to 51 cm TL) (Stebbins 2003).

Taxonomic Relationships

Salvadora hexalepis virgultea was first recognized primarily on the basis of coloration, although some scalation characters also differentiate it from other subspecies of the *S. hexalepis* complex (Bogert 1935, Bogert 1945). Genetic analysis of differentiation among the subspecies has not been undertaken and represents a clear need for future research. The placement of *Salvadora* within the colubrine phylogeny is also uncertain and requires clarification.

Life History

Salvadora hexalepis virgultea is a medium-sized, active, diurnal snake, yet remains exceedingly poorly known. Here, we use information from California populations where we can, but largely rely on information from parts of the range outside of California as well as from other species and subspecies within *Salvadora* and assume that many aspects of the life history are similar among members of the complex. This subspecies is most active in May and June, and its normal seasonal activity period appears to extend from March until October. In Ventura County, warm (and presumably active) individuals have been found under stones

between January and March (S. Sweet, pers. comm.). Individuals found in December and January in gopher burrows and woodrat nests were cold and presumably dormant (S. Sweet, pers. comm.). Specimens have been documented in the literature in all months except January and February (Klauber 1939, Grismer 2002), and more recent (1995–2011) USGS pit-fall trap data have documented captures in January (three records) and February (one record; C. Rochester, unpublished data). Breeding has been observed in late April, with hatchlings appearing between mid-July and October (Klauber 1931, Grismer 2002), perhaps indicating that multiple clutches may be laid in a single year. Other members of the genus are oviparous, and we expect that *S. h. virgultea* is as well, although eggs have never been described (Wright and Wright 1957).

Salvadora primarily eats lizards and probably specializes on whiptails (*Aspidoscelis* spp.) (Cunningham 1959b, Grismer 2002, Lemm 2006). The diet of *S. h. virgultea* has not been described, but other members of the complex feed nearly exclusively on members of the genus *Aspidoscelis*, with other small lizards being taken only when they are moving rapidly, as *Aspidoscelis* tends to do (Bogert 1939). Several members of the complex are also known to feed on other small lizards, mammals, and snakes (Stebbins 2003). The enlarged rostral scale may be an aid for digging and possibly a specialization for feeding on lizard eggs (Grismer 2002, Lemm 2006), although this behavior has never been described in *S. h. virgultea*. *Salvadora* species have enlarged rear teeth, suggesting that they may envenomate prey. The saliva of *Salvadora* has not been studied, though Grismer (2002) observed an *A. tigris* that died soon after being bitten by a Baja California patch-nosed snake (*S. h. klauberi*).

The movement ecology of *S. h. virgultea* is unknown. Like other members of the genus, it is a fast and active species that is probably active at higher temperatures than most other diurnal colubrids (Jacobson and Whitford 1971). The species readily climbs and has been found

foraging in brush, off the ground (Grinnell and Grinnell 1907, Lemm 2006; S. Sweet, pers. comm.). The species probably ranges widely, as do other large diurnal colubrids.

Habitat Requirements

Salvadora hexalepis virgulata shows an apparent preference for brushy chaparral habitat (Bogert 1935, Grismer 2002). Klauber (1939) located 35 specimens in "Heavy brush, Chaparral" habitat, which was more than the number found in all other habitat types combined. In Baja California, they are also often seen in riparian areas in the vicinity of Tecate (Grismer 2002). In California unpublished data suggest that this species has a preference for chamise and red shank and often basks at or near the tops of bushes (S. Sweet, pers. comm.). Schoenherr (1976) noted that this taxon was widespread but uncommon on the Pacific slope of the San Gabriel Mountains and that it may prefer coastal sage scrub to chaparral, at least in this area. He recorded a sighting at 830 m elevation, the highest site for coastal sage scrub habitat in the San Gabriels. At two burn sites in southern California, brush was reduced, *Aspidoscelis* numbers increased, and *S. h. virgulata* numbers remained stable or increased (C. Rochester, pers. comm.), suggesting that prey abundance may be at least as important as vegetation per se. If the species is a wide-ranging predator, it may be susceptible to habitat fragmentation. Because a large component of its diet probably consists of *Aspidoscelis* species, *S. h. virgulata* may be susceptible to decline in areas where *Aspidoscelis* are declining. The two species found within its range, the orange-throated whiptail (*A. hyperythra*) and the coastal whiptail (*Aspidoscelis tigris stejnegeri*), are both under threat. *Aspidoscelis hyperythra* is a Watch List species, while *A. t. stejnegeri* is a Priority 3 Species of Special Concern (see accounts, this volume).

Distribution (Past and Present)

The historical distribution ranges from Ventura and Los Angeles Counties south to the United States–Mexico border and south to the

vicinity of El Rosario, Baja California, Mexico. It ranges from sea level along the coast up to 2130 m (Jennings and Hayes 1994a, Lemm 2006). The eastern edge of the range extends to the vicinity of Campo, San Diego County; Banning, Riverside County; and San Bernardino, San Bernardino County (Bogert 1935).

Today, the species is declining or absent from large areas of the Los Angeles basin and along the coast to San Diego (Jennings and Hayes 1994a; R. Fisher, pers. comm.). Survey data are available from Torrey Pines State Park where this species was formerly present. It has not been re-documented there despite intensive trapping efforts (Wells 1998).

Trends in Abundance

No historical or current abundance data exist for this taxon, and anecdotal reports are rare and inconclusive. Records in southern California are infrequent (Lemm 2006), although this taxon may be more commonly encountered in riparian areas south of the United States–Mexico border (Grismer 2002). The species may have historically been rare in California. Klauber (1924) reported only two specimens despite intensive collecting efforts spanning 2 years and stated that "it seems to be uncommon." Bogert (1930) also noted that it was "uncommon in the chaparral." USGS pitfall data for San Diego, Orange, and Los Angeles Counties collected between 1995 and 2011 indicate that 2.6% of snake captures were *Salvadora hexalepis virgulata* (123 captures out of 4680 total snake captures), compared to 3.8% (61/1601) of captures for daytime road driving records for snakes recorded by Klauber (1939). These data confirm that this taxon is relatively uncommon and may suggest a moderate decline over the last 60 years.

Nature and Degree of Threat

The declines in *Salvadora hexalepis virgulata* are most likely due to the conversion of the preferred brushy habitat to other vegetation types. Development of rangeland, combined with increasingly frequent and intense wildfires, has

converted large blocks of chaparral habitat to grassland (R. Fisher, pers. comm.), which appears to be unsuitable for this species (Jennings and Hayes 1994a; S. Sweet, pers. comm.). Climate change may exacerbate the intensity of wildfires in southern California (Cayan et al. 2008b), although current models range from a 29% decrease to a 28% increase in wildfires in the region (PRBO 2011). However, climate models for 2070 project an estimated 38–44% decrease in the chaparral/coastal scrub habitat preferred by this species. Ongoing urbanization in the populated areas within this taxon's range is also destroying, degrading, and fragmenting large areas of remaining habitat. *Salvadora hexalepis virgulata*'s probable preferred prey, *Aspidoscelis* lizards, are also in decline, which could cause cascading declines in snake populations. Finally, diurnally active widely foraging snakes are particularly affected by road mortality, and the volume of vehicular traffic in much of its range is large and increasing.

Status Determination

Salvadora hexalepis virgulata has a relatively small range in California and has disappeared from significant areas centered in the southern portion of its range where it was formerly known. It continues to lose habitat, which is also causing declines in a significant component of its prey-base. Habitat loss, due to direct anthropogenic changes, climate-change-driven habitat loss, and wildfire, is unlikely to stop in the near future, which we expect will cause further declines. For all of these reasons, a Priority 2 designation is justified.

Management Recommendations

The primary management goal for *Salvadora hexalepis virgulata* should be to protect large, intact patches of brushy chaparral and/or coastal sage scrub habitat that support this snake. Ideally, these patches should be those that are least likely to be directly affected by future climate change. Pending further study of the species' movement ecology, habitat protection efforts should focus on remaining large

blocks of intact habitat. Habitat fragmentation from roads is a key issue in these efforts. Road overcrossings, if installed for other taxa, may function as a means to avoid habitat fragmentation. A key element of effective management is to maintain large, healthy populations of *Aspidoscelis hyperythra* and *Aspidoscelis tigris stejnegeri*, since they appear to form the primary prey base of this snake. As declines in those species are also linked to habitat loss, management efforts among these species should be coordinated.

Monitoring, Research, and Survey Needs

Much remains to be learned about the life history and ecology of *Salvadora hexalepis virgulata* in California. Surveys employing pitfall traps, snake traps, and daytime surveys should be undertaken to establish baseline abundance data in remaining populations, and to ascertain whether these or other survey methods are potentially biased for this species, as has been suggested by some biologists. Because these snakes are infrequently encountered, power analyses are particularly important to determine how sensitive the surveys would be in detecting declines.

Autecological research focusing specifically on diet requirements, habitat utilization, and their interaction are badly needed, and the lack of this basic knowledge undermines our ability to effectively manage this snake. Specifically, the extent to which this taxon specializes on *Aspidoscelis*, as opposed to utilizing alternative prey, needs to be determined. Basic data on home range size and movement patterns, and how they may vary as a function of vegetation, are entirely lacking, and some anecdotal observations suggest that they may vary across the species' range. These data are necessary to determine what habitat blocks are most important and how large they should be to effectively conserve this taxon.

Variation within the *S. hexalepis* complex has not been examined since the initial species description, and a genetic analysis has never been undertaken. A three-pronged genetic

analysis is critical. First, a systematic characterization of the *Salvadora* species complex should be undertaken utilizing multiple independent nuclear markers. This will serve to clarify the reality of species and subspecies as valid evolutionary units and confirm their respective boundaries and range limits. Second, a phylogeographic study, using many independent markers, within *S. h. virgulata* is

needed to identify large-scale management units within the taxon. Finally, landscape genetic studies that identify migration corridors between fragmented blocks of habitat will inform our understanding of movement ecology in this species. These would also provide an estimate of effective population sizes, augmenting ecological studies of census population sizes in protected and unprotected areas.



TWO-STRIPED GARTER SNAKE

Thamnophis hammondi (Kennicott 1860)

Status Summary

Thamnophis hammondi is a Priority 2 Species of Special Concern, receiving a Total Score/Total Possible of 57% (63/110). During the previous evaluation, it was also considered a Species of Special Concern (Jennings and Hayes 1994a).

Identification

Thamnophis hammondi is a medium-sized snake (102 cm TL) with keeled scales and a head slightly wider than its body (Stebbins 2003). It is called the two-striped garter snake because it lacks the longitudinal middorsal stripe that typifies many garter snakes. The middorsal stripe is either entirely absent or represented only by a nuchal spot at the base of the head (Fitch 1948, Stebbins 1985). Color is highly variable in this species, but there are two primary color morphs: striped/spotted and striped/non-spotted (Larson 1984, Stebbins, 2003). Both morphs have yellowish to gray stripes on each side with a ground color of olive,

brown, or brownish gray, and both lack any red coloration dorsally or laterally. The ventral coloration is dull yellowish to orange red or salmon, with or without slight dusky markings (Stebbins 2003). The striped/spotted morph

Two-Striped Garter Snake: Risk Factors

Ranking Criteria (Maximum Score)	Score
i. Range size (10)	5
ii. Distribution trend (25)	15
iii. Population concentration/ migration (10)	10
iv. Endemism (10)	3
v. Ecological tolerance (10)	3
vi. Population trend (25)	10
vii. Vulnerability to climate change (10)	7
viii. Projected impacts (10)	10
Total Score	63
Total Possible	110
Total Score/Total Possible	0.57




PHOTO ON PREVIOUS PAGE: Two-striped garter snake, Los Angeles County, California. Courtesy of Robert Hess.

has one or two rows of small, alternately spaced dark spots on each side of the dorsum between the lateral stripes (Fitch 1940, Fox 1951, Larson 1984, Stebbins 2003). The striped/non-spotted morph either lacks dark spots on the dorsum or only has very small ones next to the lateral stripes on the anterior part of the body (Larson 1984, Stebbins 2003). Other color variants include non-striped/spotted, with no lateral stripes and one or two rows of dark spots on each side, sometimes appearing checkered, and non-striped/non-spotted (Brown 1980, Larson 1984, Stebbins 2003). A melanistic morph, sometimes with obscure lateral stripes and/or spots, occurs along the outer coast from Oceano to San Simeon State Park in San Luis Obispo County, and can be expected from Gaviota State Beach in Santa Barbara County to Monterey Bay (Bellemin and Stewart 1977, Larson 1984, Stebbins 2003). All color morphs exhibit varying degrees of light flecking dorsally due to whitish pigment on the inter-scale skin and margins of scales (Larson 1984). Dark green and dull red color morphs occur in northeastern Ventura County (Stebbins 2003).

The lack of a vertebral stripe and absence of red coloration on the head and sides distinguishes *T. hammondi* from the co-occurring aquatic garter snake (*T. atratus*), the western terrestrial garter snake (*T. elegans*), and the common garter snake (*T. sirtalis*).

Taxonomic Relationships

Like several other garter snakes, *Thamnophis hammondi* has a complex taxonomic history. This species has at various times been considered a subspecies of the Sierra garter snake (*T. couchii*) (e.g., Cooper 1870, Rossman 1979, Lawson and Dessauer 1979), the northwestern garter snake (*T. ordinoides*) (e.g., Grinnell and Camp 1917), and the western terrestrial garter snake (*T. elegans*) (e.g., Fitch 1948). Rossman and Stewart (1987) most recently elevated *T. hammondi* to full species status. McGuire and Grismer (1993) synonymized the Baja California Sur garter snake (*T. digueti*) with *T. hammondi*.

Life History

In California, *Thamnophis hammondi* can be active for much of the year and has been found from January through November (R. Hansen and R. Tremper, unpublished data in Rossman et al. 1996). Ervin and Fisher (2001) reported *T. hammondi* foraging and basking at a site in San Diego County from early February to October. Rathbun et al. (1993) conducted surveys in San Simeon State Park and nearby Pico Creek in San Luis Obispo County and found that large snakes (>30 cm) were most often observed in the summer, peaking in May and June, while smaller animals were seen from late August through early November. Surface activity appears to be strongly affected by the availability of surface water (E. Ervin and R. Fisher, unpublished data). Southern populations receive less rainfall and experience greater variation in rainfall, likely resulting in shorter and less predictable activity periods than northern populations.

Mating has been observed in the field in late March (Cunningham 1959b), and females are known to store sperm (Fox 1956, Stewart 1972). Like all members of the genus, *T. hammondi* is live-bearing, with litters produced from July to late October (Ernst and Ernst 2003). Hansen and Tremper (unpublished data in Rossman et al. 1996) documented an average of 15.6 offspring from 7 litters (range 3–36). Young were born in late July and August and were 20.3–21.7 cm TL (R. Hansen and R. Tremper, unpublished data in Rossman et al. 1996). Cunningham (1959b) found a 46.1 cm SVL female that contained 6 embryos. Another 64.9 cm SVL female contained 19 eggs (Cunningham 1959b). Males mature at 37.3 cm SVL and females at 38.8 cm (Wright and Wright 1957).

Cunningham (1966) reported a mean body temperature of 14 field-active individuals of 22.6°C (range 18.6–31.8°C). Five of these animals were swimming in water between 14°C and 27°C (Cunningham 1966). Inactive snakes found under cover objects had body temperatures ranging from 7.2°C to 23.6°C (Cunningham 1966).

One radiotelemetry study has collected data on the movement ecology of this species at San Simeon State Park (Rathbun et al. 1993). Activity ranges of radio-tracked snakes were greater and more distant from water in the winter than in the summer. Average summer activity ranges for seven adult females were $1498.9 \pm 1847.6 \text{ m}^2$ (mean \pm s.d.), although the duration of the study was short (range 4–29 days), and activity may have increased with more time. Average winter activity ranges for two females and one male were $3395.7 \pm 4803.5 \text{ m}^2$ (mean \pm s.d.), with animals tracked for 29–57 days. Average daily distance to water was 7.2 m in summer, compared to 98.8 m in winter (Rathbun et al. 1993).

Thamnophis hammondi is a generalized predator on a variety of prey including fish, fish eggs, frogs, salamanders, leeches, and earthworms (Van Denburgh 1897, Klauber 1931, Fitch 1940, Fitch 1941, Cunningham 1959b, Bell and Haglund 1978, Rathbun et al. 1993, Rodríguez-Robles and Galina-Tessaro 2006). This species will eat introduced prey, such as sunfish, African clawed frogs, and bullfrogs (Ervin and Fisher 2001, Mullin et al. 2004, Ervin and Fisher 2007).

Habitat Requirements

Thamnophis hammondi is among the most aquatic of the garter snakes and is often found in or near permanent and intermittent freshwater streams, creeks, and pools (Grinnell and Grinnell 1907, Fitch 1940; R. Hansen and R. Tremper, unpublished data in Rossman et al. 1996). Associated vegetation types include willow, oak woodlands, cedar, coastal sage scrub, sparse pine, scrub oak, and chaparral (R. Hansen and R. Tremper, unpublished data, in Rossman et al. 1996, Ernst and Ernst 2003). *Thamnophis hammondi* will also use artificial aquatic habitats such as cattle ponds (Jennings and Hayes 1994a, Ervin and Fisher 2001, Ervin and Fisher 2007).

Surveys in San Simeon State Park in San Luis Obispo County from July to December 1992 resulted in 45 snake sightings: 33.3% on

land, 53.3% on banks, and 6.7% in the water (Rathbun et al. 1993). Almost all of the sightings (44/45) were in or near pooled water sources. Sixty percent of snakes were sighted in low vegetation (e.g., herbs and grasses), 28.9% in tall vegetation (e.g., cattails), 11.1% in open areas with no vegetation, and zero in wooded areas (e.g., willow; Rathbun et al. 1993).

Habitat and movement ecology may vary seasonally, although this requires further study. *Thamnophis hammondi* have been observed to concentrate their habitat use in vernal pools in the spring and in remnant pools formed from ephemeral creeks in the summer (R. Fisher, pers. comm.). Nine radio-tracked snakes in San Simeon State Park used streamside habitats more in the summer, while chaparral and grassland upland sites were used for overwintering (Rathbun et al. 1993). Ninety-five percent of diurnal locations of radio-tracked animals were on land, usually underground. Animals were underground, presumably in rodent burrows, in 87.9% of locations on land (Rathbun et al. 1993). Two of the animals had home ranges that overlapped Highway 1 (a major highway with heavy traffic), suggesting that potential road mortality may be a management issue. Although it is generally considered to be a very aquatic snake, these observations suggest that terrestrial upland habitats and rodent burrows can be important habitat components for *T. hammondi*.

Distribution (Past and Present)

Thamnophis hammondi occurs in California from Salinas, Monterey County, south along the coast into Baja California, Mexico, occurring in the South Coast, Peninsular, and Transverse ranges (Boundy 1990, Ely 1992, McGuire and Grismer 1993). Isolated populations also occur in Baja California Sur and on Santa Catalina Island (Brown 1980, Stebbins 2003). While *T. hammondi* occurs mostly west of the deserts in California, there are populations in some perennial desert slope streams in San Bernardino, Riverside, and San Diego Counties (Perkins 1938, Fitch 1940, Boundy 1990). The

elevational range is from sea level to 2450 m (Atsatt 1913). Jennings and Hayes (1994a) estimated that *T. hammondi* has been extirpated from ~40% of its historic range in California during the second half of the twentieth century. This snake may be patchily distributed even when abundant suitable habitat is available. For example, snakes were readily observed at San Simeon Creek, San Luis Obispo County, in 1992, but similar habitat about 5 km away in Pico Creek had very few snakes, even though the latter experiences less human disturbance (Rathbun et al. 1993).

Trends in Abundance

Declines in abundance appear to be less severe in the southern compared to the northern part of the range, but few quantitative data are available to support this interpretation (Jennings and Hayes 1994a). Variation in abundance over time at a particular site may be partially explained by reduction in surface activity during drought periods and not necessarily reflect mortality and declines (E. Ervin and R. Fisher, unpublished data). *Thamnophis hammondi* were rare in Carmel River fish traps in 2003–2005 (S. Barry, unpublished data) and were never encountered in extensive fieldwork in and near the Hastings Reservation in the upper Carmel Valley from 1992 to 1998 (B. Shaffer, unpublished data). Jennings and Hayes (1994a) noted that *T. hammondi* was common only in San Diego County. However, other populations in the south may be robust, such as along the Santa Clara River in Los Angeles County, along Sespe Creek in Ventura County, and in the Angeles, Los Padres, and San Bernardino national forests (S. Barry, pers. comm.). The Santa Catalina Island population was reported as small (~30 individuals) and isolated in the 1970s (Brown 1980) and is suspected to have declined since (Jennings and Hayes 1994a).

Nature and Degree of Threat

Declines in the south are thought to be due to urbanization, reservoir construction, and flood control (Jennings and Hayes 1994a). Further

north, declines are suspected to have been caused by a combination of factors including habitat modification by livestock, predation by introduced vertebrates, loss of native prey, and drought (Jennings and Hayes 1994a). However, negative interactions with nonnative species have not been well documented, and in some cases *T. hammondi* may benefit from availability of introduced prey. Reliance on aquatic habitat and prey may contribute to drought sensitivity in this species (Jennings and Hayes 1994a; R. Fisher, pers. comm.).

Under climate change, mean annual temperatures are projected to increase throughout the range of *T. hammondi*, with warmer winters and summers and earlier spring warming expected (reviewed in PRBO 2011). There is less certainty about future precipitation patterns, with estimates ranging from little change to roughly 30% decreases in rainfall (Snyder and Sloan 2005, PRBO 2011). Snowpack reductions of up to 90% are predicted in the southern part of the range (Snyder et al. 2004). Warmer and potentially drier conditions may affect availability of intermittent and ephemeral water bodies and therefore limit activity. In the more northern part of the range, the probability of large (>200 ha) fires is expected to increase (Westerling and Bryant 2008) and the area burned is expected to increase by up to 50% (Lenihan et al. 2008). Both increases and decreases in fire probability and extent have been predicted for southern California under different climate change scenarios. There is little consensus on future fire dynamics in this part of the range because of the difficulty in modeling Santa Ana weather events (Westerling et al. 2004, Westerling and Bryant 2008). How *T. hammondi* may respond to fire needs to be studied. Fire may have direct mortality effects and may alter aquatic and terrestrial habitat quality. Predicted vegetation shifts due to climate change include decreases in chaparral, shrubland, and woodland, and increases in grassland area (Lenihan et al. 2008, PRBO 2011). The impact of these shifts on *T. hammondi* populations will likely be negative.

Status Determination

Thamnophis hammondi has undergone declines and extirpations and occurs in an area of high human population density and development, resulting in a Priority 2 Species of Special Concern status.

Management Recommendations

Given this species' association with aquatic habitat and apparent willingness to use artificial habitats, restoration of aquatic habitat and supplementation with artificial wetlands should be explored as a management option in extirpated sites. Eradication efforts aimed at nonnative aquatic species should consider the potential effect on *T. hammondi* populations, particularly if native prey is not abundant (Mullin et al. 2004). Rathbun et al. (1993) documented the use of upland terrestrial habitat by *T. hammondi*, and the potential importance of rodent burrows for overwintering. In order to maintain access to these habitats, they suggested protecting terrestrial habitats within

500 m of aquatic habitats, although additional study across habitat types is needed.

Monitoring, Research, and Survey Needs

Surveys to determine the abundance and distribution of remaining populations are needed and they should be conducted by individuals that are well trained to distinguish among *Thamnophis* species. Additional data on movement ecology and habitat requirements are also necessary to facilitate the design of protected areas around known aquatic habitats and to inform possible restoration efforts. The degree to which *T. hammondi* is dependent upon introduced prey should be assessed, and the quality of those introduced prey compared to native prey should be evaluated. It may be necessary to manage for both *T. hammondi* and native prey populations simultaneously for effective recovery. Finally, landscape genetic data on the degree of differentiation at the regional and watershed levels would be valuable both for the identification of management units and for possible repatriation efforts.



COMMON GARTER SNAKE, SOUTHERN POPULATIONS

Thamnophis sirtalis (Linnaeus 1758)

Status Summary

Thamnophis sirtalis is a Priority 1 Species of Special Concern, receiving a Total Score/Total Possible of 72% (72/100). During the previous evaluation, garter snakes in this part of the range were also considered Species of Special Concern (Jennings and Hayes 1994a).

Identification

Southern coastal populations of *Thamnophis sirtalis* have not been formally described as a distinct taxon, so we limit our description here to *T. sirtalis* in general. *Thamnophis sirtalis* is a medium-sized species, with a head slightly wider than the neck and keeled dorsal scales (Stebbins 2003). *Thamnophis sirtalis* can reach up to 128 cm TL in California, with adult males from coastal California 46.3 cm SVL on average and females 58.0 cm SVL on average (J. Boundy, unpublished data). Color pattern varies widely in this species, but garter snakes typically have a dark dorsal background color with lighter dorsal and lateral stripes which can

be faint or absent. California *T. sirtalis* tend to have red or orange coloration on the head and/or sides (Stebbins 2003). *Thamnophis sirtalis* in the southern part of its California range

Common Garter Snake, Southern Populations: Risk Factors

Ranking Criteria (Maximum Score)	Score
i. Range size (10)	10
ii. Distribution trend (25)	25
iii. Population concentration/ migration (10)	Data deficient
iv. Endemism (10)	10
v. Ecological tolerance (10)	7
vi. Population trend (25)	10
vii. Vulnerability to climate change (10)	3
viii. Projected impacts (10)	7
Total Score	72
Total Possible	100
Total Score/Total Possible	0.72




PHOTO ON PREVIOUS PAGE: Common garter snake, Orange County, California. Courtesy of Jeff Lemm.

potentially co-occurs with the coast garter snake (*Thamnophis elegans terrestris*) and the two-striped garter snake (*T. hammondi*) (Stebbins 2003).

Taxonomic Relationships

Some sources consider coastal garter snake populations from southern California to be California red-sided garter snakes (*Thamnophis sirtalis infernalis*) (e.g., Barry and Jennings 1998, Stebbins 2003) and others refer to them as red-spotted garter snakes (*T. s. concinnus*) (e.g., Boundy and Rossman 1995, Janzen et al. 2002). Jennings and Hayes (1994a) based their evaluation of the putative taxon, the South Coast garter snake, on personal communications with J. Boundy and S. Sweet. Morphological and genetic studies that will help to clarify the status of this taxon are still pending. One study is a comparison of color and morphological variation among *T. sirtalis* from the south coast, central coast, and central valley of California (E. Ervin, pers. comm.; C. Mahrtdt, pers. comm.). A phylogeographic study by Janzen et al. (2002) looked at *T. sirtalis* populations along the west coast of the United States but excluded populations from southern California. Another study sequencing two populations of southern *T. sirtalis* is underway for comparison with Janzen et al.'s (2002) study (R. Fisher, pers. comm.).

Life History

While the species *Thamnophis sirtalis* is one of the most well-studied North American snakes (Rossman et al. 1996), very little is known about populations from southern coastal California. Given that *T. sirtalis* is extremely widespread, occurring throughout much of Canada and in all but one state in the continental United States, life history variation among populations is pronounced (reviewed in Rossman et al. 1996, Ernst and Ernst 2003). We therefore limit our discussion to very general *T. sirtalis* biology and documented information from California where possible.

Like all members of the genus *Thamnophis*, young are live-born from midsummer to early

fall. Cunningham (1959b) reported that an 89 cm SVL female *T. sirtalis* from Tapia Park, Los Angeles County, gave birth in late August to 20 young (18 live) that were about 25 cm in TL. Another 59 cm SVL female from the same area was carrying 12 embryos (Cunningham 1959b). A single female from farther north in San Benito County gave birth to six young, also in late August (Banta and Morafka 1968). Elsewhere, average litter sizes range from 7.6 in British Columbia (Gregory and Larsen 1993) to 32.5 in Maryland (McCauley 1945). Neonates range in size from 15 cm SVL (Manitoba; Gregory 1977, Gregory and Larsen 1993, Larsen et al. 1993) to 20 cm SVL (Lassen County, California; Jayne and Bennett 1990). Females mature at SVL of 43–57 cm, and males mature at 36–38 cm SVL, although this trait is highly variable across populations (Rossman et al. 1996). Time to maturity can take up to 4 years in some populations (e.g., Lassen County; Jayne and Bennett 1990). The proportion of females that are reproductively active in a given year ranges from 29% to 88% across populations (summarized in Rossman et al. 1996), suggesting that not every female breeds every year.

Thamnophis sirtalis can be active year-round in some southern localities (e.g., the Florida Everglades; Dalrymple et al. 1991). R uthling (1915) anecdotally reported that *T. sirtalis* was rarely encountered around Los Angeles in the winter. Hansen and Tremper (unpublished data in Rossman et al. 1996) note that lowland California *T. sirtalis* are active from February to October, but there is a post-August drop in activity associated with a seasonal reduction in aquatic habitat. Most activity is diurnal, although crepuscular and nocturnal activity has been observed when anurans, a primary prey item, are breeding (Ernst and Ernst 2003). Nocturnal activity has also been observed in lowland California, with *T. sirtalis* active at night during warm rains (R. Hansen and R. Tremper, unpublished data in Rossman et al. 1996).

Thamnophis sirtalis are generalized predators (reviewed in Rossman et al. 1996).

However, diet data are not available from the southern range of *T. sirtalis* in California. Data from northern populations show that anurans are a large part of the diet. Anurans were the most common prey observed eaten by California *T. sirtalis*, comprising 58% of prey items ($n = 48$ snakes, localities include Siskiyou and Humboldt Counties; Fitch 1941). Also consumed were earthworms (24% of prey items), and rarer prey (5% or less of prey items) such as fish, leeches, and slugs (Fitch 1941). In northern California at Eagle Lake (Lassen County, 1555 m), regurgitation of 36 adults revealed that 33% of individuals contained anurans (mostly western toads, *Bufo boreas*), and 90% of prey items were anurans (Kephart and Arnold 1982). Fish (6% of animals, 2% of prey items) and leeches (11% of animals and 8% of prey items) were taken less frequently (Kephart and Arnold 1982). In the northern Sierra Nevada near Truckee, Nevada County, anurans comprised 56% of prey volume (mostly Pacific tree frogs, *Pseudacris regilla*), while 33% of prey volume was fish. Rarer prey items (5% or less of total prey volume) included mice and leeches ($n = 88$ snakes; White and Kolb 1974). Juvenile *Thamnophis sirtalis* in California have also been observed to consume newly metamorphosed newts (*Taricha torosa*) (S. Barry, unpublished data).

Habitat Requirements

Thamnophis sirtalis in southern California is thought to be restricted to marsh and upland habitats near permanent water and riparian vegetation (Grinnell and Grinnell 1907, Fitch 1941, Von Bloeker 1942; S. Sweet, pers. comm., in Jennings and Hayes 1994a). Data are scarce, but habitat preferences may be quite narrow. Some observational data suggest that this taxon may avoid restored marshlands, although the reasons for this are not clear (R. Fisher, pers. comm.).

Distribution (Past and Present)

Thamnophis sirtalis was historically known from scattered localities along the southern coastal

plain from the Santa Clara River Valley in Ventura County to around San Pasqual in San Diego County (Klauber 1929, Jennings and Hayes 1994a; S. Sweet, pers. comm. in Jennings and Hayes 1994a; E. Ervin and C. Mahrtdt, unpublished data). The historical elevation range is thought to be from near sea level at Ballona Creek and Playa del Ray Marsh in Los Angeles County to ~832 m at Lake Henshaw in San Diego County (Von Bloeker 1942; R. Fisher, pers. comm. in Jennings and Hayes 1994a). Jennings and Hayes (1994a) estimated that 75% (18/24) of historic localities no longer supported populations due to anthropogenic and natural habitat loss (e.g., urbanization, flooding). Of the six extant localities identified by Jennings and Hayes (1994a), it is now suspected that populations remain in only three localities, with possible extirpations including Camp Pendleton and San Luis Rey (R. Fisher, pers. comm.).

Trends in Abundance

Historical accounts suggest that *Thamnophis sirtalis* was once quite common (Grinnell and Grinnell 1907, Bogert 1930, Von Bloeker 1942). Current populations are thought to be abundant at Lake Henshaw in San Diego County, rare along the Santa Clara River, and virtually extirpated elsewhere (S. Barry, pers. comm., R. Fisher, pers. comm.).

Nature and Degree of Threat

Extirpations and population declines in this taxon have been attributed to habitat loss and fragmentation due to urbanization, agriculture, and flood control projects, as well as natural events such as floods and droughts (De Lisle et al. 1986, Jennings and Hayes 1994a). At remaining sites, urbanization in Riverside County continues to impact the Santa Margarita River wetlands at Camp Pendleton, and increased dam height in the Prado Basin may have a negative flooding impact (R. Fisher, pers. comm.). Introduced aquatic predators and water snakes (genus *Nerodia*) may also negatively impact *Thamnophis sirtalis* (Jennings and Hayes 1994a; R. Fisher, pers. comm.).

Under climate change, mean annual temperatures are projected to increase throughout the southern California range of *T. sirtalis*, with warmer winters and summers and earlier spring warming expected (reviewed in PRBO 2011). There is less certainty about future precipitation patterns, with estimates ranging from little change to roughly 30% decreases in rainfall (Snyder and Sloan 2005, PRBO 2011). Snowpack reductions of up to 90% are predicted in southern California (Snyder et al. 2004). Warmer and potentially drier conditions may affect availability of intermittent and ephemeral water bodies and therefore limit activity. Increases and decreases in fire probability and extent have been predicted for southern California. There is little consensus on future fire dynamics because of the difficulty in modeling Santa Ana weather events (Westerling et al. 2004, Westerling and Bryant 2008). How *T. sirtalis* responds to fire is unknown. Fire may have direct mortality effects, and may alter aquatic and terrestrial habitat quality. Predicted vegetation shifts due to climate change include decreases in chaparral and shrubland and increases in grassland area (Lenihan et al. 2008, PRBO 2011). The potential impact of such vegetation shifts on *T. sirtalis* populations is unknown.

Status Determination

Thamnophis sirtalis in southern California has a very small range in a heavily human-impacted part of the state. In addition, these populations have been extirpated from most of their histori-

cal range, which justifies a Priority 1 Species of Special Concern designation.

Management Recommendations

Given the paucity of ecological information on southern populations, it is difficult to make management recommendations beyond the protection of existing habitat at this time. Future management strategies may include removal of invasive animals and plants, restoration of flow regimes, and repatriation of extirpated sites. The research needs outlined below will help to inform the eventual development of a management strategy for this taxon.

Monitoring, Research, and Survey Needs


Almost no ecological or life history information is available for this taxon, and this data gap needs to be addressed at the few remaining sites in southern California where *Thamnophis sirtalis* persists. Monitoring to determine population abundance and to verify extirpation is needed across sites. As remaining habitat is identified and extant populations are found and stabilized, human-mediated repatriation, perhaps in combination with captive breeding, may be the most effective strategy to repopulate extirpated sites. Studies on movement and dispersal are needed to determine connectivity among remaining populations, and genetic studies on both the differentiation of this taxon from other *T. sirtalis* populations and the level of among-population variability are needed. Finally, the importance and impacts of nonnative species as predators and prey should be investigated further.



WESTERN POND TURTLE

Emys [= *Actinemys*] *marmorata* Baird and Girard 1852

PHOTOS: (top) Western pond turtle, Solano County, California. Courtesy of Adam Clause.
(bottom) Western pond turtle, Santa Barbara County, California. Courtesy of Robert Hansen.



Status Summary

Emys marmorata is a Priority 1 Species of Special Concern in the southern part of the range (roughly corresponding to the range of the southwestern pond turtle, *E. m. pallida*) and a Priority 3 Species of Special Concern elsewhere (roughly corresponding to the range of the northwestern pond turtle, *E. m. marmorata*; see below for additional detail). These two populations received a Total Score/Total Possible of 81% (89/110) and 65% (71/110), respectively. During the previous evaluation, both populations were considered Species of Special Concern, also with different overall levels of threat (Jennings and Hayes 1994a).

Identification

Emys marmorata is a small to medium-sized (generally 17–18 cm, rarely to 24 cm, straight carapace length) brown, tan, or olive turtle (Stebbins 2003). The carapace is low, keelless, and often marked with a pattern of dark lines and/or dots, sometimes forming a pattern that radiates from the centers of each scute. The posterior edge of the carapace forms a smooth, non-serrated rim. In some individuals, the carapace has no patterning. The plastron is lighter tan or beige, hingeless, and often marked with

dark blotches (Stebbins 2003). The shell shape varies among habitat types, with turtles from foothill streams being flatter and narrower than individuals occurring at lower elevations in canals and sloughs (Lubcke and Wilson 2007).

This species is unlikely to be confused with other turtles within its range with the possible exception of melanistic individuals of the non-native red-eared slider (*Trachemys scripta elegans*). This latter species has a much shorter tail, attains larger overall body sizes, and has a serrated rim around the posterior edge of the carapace. Most individuals of this species also have prominent yellow stripes on the neck and shell and a broad red stripe over the temporal region of the head, although older individuals often develop an overall dark melanistic coloration.

Taxonomic Relationships

Emys marmorata is a member of the family Emydidae, which encompasses the majority of North American turtle species. The relationships within this group have undergone extensive revision in recent years, leading to many taxonomic changes and some instability. Formerly, this species was included in the genus

Northern Western Pond Turtle: Risk Factors		Southern Western Pond Turtle: Risk Factors	
Ranking Criteria (Maximum Score)	Score	Ranking Criteria (Maximum Score)	Score
i. Range size (10)	5	i. Range size (10)	5
ii. Distribution trend (25)	15	ii. Distribution trend (25)	25
iii. Population concentration/ migration (10)	10	iii. Population concentration/ migration (10)	10
iv. Endemism (10)	7	iv. Endemism (10)	7
v. Ecological tolerance (10)	0	v. Ecological tolerance (10)	0
vi. Population trend (25)	20	vi. Population trend (25)	25
vii. Vulnerability to climate change (10)	7	vii. Vulnerability to climate change (10)	7
viii. Projected impacts (10)	7	viii. Projected impacts (10)	10
Total Score	71	Total Score	89
Total Possible	110	Total Possible	110
Total Score/Total Possible	0.65	Total Score/Total Possible	0.81

Clemmys along with the bog turtle (now *Glyptemys muhlenbergii*), the wood turtle (now *G. insculpta*), and the spotted turtle (now *C. guttata*). Recent molecular analyses have suggested a close relationship between *E. marmorata*, Blanding's turtle (*Emys* [= *Emydoidea*] *blandingii*), and the European pond turtles (*E. orbicularis* and *E. trinacris*) (Bickham et al. 1996, Burke et al. 1996, Feldman and Parham 2002, Spinks and Shaffer 2009, Spinks et al. 2009). This species is now generally placed in either the monotypic genus *Actinemys* (Holman and Fritz 2001) or the genus *Emys* (the arrangement that we follow here).

Intraspecific variation within *E. marmorata* is also undergoing intensive study. Two subspecies have traditionally been recognized, *E. m. marmorata* (Baird and Girard 1852) and *E. m. pallida* (Seeliger 1945). These subspecies were initially distinguished by the presence or absence of inguinal scutes in the shell and coloration of the throat and neck. Subsequent studies also detected substantial morphological variation present across the range (Holland 1992a). Genetic analyses of intraspecific variation suggest that substantial variation is present, which is generally, but not precisely, concordant with the traditionally defined subspecies (Spinks and Shaffer 2005, Spinks et al. 2010). Spinks et al. (2014) analyzed a large panel of SNPs and concluded that *E. m. sensu lato* should be divided into two species. Because this arrangement is very recent, here we follow the earlier arrangement (of a single species) but consider threats separately for southern and northern populations as was done by Jennings and Hayes (1994a).

Life History

Emys marmorata is a highly aquatic species and basks frequently. In the northern part of the range (particularly at higher elevations), this species enters a period of dormancy throughout much of the winter. It is one of relatively few emydid turtles that regularly overwinter on land (Ultsch 2006), perhaps as a mechanism to avoid mortality from increased winter water

flows in the Mediterranean climate. Where it overwinters terrestrially, the species uses a variety of habitat types but chooses sites above the normal high water mark and burrows into loose soils and leaf litter (Reese 1996). In aquatic habitats that experience little change in water level (lakes, ponds, and reservoirs), pond turtles are known to overwinter in the water and will choose undercut banks, bottom mud, "snags" of downed wood, or rocks (Nussbaum et al. 1983, Ernst and Lovich 2009). Movement to overwintering sites occurs at the end of summer, most often in September, although the timing varies with the particular habitat and area (Reese 1996, Reese and Welsh 1997). In warmer areas, particularly in the southern part of the range, this species may remain active year-round.

Western pond turtles are known to mate throughout the spring, summer, and fall. Nesting usually occurs in the spring or early summer, although double clutching has been reported from several parts of the range (Goodman 1997, Germano and Bury 2001, Germano and Rathbun 2008, Scott et al. 2008). Females usually select nest sites within 100 m of a water body, although nests as far away as 500 m have occasionally been reported (Storer 1930, Holland 1994, Reese 1996, Holte 1998, Lovich and Meyer 2002). Clutch sizes vary from 1 to 13 eggs and vary depending on local conditions (Holland 1994, Lovich and Meyer 2002, Germano and Rathbun 2008). The eggs hatch in the fall and, at least in the northern part of the range, hatchlings often remain in the nest through the first winter, emerging the following spring (Holland 1994).

The diet is generalized and consists of a variety of small aquatic invertebrates (including insects, crustaceans, and mollusks) and a wide variety of algae and other plant material (Bury 1986). Carrion and small vertebrates are also occasionally consumed (Bury 1986). Growth rates vary widely depending on local conditions but appear to be highest in hatchlings and then gradually slow in adults. Reproductive maturity is widely variable and appears

to be linked to size. Females generally mature at slightly over 13 cm SCL as young as 4–5 years of age, while males mature at about 12.5 cm SCL at 6–8 years of age (Holland 1994, Reese 1996, Germano and Bury 2001, Germano and Rathbun 2008, Germano and Bury 2009; T. Engstrom, pers. comm.), although maturation can happen more quickly depending on local conditions in some areas (e.g., Germano 2010).

Habitat Requirements

Emys marmorata is generalized in its habitat requirements, occurring in a broad range of aquatic water bodies including flowing rivers and streams, permanent lakes, ponds, reservoirs, settling ponds, marshes, and other wetlands. This species will also temporarily use semipermanent or ephemeral water bodies, including stock ponds, vernal pools, and seasonal wetlands (Stebbins 2003, Bury and Germano 2008). This species will also at least occasionally enter sea water (Stebbins 1954, Holland 1989). Pond turtles require upland habitat that is suitable for nesting and overwintering use. Localized soil conditions, as well as the frequency and degree of disturbance in the upland habitat, probably limit their distribution. Soils need to be loose enough to allow nest excavation, while disturbance needs to be infrequent enough or of sufficiently low intensity that nests are not disturbed (Ernst and Lovich 2009).

This species is most frequently found in quiet reaches that experience little human impact and have abundant basking substrate in the form of downed wood and large rocks (Bury and Germano 2008, Thomson et al. 2010). The species can persist, at least over moderate periods of time, in highly modified habitats with high human traffic and/or little basking substrate (Spinks et al. 2003, Germano 2010).

Distribution (Past and Present)

Emys marmorata ranges widely along the Pacific coast from western Washington to the northern part of the Baja California Peninsula in Mexico. Within California, the species

ranges from the Pacific coast inland to the Sierra Nevada foothills up to elevations of 2048 m (Ernst and Lovich 2009). Further south, it ranges from the coast inland to the peninsular ranges. Scattered populations exist in the Mojave River (e.g., Victorville, Camp Cady, and Afton Canyon, San Bernardino County, California) and in some Great Basin drainages including the Susan River (Lassen County, California), and the Truckee and Carson Rivers (Nevada, possibly extending into Nevada County, California, although this has not been documented) (Holland 1992b, Lovich and Meyer 2002). Additional scattered populations are known from the Klamath Basin (R. Bury, pers. comm.). Some or all of these populations could represent introductions. One hundred and eighty individuals of this species were introduced in the state of Nevada in 1887, and these may be the source of the population in the Truckee and/or Carson Rivers (Gary 1889).

Within *E. marmorata*, the southern subspecies (*E. m. pallida*) extends from the southern range edge in Baja California, Mexico, northward in the Coast Range to San Francisco Bay, while the northern subspecies (*E. m. marmorata*) extends from San Francisco Bay north through the Sacramento Valley and Coast Range to the northern range limit in Washington. A large intergrade zone between the two subspecies has been hypothesized to exist in the San Joaquin Valley (Seeliger 1945), although recent work has shown that this area is genetically a member of the northern subspecies (Spinks et al. 2014). The populations that we recognize correspond to these subspecies distributions.

In the north, large and relatively intact populations still exist through large areas of the Coast Range and Sierra foothills, although agriculture and habitat modification have destroyed large areas of riparian and wetland habitat in the Sacramento Valley that almost certainly supported large populations of this species in the past. Scattered populations remain throughout the Sacramento Valley, but the extensive marsh habitat that dominated

much of the valley floor has been largely drained and converted to agriculture. Kelly et al. (2005) estimated that the extent of wetland habitat in the Central Valley has declined by ~80% since the 1860s when large-scale land conversion began, and this undoubtedly eliminated many *E. marmorata* populations. Holland (1992b) argued that the San Joaquin River drainage formerly represented the stronghold of this species, supporting vast numbers of individuals, and that the species has been lost from >99% of its range in the region. Overall, the number of viable populations in this area has clearly decreased, but some do remain (Holland 1992a, Jennings and Hayes 1994a, Germano 2010, Bury et al. 2012).

In the south, extensive urbanization and land conversion have caused precipitous population declines. A large fraction of remaining habitat in southern California exists as patches surrounded by large tracts of unsuitable habitat that have little suitable upland nesting habitat. Dispersal corridors between adjacent habitats have also been mostly severed by intervening urban development and heavily used roadways, resulting in heavy mortality on females searching out nest sites (R. Fisher, pers. comm.).

Trends in Abundance

Emys marmorata was formerly abundant throughout much of California. Bogert (1930) reported that *E. marmorata* was "common in larger streams along the coast and in many of the marshes adjacent to the coast," and many of these habitats still support relatively large populations (Jennings and Hayes 1994a, Germano and Rathbun 2008, Thomson et al. 2010). Elsewhere declines have occurred, particularly in southern California. Van Denburgh (1922) reported that the species was "abundant on the west fork of the San Gabriel River," but recent reports suggest that the species has declined precipitously in this area and in the Los Angeles Basin in general (Brattstrom 1988, Jennings and Hayes 1994a). Large, relatively intact populations remain through much of the northern Coast Ranges, although areas in the Central

Valley and southern California that still support the species have severely declined (Holland 1992b, Jennings and Hayes 1994a). Populations that remain in the Central Valley are undoubtedly smaller and more fragmented than they once were due to the large-scale land conversion that occurred in this area beginning in the 1860s. Further, *E. marmorata* were harvested commercially for many years, selling for 3–6 dollars per dozen in San Francisco markets during the 1920s and 1930s (Pope 1939, Nussbaum et al. 1983). The overall extent of declines in abundance caused by market collection is poorly understood. However, localized declines due to market collection were noted as early as 1879 in Sacramento (Lockington 1879), and the species' life history would make it particularly susceptible to declines from intense adult mortality.

Some published and ongoing surveys suggest that population sizes are stable in several remaining populations in the southern part of the range. In particular, southern populations near Gorman, Fresno, and along the central coast of California appear to be stable in abundance with a population structure that indicates continued breeding (Germano 2010; D. Germano, pers. comm.). Unpublished field data also indicate that the species persists in some numbers throughout Merced (particularly east of Gustine) and Fresno Counties, as well as some areas of Kern County (S. Barry, pers. comm.). At least in some areas, ongoing declines in abundance may have slowed or stopped. If additional data corroborate these observations, a decrease in the population trend scores may be warranted during the next Species of Special Concern evaluation.

Nature and Degree of Threat

The largest threats currently facing *Emys marmorata* are land use changes and fragmentation of existing habitat, as well as possible impacts via competition and predation by introduced species.

Throughout the range of *E. marmorata*, extensive wetland habitats that once supported

large numbers of this species have declined in extent and quality. Ongoing land use conversion to agriculture as well as urban development have degraded and fragmented habitat throughout virtually all of this taxon's range. These effects are most pronounced in southern California, where relatively few viable populations of this species now remain. Even in northern California, land use changes are having impacts. Reese and Welsh (1998) documented changes in the age structure of *E. marmorata* populations as a result of damming in the Trinity River drainage, suggesting negative impacts on juvenile turtles and therefore recruitment in populations affected by dams.

The impact of introduced species is largely unknown but could potentially be detrimental in several ways. The red-eared slider is widely established throughout the range of *E. marmorata* and may serve as a disease vector and competitor (Bury 2008a). The spiny softshell turtle (*Apalone spinifer*) is a more recent introduction to the Central Valley of California and is now breeding in at least one site in the Sacramento Valley (L. Patterson, pers. comm.). If this species becomes invasive on a larger scale, it is also likely to compete with and possibly prey on small *E. marmorata*. In Southern California, the range of these two species appears not to overlap, suggesting that softshells may have strong impacts on pond turtles (R. Fisher, pers. comm.). Additional introduced species that may affect *E. marmorata* are bullfrogs, crayfish, and introduced centrarchids. In the Salinas River, *E. marmorata* declined following the invasion of bullfrogs in the 1970s (B. Hubbs, pers. comm.). The strength and mechanism (predation or competition) of their impact is not currently clear, and further studies are needed. Ravens, crows, raccoons, and opossums are all known predators of *E. marmorata* adults and nests. The population sizes of these human commensal species have increased through time and may also be having impacts on *E. marmorata* populations via increased predation pressure. A very important source of this decline may operate through nest

predation that leads to reduced or failed recruitment year to year (S. Sweet, pers. comm.).

The impacts of climate change on *E. marmorata* are still poorly understood but are likely to be significant. Climate simulation models project strong changes to river hydrology in California. In particular, decreasing snowpacks and a shift to earlier and stronger river flows (and increased frequency and strength of scouring floods) are likely to negatively affect habitat and could cause local extirpations (Gayan et al. 2008b). Because the habitat is now fragmented, recolonization of these areas following localized extirpations is unlikely, particularly in southern California where the habitat is the most fragmented. Importantly, the genetic data indicate that most of the genetic diversity within this species resides in southern California. Because of this, declines in this area could result in the extirpation of much of the genetic diversity that is currently present (Spinks et al. 2010, Spinks et al. 2014; R. Fisher, unpublished data).

Status Determination

Priority 1 Species of Special Concern status is justified for *Emys marmorata* in the southern portion of the range because these populations are experiencing ongoing and strong declines in distribution and abundance (although, as noted above, some evidence indicates these declines may be slowing in some areas). Further, this area contains most of the genetic diversity that has been identified within this taxon, so entire genetic lineages are at risk. In the north, populations are experiencing declines, although to date they are less severe than in the southern portion of the range. Many of the remaining populations in the north occur in habitats that are unlikely to experience land use changes on a scale that will threaten long-term survival, so we consider this segment of the range a Priority 3 Species of Special Concern.

Management Recommendations

Our recommendations follow those of Bury et al. (2012). We outline these recommenda-

tions below and refer readers to that document for additional discussion. Protecting habitat from further degradation and fragmentation is the highest priority for this species. Following this, habitat restoration, particularly that which increases connectivity between currently isolated habitats and increases the extent of setback or buffer habitat around wetlands that is suitable for nesting, is an important management priority. Efforts to reduce or control the impact of predators (especially on nests) are also an important way to maintain current populations and increase recruitment of juveniles. Formal headstarting programs may be a useful tool for repopulating areas where local extirpations have occurred but only as a last resort and if the habitat can be restored to an extent that a population can survive with little intervention. One encouraging observation is that *Emys marmorata* can live in close proximity to human disturbance, provided that they have adequate suitable basking and nesting sites.

Monitoring, Research, and Survey Needs

Further research on the impact of invasive species is needed. In particular, the impact of reared sliders, bullfrogs, and centrarchids needs to be further characterized, to understand both to what extent these species can coexist and the effects these species have on the native populations. Both nest and hatchling habitat requirements are relatively poorly characterized, and need to be clarified if the species is to persist and thrive in human-modified habitats. The effectiveness of headstarting efforts needs to be evaluated in various habitats and predation situations. Because a large amount of life history variation is present in this taxon (particularly relating to time to maturity, body size, and clutch size; e.g., Germano 2010), researchers and managers should be cautious when applying life history data collected in one population to a different population, particularly those occurring at widely different elevations, water temperatures, or habitat types.



SONORA MUD TURTLE

Kinosternon sonoriense Le Conte 1854

Status Summary

Kinosternon sonoriense is a Priority I Species of Special Concern, receiving a Total Score/Total Possible of 66% (56/85). During the previous evaluation, it was also considered a Species of Special Concern (Jennings and Hayes 1994a). It has not been recorded from its historic range along the California–Arizona border since 1962.

Identification

Kinosternon sonoriense is a small (maximum size ~17 cm SCL) black or brown turtle, with prominent barbels on the chin and neck and usually with yellow or cream mottling on the sides of the head that form broken stripes (Ernst and Lovich 2009). The plastron is lighter than the carapace, usually pale brown to yellow, with dark pigmentation along the scute seams and well-developed anterior and posterior hinges (Ernst and Lovich 2009). The overall carapace shape is oval and moderately domed. This species is unlikely to be confused with other native

California turtles, since it is the only California native that possesses barbels and the only native aquatic turtle that occurs within its range. However, many kinosternid turtle species are

Sonora Mud Turtle: Risk Factors

Ranking Criteria (Maximum Score)	Score
i. Range size (10)	10
ii. Distribution trend (25)	20
iii. Population concentration/ migration (10)	10
iv. Endemism (10)	0
v. Ecological tolerance (10)	3
vi. Population trend (25)	Data deficient
vii. Vulnerability to climate change (10)	3
viii. Projected impacts (10)	10
Total Score	56
Total Possible	85
Total Score/Total Possible	0.66




PHOTO ON PREVIOUS PAGE: Sonora mud turtle, Santa Cruz County, Arizona. Courtesy of Jeff Lemm.

difficult to distinguish, and some of these have been sporadically introduced around the state. The most common introduced kinosternid is likely the common musk (or stinkpot) turtle (*Sternotherus odoratus*). This species has two broken light stripes on each side of the head and has only a single, anterior hinge on the plastron. Other species in the genus *Kinosternon* have also been introduced (*K. flavescens* in particular; S. Sweet, pers. comm.) but are not common and will often require expert identification (Spinks et al. 2003, Spinks et al., pers. obs.). See Stebbins (2003) for additional details.

Taxonomic Relationships

Two subspecies have been described, one of which historically occurred in California. The Sonora mud turtle (*Kinosternon sonoriense sonoriense*) includes California as well as the majority of the species' range in the southwestern United States and northern Mexico. The Sonoyta mud turtle (*K. s. longifemorales*) is restricted to the Rio Sonoyta drainage in Mexico and southern Arizona (Iverson 1976). Intraspecific, including subspecific, variation has not yet been investigated genetically.

Life History

The life history of this species has not been studied in California. Life history studies in Arizona and New Mexico suggest that there is some interpopulation variation in basic life history parameters of this species. We base our life history description on work conducted primarily in Arizona and New Mexico but recognize that these data should be regarded as tentative for California populations.

Kinosternon sonoriense is active throughout the year as long as water is present, though in warmer months it may become active primarily at night (Hulse 1974, Hulse 1982). Hibernation is known to occur in high-elevation populations in New Mexico (Degenhardt et al. 1996), although it is unlikely that this occurs in California populations, which were exclusively low elevation. *Kinosternon sonoriense* aestivates terrestrially in response to seasonal drying in sev-

eral populations (Ligon and Stone 2003, Hall and Steidl 2007, Hensley et al. 2010) but elsewhere may be more closely tied to permanent water (Ligon and Peterson 2002). In Arizona, females come into reproductive condition after a minimum of 5 years or with a carapace length between 115 and 125 mm, after which they produce one to four clutches per year although this varies depending on location (Van Loben Sels et al. 1997, Ernst and Lovich 2009, Lovich et al. 2012). Females become gravid between April and September, although most frequently in June and July (Lovich et al. 2012). The developing embryos apparently require a period of cooling before development restarts in the spring (Hulse 1982, Ewert 1991, Ernst and Lovich 2009). In Arizona, hatching may be associated with the summer monsoon in late summer (van Loben Sels et al. 1997).

Kinosternon sonoriense can attain high local population densities. One population in Hidalgo County, New Mexico, contained 212 turtles (Stone 2001). Another population in Yavapai County, Arizona, reached 750 individuals/ha of aquatic habitat (Hulse 1982). Individuals are known to undertake long terrestrial movements (>1 km) when water becomes limiting (Stone 2001, Hall and Steidl 2007), and Stone (2001) found that 26% of recaptured individuals had moved overland between aquatic capture sites. In the Santa Catalina Mountains (Pima County, Arizona), where the aquatic habitat consists of small and discrete pools, the presence of two or more adult turtles of the same sex within single pools was rare, suggesting that the species may be territorial where resources are limiting (Hall and Steidl 2007).

Kinosternon sonoriense is primarily carnivorous, feeding on a variety of invertebrates. It is known to shift to omnivory in suboptimal habitat (Hulse 1974) and to feed on or scavenge small vertebrates (Stone et al. 2005, Lovich et al. 2010).

Habitat Requirements

Habitat requirements for *Kinosternon sonoriense* in California are unknown but are likely tied to

the presence of a reliable water source and a suitable prey base. Elsewhere in its range, it inhabits a wide variety of both permanent and temporary aquatic habitats including streams, creeks, stock ponds, and natural ponds (van Loben Sels et al. 1997, Ernst and Lovich 2009, Stanila 2009, Hensley et al. 2010, Stone et al. 2011). In California, it was known to enter artificial water bodies, although the long-term suitability of this habitat is unknown. Optimal habitat appears to be slow-moving, permanent water with a high density of aquatic invertebrates and a muddy bottom (Jennings and Hayes 1994a).

Distribution (Past and Present)

Historically, this species occurred in California along the Lower Colorado River drainage (Van Denburgh and Slevin 1913, Grinnell and Camp 1917, Dill 1944). La Rivers (1942) reported the northernmost record for the species in the Colorado River drainage from Clark County, Nevada. Cooper (1870) mentioned a specimen from an unspecified locality in the Colorado River Valley, collected while he was stationed at Fort Mohave, Arizona. Several more individuals were collected from the vicinity of Yuma, Arizona, and Palo Verde, California, in the early 1900s (Van Denburgh and Slevin 1913, Van Denburgh 1922). A 1942 record (SDNHM 17897) extended the western range in California to within ~20 km of Calexico, suggesting that this taxon was present in ditches and canals in the Imperial Valley for at least some period of time. Klauber (1934) indicates that it was not “yet” present in the Imperial Valley, though by 1942 it clearly was. The overall extent and timing of its expansion into the Imperial Valley is essentially unknown. In the Lower Colorado River Valley, the species was present at least until 1941 near Bard, Imperial County (SDNHM 33866).

The last published record of *Kinosternon sonoriense* in the Lower Colorado River drainage occurred on the Arizona side of the river ~1.6 km southwest of Laguna Dam on 31 March 1962 (Funk 1974, Lovich and Beaman 2008).

Turtle trapping surveys were conducted in April of 1991 throughout much of the historic California range and failed to detect the species (King and Robbins 1991). The presence of “small black turtles along the Coachella Canal” was rumored in the 1990s, but these reports were never verified and could have been misidentified *Trachemys scripta* or *Apalone spinifera* (J. Lovich, pers. comm.).

Outside of California, *K. sonoriense* ranges through much of southern Arizona, into the southwestern corner of New Mexico and south into northern Sonora and Chihuahua, Mexico, from sea level to 2040 m (Stebbins 2003, Lovich and Beaman 2008, Ernst and Lovich 2009).

Trends in Abundance

There is no information concerning historical abundance of this species in California. Only five reliable localities have been recorded in California, and historical accounts from the early twentieth century contain few data on abundance. Van Denburgh and Slevin (1913) reported that “six or eight” specimens were collected near Yuma before 1906, and stated that “whether it ascends the Colorado River above the Gila is not known.” Van Denburgh (1922) stated that the species occurred in the Lower Colorado River drainage but was aware of records only near Yuma and at Palo Verde in Imperial County. The Clark County Nevada record had not yet been reported at this time (La Rivers 1942). Dill (1944) mentioned only that this taxon occasionally stole bait from fishermen (presumably implying that it was fairly well known to fishermen). The paucity of records from California suggests that populations here may not have occurred in the high densities documented elsewhere, although this species is difficult to detect without specific trapping efforts, and it is not clear that these efforts were ever made while the species was known to be present. Thus, the historical data on abundance are inconclusive. *Kinosternon sonoriense* has not been collected in or near California in nearly 50 years, despite extensive

surveys (King and Robbins 1991). It is clear that declines, and possibly extirpation, have occurred during the last century.

Nature and Degree of Threat

The causes of decline of *Kinosternon sonoriense* in California are poorly understood, but may be associated with habitat modification and water diversion along the Colorado River and the Imperial Valley (Ohmart et al. 1988). Increased use of pesticides may have modified the available prey base, forcing the species to shift to a suboptimal herbivorous diet, which has been suggested as a factor in other *K. sonoriense* declines (King et al. 1996). The impact of introduced exotic crayfish, bullfrogs, warm water fishes, and softshelled turtles, all of which were well established around the time of *K. sonoriense* declines (Dill 1944, Lovich and Beaman 2008), is unknown, but they could plausibly have had a negative impact on *K. sonoriense*. At one site in Arizona, reduced *K. sonoriense* densities appear to be associated with the presence of introduced crayfish (Lazaroff et al. 2006).

Between 1941 and 1943, the Imperial Irrigation District burned and sprayed oil on 13,000 km of ditches and canals in the Imperial Valley in an effort to control the damage being done by spreading muskrat populations (Twining and Hensley 1943). These efforts certainly destroyed a great deal of aquatic habitat in the region, and the effect of the oil residues may have also had strong impacts on *K. sonoriense* and other taxa that disappeared from this area during the same time period (e.g., *Rana yavapaiensis*, *Bufo alvarius*).

Status Determination

A Priority 1 Species of Special Concern designation is justified by the complete absence of records for this species since the 1960s. This is the primary cause for concern. Little understanding of *Kinosternon sonoriense*'s habitat requirements or factors leading to decline in California currently exists. However, given the survey efforts that have been conducted to date, we assume that any remaining California pop-

ulations are small, fragmentary, and vulnerable to extirpation. The species may also be vulnerable to increasing temperatures and changing hydrology due to climate change.

Management Recommendations

If future surveys detect any remaining populations, initial management efforts should focus on protecting those populations while research is performed that focuses on expanding suitable habitat and rebuilding local populations. If initial estimates of population structure indicate that reproduction and/or recruitment is not occurring, a headstarting program could be effective as a stopgap measure to prevent local extirpation. Many aquatic turtles have very different habitat requirements for hatchlings and adults, and ecological studies of both age classes will almost certainly be necessary to ensure the survival of remnant native populations.

Monitoring, Research, and Survey Needs

Although surveys have been performed for *Kinosternon sonoriense* in California, these efforts are not yet comprehensive. As this species is generally easily captured using submersible turtle traps, more complete survey efforts will help to clarify the species' status in California. Areas that have not yet been systematically surveyed include the backwaters of the Colorado River below Needles and along Lake Havasu (R. Fisher, pers. comm.); Haughtelin, Ferguson, Taylor, Draper, and Walker Lakes (King and Robbins 1991); the Coachella Canal; and any riparian habitat remaining in the area of Laguna Dam, as well as at Topock Marsh in the Havasu National Wildlife Refuge. Because the Lower Colorado River segment of the species' range spanned both California and Arizona, additional surveys should be coordinated with wildlife managers in Arizona to search potential habitat on the eastern side of the Colorado River.

If surveys do detect any individuals, managers should immediately initiate a monitoring program to determine the size and stability of

the population, as well as an ecological study of population structure and life history. This will almost certainly involve individually marking turtles with shell notches and/or PIT tags and performing mark-recapture surveys to estimate population size and individual growth rate. In particular, whether, and how much, reproduction is taking place in existing populations will be critical to determine. Juvenile turtles rarely enter submersible traps; thus, alternative methods should be employed to search for them (such as seining or snorkeling). Female turtles should also be checked for eggs using either palpation or radiographs, preferably with portable field-capable digital X-ray units.

Genetic samples from the Lower Colorado River do not exist and should be collected,

should remaining populations be found. These samples will be valuable to researchers working on *Kinosternon* phylogenetics and phylogeography and will also be critical in assessing the existing diversity within remaining populations and the divergence between these and more abundant populations to the east in Arizona.

Finally, researchers should attempt to characterize differences between habitat that supports this species and nearby habitats that do not. The causes of decline are still poorly understood, so management efforts that focus on rebuilding populations must be informed with strong data on the impact of introduced predators, pesticide, and herbicide drift, introduced aquatic plants, and habitat modification on *K. sonoriense* population persistence.

This page intentionally left blank

LITERATURE CITED

- Adalsteinsson, S.A., Branch, W.R., Trape, S., Vitt, L.J., and Hedges, S.B. 2009. Molecular phylogeny, classification, and biogeography of snakes of the family Leptotyphlopidae (Reptilia, Squamata). *Zootaxa* 2240:1–50.
- Adams, M.J. 1999. Correlated factors in amphibian decline: exotic species and habitat change in western Washington. *Journal of Wildlife Management* 63:1162–1171.
- Adams, M.J. 2000. Pond permanence and the effects of exotic vertebrates on anurans. *Ecological Applications* 10:559–568.
- Adams, M.J. and Bury, R.B. 2002. The endemic headwater stream amphibians of the American Northwest: associations with environmental gradients in a large forested preserve. *Global Ecology and Biogeography* 11:169–178.
- Adams, M.J., Bury, R.B., and Swarts, S.A. 1998. Amphibians of the Fort Lewis Military Reservation, Washington: sampling techniques and community patterns. *Northwestern Naturalist* 79:12–18.
- Adams, M.J., Pearl, C.A., and Bury, R.B. 2003. Indirect facilitation of an anuran invasion by non-native fishes. *Ecology Letters* 6:343–351.
- Adams, M.J., Pearl, C.A., Galvan, S., and McCreary, B. 2011. Non-native species impacts on pond occupancy by an anuran. *The Journal of Wildlife Management* 75:30–35.
- Adams, S.B. and Frissell, C.A. 2001. Thermal habitat use and evidence of seasonal migration by Rocky Mountain tailed frogs, *Ascaphus montanus*, in Montana. *Canadian Field Naturalist* 115:251–256.
- Adest, G.A. 1977. Genetic relationships in the genus *Uma* (Iguanidae). *Copeia* 1977:47–52.
- Aguilar, A., Douglas, R.B., Gordon, E., Baumsteiger, J., and Goldsworthy, M.O. 2013. Elevated genetic structure in the coastal tailed frog (*Ascaphus truei*) in managed redwood forests. *Journal of Heredity* 104:202–216.
- Alberts A.C., Brown, T.K., Grant, T.D., Lemm, J.M., Montagne, J.P., Milroy, L.G., III, and Jackintell, L.A. 2004. Conservation of the San Diego coast horned lizard on the Southwestern Riverside County Multi-Species Reserve. Final Project Report to the Metropolitan Water District of Southern California for Agreement No. 1550. Applied Conservation Division, Zoological Society of San Diego, San Diego, CA.
- Aldridge, R.D. 1979. Seasonal spermatogenesis in sympatric *Crotalus viridis* and *Arizona elegans* in New Mexico. *Journal of Herpetology* 13:187–192.
- Altig, R. and Brodie, E.D., Jr. 1972. Laboratory behavior of *Ascaphus truei* tadpoles. *Journal of Herpetology* 6:21–24.
- Anderson, B.W. and Ohmart, R.D. 1982. Revegetation for Wildlife Enhancement along the Lower Colorado River. Bureau of Reclamation, Boulder City, NV.
- Anderson, J.D. 1960. Cannibalism in *Dicamptodon ensatus*. *Herpetologica* 16:260.
- Anderson, J.D. 1961. The courtship behavior of *Ambystoma macrodactylum croceum*. *Copeia* 1961:132–139.
- Anderson, J.D. 1967. A comparison of the life histories of coastal and montane populations of

- Ambystoma macrodactylum* in California. The American Midland Naturalist 77:323–355.
- Anderson, J.D. 1968a. A comparison of the food habits of *Ambystoma macrodactylum sigillatum*, *Ambystoma macrodactylum croceum*, and *Ambystoma tigrinum californiense*. Herpetologica 24:273–284.
- Anderson, J.D. 1968b. Thermal histories of two populations of *Ambystoma macrodactylum*. Herpetologica 24:29–35.
- Andre, S.E., Parker, J., and Briggs, C.J. 2008. Effect of temperature on host response to *Batrachochytrium dendrobatidis* infection in the mountain yellow-legged frog (*Rana muscosa*). Journal of Wildlife Diseases 44:717–720.
- Andrews, K.M., Gibbons, J.W., and Jochimsen, D.M. 2008. Ecological effects of roads on amphibians and reptiles: a literature review, in Mitchell, J.C., Brown, R.E.J., and Bartholomew, B. (Eds): Urban Herpetology. Society for the Study of Amphibians and Reptiles, Salt Lake City, UT, pp.121–143.
- Anton, T.G. 1994. Observation of predatory behavior in the regal ringneck snake (*Diadophis punctatus regalis*) under captive conditions. Bulletin of the Chicago Herpetological Society 29:95.
- Anzalone, C., Kats, L., and Gordon, M. 1998. Effects of solar UV-B radiation on embryonic development in *Hyla cadaverina*, *Hyla regilla*, and *Taricha torosa*. Conservation Biology 12:646–653.
- Arnold, S.J. and Halliday, T. 1986. Natural history notes: *Hyla regilla*. Predation. Herpetological Review 17:44.
- Ashton, D.T., Marks, S.B., and Welsh, H.H., Jr. 2006. Evidence of continued effects from timber harvesting on lotic amphibians in redwood forests of northwestern California. Forest Ecology and Management 221:183–193.
- Atsatt, S.R. 1913. The reptiles of the San Jacinto area of southern California. University of California Publications in Zoology 12:31–50.
- Baird, S.F. 1854. Descriptions of new genera and species of North American frogs. Proceedings of the Academy of the Natural Sciences of Philadelphia 7:59–62.
- Baird, S.F. 1858. Description of new genera and species of North American lizards in the Museum of the Smithsonian Institution. Proceedings of the Academy of the Natural Sciences of Philadelphia 10:253–256.
- Baird, S.F. 1859. Report upon the reptiles of the route, in Explorations and Surveys, R.R. Route from the Mississippi River to the Pacific Ocean, 1853–56, Vol. 10, Williamson's Route. Zoological Report, Part 6, Number 4, Washington, DC, pp.37–45.
- Baird, S.F. and Girard, C. 1852. Descriptions of new species of reptiles collected by the U.S. exploring expedition under the command of Capt. Charles Wilkes. Proceedings of the Academy of the Natural Sciences of Philadelphia 6:174–177.
- Baird, S.F. and Girard, C. 1853a. Catalogue of North American Reptiles in the Museum of the Smithsonian Institution. Part I: Serpents. Smithsonian Institution, Washington, DC.
- Baird, S.F. and Girard, C. 1853b. Communication by Mr. Charles Girard on behalf of Prof. Baird and himself, upon a species of frog and another toad. Proceedings of the Academy of Natural Sciences of Philadelphia 6:378–379.
- Baldwin, K.S. and Stanford, R.A. 1987. Natural history notes: *Ambystoma tigrinum californiense*. Predation. Herpetological Review 18:33.
- Balfour, P.S. and Ranlett, J. 2006. Natural history notes: *Spea hammondi*. Predation. Herpetological Review 37:212.
- Ballesteros-Barrera, C., Martínez-Meyer, E., and Gadsden, H. 2007. Effects of land-cover transformation and climate change on the distribution of two microendemic lizards, genus *Uma*, of northern Mexico. Journal of Herpetology 41:733–740.
- Bancroft, B.A., Baker, N.J., Searle, C.L., Garcia, T.S., and Blaustein, A.R. 2008. Larval amphibians seek warm temperatures and do not avoid harmful UVB radiation. Behavioral Ecology 19:879–886.
- Banta, B.H. 1963. Remarks upon the natural history of *Gerrhonotus panamintinus* Stebbins. Occasional Papers of the California Academy of Science 36:1–12.
- Banta, B.H. and Leviton, A.E. 1961. Mating behavior of the Panamint Lizard, *Gerrhonotus panamintinus* Stebbins. Herpetologica 17:204–206.
- Banta, B.H., Mahrtdt, C.R., and Beaman, K.R. 1996. *Elgaria panamintina*. Catalogue of American Amphibians and Reptiles 629:1–4.
- Banta, B.H. and Morafka, D.J. 1968. An annotated checklist of the recent amphibians and reptiles of the Pinnacles National Monument and Bear Valley, San Benito and Monterey counties, California, with some ecological observations. The Wasmann Journal of Biology 26:161–183.
- Barrows, C.W. 2006. Population dynamics of a threatened sand dune lizard. The Southwestern Naturalist 51:514–523.
- Barrows, C.W. 2012. Temporal patterns of abundance of arthropods on sand dunes. The Southwestern Naturalist 57:262–266.
- Barrows, C.W. and Allen, M.F. 2007. Persistence and local extinctions of endangered lizard *Uma*

- inornata* on isolated habitat patches. *Endangered Species Research* 3:61–68.
- Barrows, C.W. and Allen, M.F. 2009. Conserving species in fragmented habitats: population dynamics of the flat-tailed horned lizard, *Phrynosoma mcallii*. *The Southwestern Naturalist* 54:307–316.
- Barrows, C.W. and Allen, M.F. 2010. Patterns of occurrence of reptiles across a sand dune landscape. *Journal of Arid Environments* 74:186–192.
- Barrows, C.W., Allen, E.B., Brooks, M.L., and Allen, M.F. 2009. Effects of an invasive plant on a desert sand dune landscape. *Biological Invasions* 11:673–686.
- Barrows, C.W., Allen, M.F., and Rotenberry, J.T. 2006. Boundary processes between a desert sand dune community and an encroaching suburban landscape. *Biological Conservation* 131:486–494.
- Barrows, C.W., Preston, K., Rotenberry, J.T., and Allen, M. 2008. Using occurrence records to model historic distributions and estimate habitat losses for two psammophilic lizards. *Biological Conservation* 141:1885–1893.
- Barrows, C.W., Rotenberry, J.T., and Allen, M.F. 2010. Assessing sensitivity to climate change and drought variability of a sand dune endemic lizard. *Biological Conservation* 143:731–736.
- Barry, S.J. and Fellers, G.M. 2013. History and status of the California red-legged frog (*Rana draytonii*) in the Sierra Nevada, California, USA. *Herpetological Conservation and Biology* 8:456–502.
- Barry, S.J. and Jennings, M.R. 1998. *Eutaenia sirtalis tetrataenia* Cope in Yarrow, 1875 and *Coluber infernalis* Blainville, 1835 [Currently *Thamnophis s. tetrataenia* and *T. s. infernalis*]: proposed conservation of usage of the subspecific names by the designation of a neotype for *T. s. infernalis* (Case 3012). *Bulletin of Zoological Nomenclature* 55:224–228.
- Beauchamp, B., Wone, B., Bros, S., and Kutilek, M. 1998. Habitat use of the flat-tailed horned lizard (*Phrynosoma mcallii*) in a disturbed environment. *Journal of Herpetology* 32:210–216.
- Beck, D.D. 1985. The Natural History, Distribution and Present Status of the Gila Monster in Utah. Utah Division of Wildlife Resources, Nongame Section, Salt Lake City.
- Beck, D.D. 1990. Ecology and behavior of the Gila monster in southwestern Utah. *Journal of Herpetology* 24:54–68.
- Beck, D.D. 2005. *Biology of Gila Monsters and Beaded Lizards*. University of California Press, Berkeley.
- Becker, D. and Henderson, B. 2010. Notes on Owens River *Rana pipiens* Habitat Assessment. Unpublished Report prepared on April 21.
- Belden, L.K., Moore, I.T., Mason, R.T., Wingfield, J.C., and Blaustein, A.R. 2003. Survival, the hormonal stress response and UV-B avoidance in cascades frog tadpoles (*Rana cascadae*) exposed to UV-B radiation. *Functional Ecology* 17:409–416.
- Bell, J.L., Sloan, L.C., and Snyder, M.A. 2004. Regional changes in extreme climatic events: a future climate scenario. *Journal of Climate* 17:81–87.
- Bell, M.A. and Haglund, T.R. 1978. Selective predation of threespine sticklebacks (*Gasterosteus aculeatus*) by garter snakes. *Evolution* 32:304–319.
- Bellemin, J.M. and Stewart, G.R. 1977. Diagnostic characters and color convergence of the garter snakes *Thamnophis elegans terrestris* and *Thamnophis couchii atratus* along the central California coast. *Bulletin of the Southern California Academy of Sciences* 76:73–84.
- Berger, L., Speare, R., Daszak, P., Green, D.E., Cunningham, A.A., Goggin, C.L., Slocumbe, R., et al. 1998. Chytridiomycosis causes amphibian mortality associated with population declines in the rain forests of Australia and Central America. *Proceedings of the National Academy of Sciences of the United States of America* 95:9031–9036.
- Bezy, R.L. 1967. A new night lizard (*Xantusia vigilis sierrae*) from the southern Sierra Nevada in California. *Journal of the Arizona Academy of Science* 4:163–167.
- Bezy, R.L. 2009. *Xantusia sierra*, in Jones, L. and Lovich, R. (Eds): *Lizards of the American Southwest: A Photographic Field Guide*, Rio Nuevo Publishing, Tucson, AZ, pp.432–435.
- Bezy, R.L., Gorman, G.C., Kim, Y.J., and Wright, J.W. 1977. Chromosomal and genetic divergence in the fossorial lizards of the family Anniellidae. *Systematic Zoology* 26:57–71.
- Bezy, R.L. and Sites, J.W. 1987. A preliminary study of allozyme evolution in the lizard family Xantusiidae. *Herpetologica* 43:280–292.
- Bezy, R.L. and Wright, J. 1971. Karyotypic variation and relationships of the California legless lizard, *Anniella pulchra* Gray (Reptilia: Anniellidae). *Herpetological Review* 3:71–72.
- Bicket, J.C. 1982. Geographic distribution: *Meloderma suspectum cinctum*. *Herpetological Review* 13:131.
- Bickham, J.W., Lamh, T., Minx, P., and Patton, J.C. 1996. Molecular systematics of the genus *Clemmys* and the intergeneric relationships of emydid turtles. *Herpetologica* 52:89–97.

- Biek, R., Mills, L., and Bury, R. 2002. Terrestrial and stream amphibians across clearcut-forest interfaces in the Siskiyou Mountains, Oregon. *Northwest Science* 76:129–140.
- Blanchard, F.N. 1924. A new snake of the genus *Arizona*. *Occasional Papers of the Museum of Zoology* 150:1–5.
- Blanchard, F.N. 1942. The ring-neck snakes, genus *Diadophis*. *Bulletin of the Chicago Academy of Sciences* 7:1–144.
- Blaustein, A.R. and O'Hara, R.K. 1987. Aggregation behaviour in *Rana cascadae* tadpoles: association preferences among wild aggregations and responses to non-kin. *Animal Behaviour* 35:1549–1555.
- Blaustein, A.R., Romansic, J.M., Scheessele, E.A., Han, B.A., Pessier, A.P., and Longcore, J.E. 2005. Interspecific variation in susceptibility of frog tadpoles to the pathogenic fungus *Batrachochytrium dendrobatidis*. *Conservation Biology* 19:1460–1468.
- Blouin, M.S., Phillipsen, I.C., and Monsen, K.J. 2010. Population structure and conservation genetics of the Oregon spotted frog, *Rana pretiosa*. *Conservation Genetics* 11:2179–2194.
- Bobzien, S. and DiDonato, J. 2007. The status of California tiger salamander (*Ambystoma californiense*), California red-legged frog (*Rana draytonii*), foothill yellow-legged frog (*Rana boylei*), and other aquatic herpetofauna in the East Bay Regional Park District, California. Report available from EBRP, Oakland, CA.
- Bogert, C.M. 1930. An annotated list of the amphibians and reptiles of Los Angeles County, California. *Bulletin of the Southern California Academy of Sciences* 29:3–14.
- Bogert, C.M. 1935. *Salvadora grahamiae virgulata*, a new subspecies of the patch-nosed snake. *Bulletin of the Southern California Academy of Sciences* 34:88–94.
- Bogert, C.M. 1939. A study of the genus *Salvadora*, the patch nosed-snakes. Publication in *Biological Sciences*, University of California at Los Angeles 1:177–236.
- Bogert, C.M. 1945. Two additional races of the patch-nosed snake, *Salvadora hexalepis*. *American Museum Novitates* 1285:1–14.
- Bogert, C.M. and Martin del Campo, R. 1956. The Gila monster and its allies: the relationships, habits, and behavior of the lizards of the family Helodermatidae. *Bulletin of the American Museum of Natural History* 109:1–238.
- Bolster, B.C. (Ed.). 1998. Terrestrial Mammal Species of Special Concern in California. Draft Final Report prepared by P.V. Brylski, P.W. Collins, E.D. Pierson, W.E. Rainey and T.E. Kucera. Report submitted to California Department of Fish and Game Wildlife Management Division, Nongame Bird and Mammal Conservation Program for Contract No. FG3146WM.
- Bolster, B.C. 2010. A status review of the California tiger salamander (*Ambystoma californiense*). California Department of Fish and Game, Nongame Wildlife Program Report 2010-4. 99pp.
- Bolster, B.C., King, T., and Robbins, M. 2000. Distribution of Mojave and Colorado Desert fringe-toed lizards, *Uma scoparia* and *Uma notata notata*. California Department of Fish and Game, Habitat Conservation Division, Sacramento.
- Bonham, C. and Lockhart, M. 2011. A Status Review of the Mountain Yellow-Legged Frog. Report to the Fish and Game Commission, California Department of Fish and Game, Sacramento.
- Borney, R., Cooper, C.B., Dickinson, J., Kelling, S., Phillips, T., Rosenberg, K.V., and Shirk, J. 2009. Citizen science: a developing tool for expanding science knowledge and scientific literacy. *BioScience* 59:977–984.
- Boorse, G.C. and Denver, R.J. 2003. Endocrine mechanisms underlying plasticity in metamorphic timing in spadefoot toads. *Integrative and Comparative Biology* 43:646–657.
- Bostic, D.L. 1971. Herpetofauna of the Pacific Coast of north central Baja California, Mexico, with a description of a new subspecies of *Phyllodactylus xanti*. *Transactions of the San Diego Society of Natural History* 16:237–263.
- Boundy, J. 1990. Biogeography and Variation in Southern Populations of the Garter Snake *Thamnophis atratus*, with a Synopsis of the *T. couchii* Complex. Master's Thesis. California State University, San Jose.
- Boundy, J. and Balgooyen, T.G. 1988. Record lengths for some amphibians and reptiles from the western United States. *Herpetological Review* 19:26–27.
- Boundy, J. and Rossman, D.A. 1995. Allocation and status of the garter snake names *Coluber infernalis* Blainville, *Eutaenia sirtalis tetrataenia* Cope and *Eutaenia imperialis* Coues and Yarrow. *Copeia* 1995:236–240.
- Bourque, R.M. 2008. Spatial Ecology of an Inland Population of the Foothill Yellow-Legged Frog (*Rana boylei*) in Tehama County, California. Master's Thesis. Humboldt State University, Arcata, CA.
- Bowerman, J. and Pearl, C.A. 2010. Ability of Oregon spotted frog (*Rana pretiosa*) embryos

- from central Oregon to tolerate low temperatures. *Northwestern Naturalist* 91:198-202.
- Bradford, D.F., Gordon, M.S., Johnson, D.F., Andrews, R.D., and Jennings, W.B. 1994. Acidic deposition as an unlikely cause for amphibian population declines in the Sierra Nevada, California. *Biological Conservation* 69:155-161.
- Bradford, D.F., Jaeger, J.R., and Jennings, R.D. 2004. Population status and distribution of a decimated amphibian, the relict leopard frog (*Rana onca*). *The Southwestern Naturalist* 49:218-228.
- Bradford, D.F., Swanson, C., and Gordon, M.S. 1992. Effects of low pH and aluminum on two declining species of amphibians in the Sierra Nevada, California. *Journal of Herpetology* 26:369-377.
- Bradley, G.A., Rosen, P.C., Sredl, M.J., Jones, T.R., and Longcore, J.E. 2002. Chytridiomycosis in native Arizona frogs. *Journal of Wildlife Diseases* 38:206-212.
- Bradley, W.G. and Deacon, J.E. 1966. Distribution of the Gila monster in the northern Mojave Desert. *Copeia* 1966:365-366.
- Brame, A.H., Jr. 1956. The number of eggs laid by the California newt, *Taricha torosa*. *Herpetologica* 12:325.
- Brame, A.H., Jr. 1968. The number of egg masses and eggs laid by the California newt, *Taricha torosa*. *Journal of Herpetology* 2:169-170.
- Brame, A.H. and Murray, K.F. 1968. Three new slender salamanders (*Batrachoseps*) with a discussion of relationships and speciation within the genus. *Bulletin of the Natural History Museum of Los Angeles County* 4:1-35.
- Brattstrom, B.H. 1951. The number of young of *Xantusia*. *Herpetologica* 7:143-144.
- Brattstrom, B.H. 1952. The food of the nightlizards, genus *Xantusia*. *Copeia* 1952:168-172.
- Brattstrom, B.H. 1963. A preliminary review of the thermal requirements of amphibians. *Ecology* 44:238-255.
- Brattstrom, B.H. 1965. Body temperatures of reptiles. *The American Midland Naturalist* 73:376-422.
- Brattstrom, B.H. 1988. Habitat destruction in California with special reference to *Clemmys marmorata*: a perspective, in De Lisle, H.F., Brown, P.R., Kaufman, B., and McGurty, B.M. (Eds): *Proceedings of the Conference on California Herpetology*. Special Publication no. 4. Southwestern Herpetologists Society, Van Nuys, CA, pp.13-24.
- Brattstrom, B.H. 1997. Status of the subspecies of the coast horned lizard, *Phrynosoma coronatum*. *Journal of Herpetology* 31:434-436.
- Brattstrom, B.H. and Bondello, M.C. 1979. The effect of ORV sounds on the emergence of Couch's spadefoot toad, *Scaphiopus couchii*. Bureau of Land Management, Riverside, CA.
- Brattstrom, B.H. and Bondello, M.C. 1983. Effects of off-road vehicle noise on desert vertebrates, in Webb, R.H. and Wilshire, H.G. (Eds): *Environmental Effects of Off-Road Vehicles: Impacts and Management in Arid Regions*, Springer-Verlag, New York, pp.167-206.
- Brattstrom, B.H. and Warren, J.W. 1953. A new subspecies of racer, *Masticophis flagellum*, from the San Joaquin Valley of California. *Herpetologica* 9:177-179.
- Brehme, C.S. 2003. Responses of Small Terrestrial Vertebrates to Roads in a Coastal Sage Scrub Ecosystem. Master's Thesis. San Diego State University, San Diego, CA.
- Briggs, C.J., Knapp, R.A., and Vredenburg, V.T. 2010. Enzootic and epizootic dynamics of the chytrid fungal pathogen of amphibians. *Proceedings of the National Academy of Sciences* 107:9695-9700.
- Briggs, L.J. 1987. Breeding biology of the cascade frog, *Rana cascadae*, with comparisons to *R. aurora* and *R. pretiosa*. *Copeia* 1987:241-245.
- Brode, J.M. 1967. Occurrence of *Ambystoma macrodactylum* in the Warner Mountains of northeastern California. *Herpetologica* 23:315-316.
- Brown, D.E. and Carmony, N.B. 1991. Gila Monster: Facts and Folklore of America's Aztec lizard, High Lonesome Books, Silver City, NM, 130pp.
- Brown, H.A. 1967. Embryonic temperature adaptations and genetic compatibility of two allopatric populations of the spadefoot toad, *Scaphiopus hammondi*. *Evolution* 21:742-761.
- Brown, H.A. 1969. The heat resistance of some anuran tadpoles (Hylidae and Pelobatidae). *Copeia* 1969:138-147.
- Brown, H.A. 1975a. Temperature and development of the tailed frog *Ascaphus truei*. *Comparative Biochemistry and Physiology A*. 50:397-406.
- Brown, H.A. 1975b. Reproduction and development of the red-legged frog, *Rana aurora*, in northwestern Washington. *Northwest Science* 49:241-252.
- Brown, H.A. 1976. The status of California and Arizona populations of the western spadefoot toads (genus *Scaphiopus*). *Natural History Museum of Los Angeles County Contributions in Science* 286:1-15.
- Brown, H.A. 1977. Oxygen-consumption of a large, cold-adapted frog egg (*Ascaphus truei* (Amphibia-Ascaphidae)). *Canadian Journal of Zoology* 55:343-348.

- Brown, H.A. 1990. Morphological variation and age-class determination in overwintering tadpoles of the tailed frog *Ascaphus truei*. *Journal of Zoology* 220:171–184.
- Brown, T.K., Lemm, J.M., Montagne, J., Tracey, J.A., and Alberts, A.C. 2008. Spatial ecology, habitat use, and survivorship of resident and translocated red diamond rattlesnakes (*Crotalus ruber*), in Hayes, W.K., Bearman, K.R., Cardwell, M.D., and Bush, S.P. (Eds): *The Biology of Rattlesnakes Symposium*, Loma Linda University Press, Loma Linda, CA.
- Brown, T.W. 1980. Present status of the garter snake on Santa Catalina Island, in Power, D.M. (Ed.): *The California Islands: Proceedings of a Multidisciplinary Symposium*, Santa Barbara Museum of Natural History, Santa Barbara, CA, pp.585–595.
- Bryant, H.C. 1911. The horned lizards of California and Nevada of the genera *Phrynosoma* and *Anota*. *University of California Publications in Zoology* 9:1–84.
- Bryant, H.C. 1917. The leopard frog in California. *California Fish and Game* 3:91.
- Buck, J.C., Truong, L., and Blaustein, A.R. 2011. Predation by zooplankton on *Batrachochytrium dendrobatidis*: biological control of the deadly amphibian chytrid fungus? *Biodiversity and Conservation* 20:3549–3553.
- Bulger, J.B., Scott, N.J., and Seymour, R.B. 2003. Terrestrial activity and conservation of adult California red-legged frogs *Rana aurora draytonii* in coastal forests and grasslands. *Biological Conservation* 110:85–95.
- Bull, E.L. and Carter, B.E. 1996. *Tailed Frogs: Distribution, Ecology, and Association with Timber Harvest in Northeastern Oregon*. USDA Forest Service Research Paper PNW-RP-4971–11.
- Bunck, C.M., Burns, K.L., Green, D.E., Knickelbein, B.A., Schneider, J.E., Smith, D.J., and Wright, S.D. 2009. *National Wildlife Health Center: Health & Disease Surveys of Declining Amphibian Populations at Regional ARMI Monitoring Sites*. Unpublished Report.
- Bunn, D., Mummert, A., Hoshovsky, M., Gilardi, K., and Shanks, S. 2007. *California Wildlife: Conservation Challenges; California's Wildlife Action Plan*, Sacramento, CA.
- Burgess, R.C., Jr. 1950. Development of spade-foot toad larvae under laboratory conditions. *Copeia* 1950:49–51.
- Burke, R.L., Leuteritz, T.E., and Wolf, A.J. 1996. Phylogenetic relationships of emydine turtles. *Herpetologica* 52:572–584.
- Burkholder, L.L. and Diller, L.V. 2007. Life history of postmetamorphic coastal tailed frogs (*Ascaphus truei*) in northwestern California. *Journal of Herpetology* 41:251–262.
- Burrage, B.R. 1965. Notes on the eggs and young of the lizards *Gerrhonotus multicarinatus webbi* and *G. m. nanus*. *Copeia* 1965:512.
- Burt, C.E. 1931. An interpretation of certain experimental and observational data on the limbless lizard, *Anniella pulchra* Gray. *Copeia* 1931:105–106.
- Bury, R.B. 1968. The distribution of *Ascaphus truei* in California. *Herpetologica* 24:39–46.
- Bury, R.B. 1970a. A biogeographic analysis of the herpetofauna of Trinity County, California. *Journal of Herpetology* 4:165–178.
- Bury, R.B. 1970b. Food similarities in the tailed frog, *Ascaphus truei*, and the Olympic salamander, *Rhyacotriton olympicus*. *Copeia* 1970:170–171.
- Bury, R.B. 1972. Small mammals and other prey in diet of Pacific giant salamander (*Dicamptodon ensatus*). *The American Midland Naturalist* 87:524–526.
- Bury, R.B. 1986. Feeding ecology of the turtle, *Clemmys marmorata*. *Journal of Herpetology* 20:515–521.
- Bury, R.B. 2005. *Dicamptodon ensatus*, in Lannoo, M.J. (Ed.): *Amphibian Declines: The Conservation Status of United States Species*, University of California Press, Berkeley, pp.653–654.
- Bury, R.B. 2008a. Do urban areas favor invasive turtles in the Pacific Northwest, in Mitchell, J.C., Brown, R.E.J., and Bartholomew, B. (Eds): *Urban Herpetology, Society for the Study of Amphibians and Reptiles*, Salt Lake City, UT, pp.343–345.
- Bury, R.B. 2008b. Low thermal tolerances of stream amphibians in the Pacific Northwest: implications for riparian and stream management. *Applied Herpetology* 5:63–74.
- Bury, R.B. and Adams, M.J. 1999. Variation in age at metamorphosis across a latitudinal gradient for the tailed frog, *Ascaphus truei*. *Herpetologica* 55:283–291.
- Bury, R.B. and Balgooyen, T. 1976. Temperature selectivity in legless lizard, *Anniella pulchra*. *Copeia* 1976:152–155.
- Bury, R.B. and Corn, P.S. 1988. Douglas-fir forests in the Oregon and Washington Cascades: relation of the herpetofauna to stand age and moisture, in Szaro, R.C., Severson, K.E., and Patton, D.R. (Eds): *Management of Amphibians, Reptiles, and Small Mammals in North America*. General Technical Report RM-166. United States Department of Agriculture Forest Service, Rocky

- Mountain Forest and Range Experiment Station, Fort Collins, CO, pp.11–22.
- Bury, R.B., Corn, P.S., Aubry, K.B., Gilbert, F.F., and Jones, L.L.C. 1991. Aquatic amphibian communities in Oregon and Washington, in Ruggiero, L.F., Aubry, K.B., Carey, A.B., and Huff, M.H. (Eds): *Wildlife and Vegetation of Unmanaged Douglas-Fir Forests*, United States Department of Agriculture, Forest Service, General Technical Report PNW-GTR-285, pp.353–362.
- Bury, R.B., Fellers, G.M., and Ruth, S.B. 1969. First records of *Plethodon dunni* in California, and new distributional data on *Ascaphus truei*, *Rhyacotriton olympicus*, and *Hydromantes shastae*. *Journal of Herpetology* 3:157–161.
- Bury, R.B. and Germano, D.J. 2008. *Actinemys marmorata* (Baird and Girard 1852) western pond turtle, Pacific pond turtle, in Rhodin, A.G.J., Pritchard, P.C.H., van Dijk, P.P., Samure, R.A., Buhlmann, K.A. and Iverson, J.B. (Eds): *Conservation Biology of Freshwater Turtles and Tortoises*, Chelonian Monograph No. 5. Chelonian Research Foundation, Lunenburg, MA, pp.1.1–1.9.
- Bury, R.B. and Martin, M. 1967. The food of the salamander *Rhyacotriton olympicus*. *Copeia* 1967:487.
- Bury, R.B., Welsh, H.H., Jr., Germano, D.J., Ashton, D.T. (Eds). 2012. *Western Pond Turtle- Biology, Sampling Techniques, Inventory and Monitoring, Conservation, and Management: Northwest Fauna No. 7*, The Society for Northwestern Vertebrate Biology, Olympia, WA, 128pp.
- Calef, G.W. 1973. Natural mortality of tadpoles in a population of *Rana aurora*. *Ecology* 54:741–758.
- California Climate Action Team. 2006. *Climate Action Team Report to Governor Schwarzenegger and the Legislature*. California Environmental Protection Agency, Sacramento.
- Caltrans. 2012. *Highway Crossings for Herptiles (Reptiles and Amphibians)*. Preliminary Investigation requested by Margaret Gabil, Caltrans District 4, Division of Environmental Planning & Engineering.
- Camp, C.L. 1915. *Batrachoseps major* and *Bufo cognatus californicus*, new amphibia from southern California. *University of California Publications in Zoology* 12:327–334.
- Camp, C.L. 1916a. Description of *Bufo canorus*, a new toad from the Yosemite National Park. *University of California Publications in Zoology* 17:59–62.
- Camp, C.L. 1916b. Notes on the local distribution and habits of the amphibians and reptiles of southeastern California in the vicinity of the Turtle Mountains. *University of California Publications in Zoology* 12:503–544.
- Camp, C.L. 1917. Notes on the systematic status of the toads and frogs of California. *University of California Publications in Zoology* 17:115–125.
- Carothers, J.H. 1986. An experimental confirmation of morphological adaptation: toe fringes in the sand-dwelling lizard *Uma scoparia*. *Evolution* 40:871–874.
- Carpenter, C.C. 1963. Patterns of behavior in three forms of the fringe-toed lizards (*Uma*-Iguaniidae). *Copeia* 1963:406–412.
- Carroll, S., Ervin, E., and Fisher, R. 2005. Natural history notes: *Taricha torosa torosa*. Overwintering larvae. *Herpetological Review* 36:297.
- Cary, J.A. 2010. *Determining Habitat Characteristics That Predict Oviposition Site Selection for Pond-Breeding Northern Red-Legged Frogs (Rana aurora) in Humboldt County, California*. Humboldt State University, Arcata, CA.
- Cary, W. 1889. Biennial report of the fish commissioner of the state of Nevada. Appendix to the Journals of the Senate and Assembly 14th session, pp.3–7.
- Case, S.M. 1978. Biochemical systematics of members of the genus *Rana* native to western North America. *Systematic Zoology* 27:299–311.
- Case, T.J. and Fisher, R.N. 2001. Measuring and predicting species presence: coastal sage scrub case study, in Hunsaker, C.T., Goodchild, M.F., Friedl, M.A., and Case, T.J. (Eds): *Spatial Uncertainty in Ecology: Implications for Remote Sensing and GIS Applications*, Springer, New York, pp.47–71.
- Catenazzi, A. and Kupferberg, S.J. 2013. The importance of thermal conditions to recruitment success in stream-breeding frog populations distributed across a productivity gradient. *Biological Conservation* 168:40–48.
- Cayan, D.R., Luers, A., Franco, G., Hanemann, M., Croes, B., and Vine, E. 2008a. Overview of the California climate change scenarios project. *Climatic Change* 87:1–6.
- Cayan, D.R., Luers, A., Hanemann, M., Franco, G., and Croes, B. 2006. Possible Scenarios of Climate Change in California: Summary and Recommendations. White paper from California Climate Change Center.
- Cayan, D.R., Maurer, E.P., Dettinger, M.D., Tyree, M. and Hayhoe, K. 2008b. Climate change scenarios for the California region. *Climatic Change* 87 (Suppl. 1):S21–S42.
- CBD (Center for Biological Diversity). 2006. Petition to list the Amargosa River distinct

- population segment of the Mojave fringe-toed Lizard (*Uma scoparia*) as threatened or endangered under the U.S. Endangered Species Act. CDFG (California Department of Fish and Game). 2008. California Aquatic Invasive Species Management Plan, CDFW, Sacramento, CA.
- CDFW. 2013. Report: A Look Back at CDFW in 2012. CDFG, Sacramento, CA.
- Chan, L.M. and Zamudio, K.R. 2009. Population differentiation of temperate amphibians in unpredictable environments. *Molecular Ecology* 18:3185–3200.
- Chan-McLeod, A.C.A. 2003. Factors affecting the permeability of clearcuts to red-legged frogs. *The Journal of Wildlife Management* 67:663–671.
- Chan-McLeod, A.C.A. and Moy, A. 2007. Evaluating residual tree patches as stepping stones and short-term refugia for red-legged frogs. *Journal of Wildlife Management* 71:1836–1844.
- Chan-McLeod, A.C.A. and Wheeldon, B. 2004. Natural history notes: *Rana aurora*. Habitat and movement. *Herpetological Review* 35:375.
- Chang, A.T. and Shaffer, H.B. 2010. Population Genetics and Phylogenetics of *Rana cascadae* in the Lassen and Klamath Region. Unpublished Report to the United States Forest Service.
- Chelgren, N.D., Pearl, C.A., Adams, M.J., and Bowerman, J. 2008. Demography and movement in a relocated population of Oregon spotted frogs (*Rana pretiosa*): influence of season and gender. *Copeia* 2008:742–751.
- Chelgren, N.D., Rosenberg, D., Heppell, S., and Gitelman, A. (2006) Carryover aquatic effects on survival of metamorphic frogs during pond emigration. *Ecological Applications* 16:250–261.
- Chen, X., Barrows, C.W., and Li, B. 2006. Is the Coachella Valley fringe-toed lizard (*Uma inornata*) on the edge of extinction a Thousand Palms Preserve in California? *The Southwestern Naturalist* 51:28–34.
- Christin, M., Gendron, A.D., Brousseau, P., Ménard, L., Marcogliese, D.J., Cyr, D., Ruby, S., and Fournier, M. 2003. Effects of agricultural pesticides on the immune system of *Rana pipiens* and on its resistance to parasitic infection. *Environmental Toxicology and Chemistry* 22:1127–1133.
- Clark, R.W., Brown, W.S., Storch, R., and Zamudio, K.R. 2010. Roads, interrupted dispersal, and genetic diversity in timber rattlesnakes. *Conservation Biology* 24:1059–1069.
- Clarke, R.F. 1958. An ecological study of reptiles and amphibians in Osage County, Kansas. *Emporia State Research Studies* 7:1–52.
- Clarkson, R.W. and Rorabaugh, J.C. 1989. Status of leopard frogs (*Rana pipiens* complex, Ranidae) in Arizona and southeastern California. *The Southwestern Naturalist* 34:531–538.
- Claussen, D.L. 1973. The thermal relations of the tailed frog *Ascaphus truei* and the Pacific tree frog *Hyla regilla*. *Comparative Biochemistry and Physiology A* 44:137–153.
- Cliff, F.S. 1954. Snakes of the islands in the Gulf of California, Mexico. *Transactions of the San Diego Society of Natural History* 12:67–98.
- Coe, W.R. and Kunkel, B.W. 1906. Studies on the California limbless lizard, *Anniella*. *Transactions of the Connecticut Academy of Arts and Sciences* 12:349–403.
- Cole, C.J. 1962. Notes on the distribution and food habits of *Bufo alvarius* at the eastern edge of its range. *Herpetologica* 18:172–175.
- Coleman, J.L., Ford, N.B., and Herriman, K. 2008. A road survey of amphibians and reptiles in a bottomland hardwood forest. *Southeastern Naturalist* 7:339–348.
- Collins, J.P. and Lewis, M.A. 1979. Overwintering tadpoles and breeding season variation in the *Rana pipiens* complex in Arizona. *The Southwestern Naturalist* 24:371–373.
- Collins, J.T. and Taggart, T.W. 2009. Standard Common and Current Scientific Names for North American Amphibians, Turtles, Reptiles and Crocodylians. Center for North American Herpetology, Lawrence, KS.
- Cornrack, L., Bolster, B., Gustafson, J., Steele, D., and Burket, E. 2008. Species of special concern: a brief description of an important California Department of Fish and Game designation. California Department of Fish and Game, Wildlife Branch, Nongame Wildlife Program, Sacramento.
- Cooper, D.S. and Mathewson, P. 2008. Griffith Park Wildlife Management Plan. Report from Cooper Ecological Monitoring, Inc., Los Angeles, CA.
- Cooper, J.G. 1869. The naturalist in California. *The American Naturalist* 3:470–481.
- Cooper, J.G. 1870. The fauna of California and its geographical distribution. *Proceedings of the California Academy of Sciences* 1:118–123.
- Cope, E.D. 1869. Diagnosis of *Heloderma suspectum*. *Proceedings of the Academy of Natural Sciences of Philadelphia* 21:4–5.
- Cope, E.D. 1883. Notes on the geographical distribution of Batrachia and Reptilia in western North America. *Proceedings of the Academy of Natural Sciences of Philadelphia* 35:10–35.
- Cope, E.D. 1892. A critical review of the characters and variations of the snakes of North America.

- Proceedings of the United States National Museum 14:589-694.
- Cope, E.D. 1894. On the iguanian genus *Uma* Baird. The American Naturalist 28:434-435.
- Cope, E.D. 1895a. On some new North American snakes. The American Naturalist 29:676-680.
- Cope, E.D. 1895b. On the species of *Uma* and *Xantusia*. The American Naturalist 29:938-939.
- Corn, P.S. and Bury, R. 1989. Logging in western Oregon: responses of headwater habitats and stream amphibians. Forest Ecology and Management 29:39-57.
- Corn, P.S. and Fogleman, J.C. 1984. Extinction of montane populations of the northern leopard frog (*Rana pipiens*) in Colorado. Journal of Herpetology 18:147-152.
- Corn, P.S. and Livo, L.J. 1989. Leopard frog and wood frog reproduction in Colorado and Wyoming. Northwestern Naturalist 70:1-9.
- Cowles, R.B. and Bogert, C.M. 1944. A preliminary study of the thermal requirements of desert lizards. Bulletin American Museum Natural History 83:265-296.
- Crayon, J.J. 1988. Natural history notes: *Rana catesbiana*. Diet. Herpetological Review 29:232.
- Crother, B.I. 2009. Are standard names lists taxonomic straightjackets? Herpetologica 65:129-135.
- Cunningham, J.D. 1955. Arboreal habits of certain reptiles and amphibians in southern California. Herpetologica 11:217-220.
- Cunningham, J.D. 1956. Food habits of the San Diego alligator lizard. Herpetologica 12:225-230.
- Cunningham, J.D. 1959a. Notes on *Anniella*. Herpetologica 15:19-20.
- Cunningham, J.D. 1959b. Reproduction and food of some California snakes. Herpetologica 15:17-19.
- Cunningham, J.D. 1962. Observations on the natural history of the California toad, *Bufo californicus* Camp. Herpetologica 17:255-260.
- Cunningham, J.D. 1963. Additional observations on the ecology of the Yosemite toad, *Bufo canorus*. Herpetologica 19:56-61.
- Cunningham, J.D. 1966. Additional observations on the body temperatures of reptiles. Herpetologica 22:184-189.
- Cushman, K.A. and Pearl, C.A. 2007. A Conservation Assessment for the Oregon Spotted Frog (*Rana pretiosa*). Unpublished Report to USDA Forest Service Region 6. 46pp.
- Dalrymple, G.H., Steiner, T.M., Nodell, R.J., and Bernardino, F.S., Jr. 1991. Seasonal activity of the snakes of Long Pine Key, Everglades National Park. Copeia 1991:294-302.
- D'Amore, A., Hemingway, V., and Wasson, K. 2010. Do a threatened native amphibian and its invasive congener differ in response to human alteration of the landscape? Biological Invasions 12:145-154.
- D'Amore, A., Kirby, E., and McNicholas, M. 2009. Invasive species shifts ontogenetic resource partitioning and microhabitat use of a threatened native amphibian. Aquatic Conservation: Marine and Freshwater Ecosystems 19:534-541.
- Davidson, C. 2004. Declining downwind: amphibian population declines in California and historical pesticide use. Ecological Applications 14:1892-1902.
- Davidson, C. 2010. Natural history notes: *Rana draytonii*. Prey. Herpetological Review 41:66.
- Davidson, C., Benard, M., Shaffer, H., Parker, J., O'Leary, C., Conlon, J., and Rollins-Smith, L. 2007. Effects of chytrid and carbaryl exposure on survival, growth and skin peptide defenses in foothill yellow-legged frogs. Environmental Science and Technology 41:1771-1776.
- Davidson, C. and Fellers, G.M. 2005. *Bufo canorus*, in Lannoo, M.J. (Ed.): Amphibian Declines: The Conservation Status of United States Species, University of California Press, Berkeley, pp.400-401.
- Davidson, C., Shaffer, H.B., and Jennings, M.R. 2002. Spatial tests of the pesticide drift, habitat destruction, UV-B, and climate change hypotheses for California amphibian declines. Conservation Biology 16:1588-1601.
- Davidson, C., Stanley, K., and Simonich, S.M. 2012. Contaminant residues and declines of the Cascades frog (*Rana cascadae*) in the California Cascades, USA. Environmental Toxicology and Chemistry 31:1895-1902.
- Davis, M.A. and Heathcote, J. 2007. The price and quantity of residential land in the United States. Journal of Monetary Economics 54:2595-2620.
- De Lisle, H.F. 1979. Gila monster (*Heloderma suspectum*) found in California. Herpetology (Southwestern Herpetologists Society) 10:5-7.
- De Lisle, H.F. 1983. Banner year for California Gilas. Herpetology (Southwestern Herpetologists Society) 13:11.
- De Lisle, H., Canton, G., Feldner, J., O'Connor, P., Peterson, M., and Brown, P. 1986. The distribution and present status of the herpetofauna of the Santa Monica Mountains of Los Angeles and Ventura counties, California. Special Publication No. 2. Southwestern Herpetologists Society, Van Nuys, CA.
- de Queiroz, K. and Reeder, T.W. 2008. Squamata: Lizards. in Crother, B.I. (Ed.): Scientific and

- Standard English Names of Amphibians and Reptiles of North America North of Mexico, with Comments Regarding Confidence in Our Understanding. Society for the Study of Amphibians and Reptiles Herpetological Circular, Salt Lake City, UT. pp.24-45.
- De Vlaming, V.L. and Bury, R.B. 1970. Thermal selection in tadpoles of the tailed frog *Ascaphus truei*. *Journal of Herpetology* 4:179-189.
- Degenhardt, W.G., Painter, C.W., and Price, A.H. 1996. Amphibians and reptiles of New Mexico. University of New Mexico Press, Albuquerque, NM.
- Dejean, T., Valentini, A., Duparc, A., Pellier-Cuit, S., Pompanon, F., Taberlet, P., and Miaud, C. 2011. Persistence of environmental DNA in freshwater ecosystems. *PLOS ONE* 6:e23398.
- Denver, R.J. 1997a. Environmental stress as a developmental cue: corticotropin-releasing hormone is a proximate mediator of adaptive phenotypic plasticity in amphibian metamorphosis. *Hormones and Behavior* 31:169-179.
- Denver, R.J. 1997b. Proximate mechanisms of phenotypic plasticity in amphibian metamorphosis. *Integrative and Comparative Biology* 37:172.
- Denver, R.J., Mirhadi, N., and Phillips, M. 1998. Adaptive plasticity in amphibian metamorphosis: response of *Scaphiopus hammondi* tadpoles to habitat desiccation. *Ecology* 79:1859-1872.
- Dethlefsen, E.S. 1948. A subterranean nest of the Pacific giant salamander, *Dicamptodon ensatus* (Eschscholtz). *The Wasmann Collector* 7:81-84.
- Dever, J.A. 2007. Fine-scale genetic structure in the threatened foothill yellow-legged frog (*Rana boylei*). *Journal of Herpetology* 41:168-173.
- Dickinson, J.L., Zuckerberg, B., and Bontar, D.N. 2010. Citizen science as an ecological research tool: challenges and benefits. *Annual Review of Ecology, Evolution, and Systematics* 41:149-172.
- Dickman, M. 1968. The effect of grazing by tadpoles on the structure of a periphyton community. *Ecology* 49:1188-1190.
- Dill, W.A. 1944. The fishery of the lower Colorado River. *California Fish and Game*. 30:109-211.
- Diller, L.V. and Wallace, R.L. 1996. Distribution and habitat of *Rhyacotriton variegatus* in managed, young growth forests in north coastal California. *Journal of Herpetology* 30:184-191.
- Diller, L.V. and Wallace, R.L. 1999. Distribution and habitat of *Ascaphus truei* in streams on managed, young growth forests in north coastal California. *Journal of Herpetology* 33:71-79.
- Dimmitt, M.A. and Ruibal, R. 1980a. Environmental correlates of emergence in spadefoot toads (*Scaphiopus*). *Journal of Herpetology* 14:21-29.
- Dimmitt, M.A. and Ruibal, R. 1980b. Exploitation of food resources by spadefoot toads (*Scaphiopus*). *Copeia* 1980:854-862.
- Dixon, J.D. 1975. Geographic distribution: *Gerrhonotus panamintinus*. *Herpetological Review* 6:45.
- Dixon, J.R. 1967. Amphibians and Reptiles of Los Angeles County. Los Angeles Museum of Natural History, Los Angeles, CA.
- Dole, J.W. 1967. The role of substrate moisture and dew in the water economy of leopard frogs, *Rana pipiens*. *Copeia* 1967:141-149.
- Dole, J.W. 1971. Dispersal of recently metamorphosed leopard frogs, *Rana pipiens*. *Copeia* 1971:221-228.
- Doubledee, R.A., Mueller, E.B., and Nisbet, R.M. 2003. Bullfrogs, disturbance regimes, and the persistence of California red-legged frogs. *Journal of Wildlife Management* 67:424-438.
- Douglas, M., Douglas, M., Schuett, G., Beck, D., and Sullivan, B. 2010. Conservation phylogenetics of helodermatid lizards using multiple molecular markers and a supertree approach. *Molecular Phylogenetics and Evolution* 55:153-167.
- Drennan, J.E., Jackman, R.E., Marlow, D.R., and Wiseman, K.D. 2006. Identifying climatic and water flow triggers associated with breeding activities of a foothill yellow-legged frog (*Rana boylei*) population on the North Fork Feather River, California. Garcia and Associates, San Francisco, CA.
- Drost, C.A. and Fellers, G.M. 1996. Collapse of a regional frog fauna in the Yosemite area of the California Sierra Nevada, U.S.A. *Conservation Biology* 10:414-425.
- Dugan, E.A., Figueroa, A., and Hayes, W.K. 2008. Home range size, movements, and mating phenology of sympatric red diamond (*Crotalus ruber*) and southern Pacific (*C. oreganus helleri*) rattlesnakes in southern California, in Hayes, W.K., Beaman, K.R., Cardwell, M.D., and Bush, S.P. (Eds): *The Biology of Rattlesnakes Symposium*, Loma Linda University Press, Loma Linda, CA, pp.353-364.
- Dugan, E.A. and Hayes, W.K. 2012. Diet and feeding ecology of the red diamond rattlesnake, *Crotalus ruber* (Serpentes: Viperidae). *Herpetologica* 2:203-217.
- Dumas, P.C. 1966. Studies of the *Rana* species complex in the Pacific northwest. *Copeia* 1966:60-74.
- Dundee, H.A. and Miller, M.C. 1968. Aggregative behavior and habitat conditioning by the prairie ringneck snake, *Diadophis punctatus arnyi*. *Tulane Studies in Zoology and Botany* 15:41-58.

- Dunlap, D.G. 1955. Inter- and intraspecific variation in Oregon frogs of the genus *Rana*. *The American Midland Naturalist* 54:314-331.
- Dupuis, L. and Steventon, D. 1999. Riparian management and the tailed frog in northern coastal forests. *Forest Ecology and Management* 124:35-43.
- Dupuis, L.A., Bunnell, F.L., and Friele, P.A. 2000. Determinants of the tailed frog's range in British Columbia, Canada. *Northwest Science* 74:109-115.
- Elliott, I., Gerhardt, C., and Davidson, C. 2009. *The Frogs and Toads of North America: A Comprehensive Guide to Their Identification, Behavior, and Calls*, Houghton Mifflin Harcourt, New York.
- Ely, E. 1992. Geographic distribution: *Thamnophis hammondi*. *Herpetological Review* 23:124-125.
- Emery, A.R., Beist, A.H., and Kodaira, K. 1972. Under-ice observations of wintering sites of leopard frogs. *Copeia* 1972:123-126.
- Emmerich, K. and Cunningham, L. 2003. Geographic distribution: *Diadophis punctatus*. *Herpetological Review* 34:169.
- Ernst, C.H. and Ernst, E.M. 2003. *Snakes of the United States and Canada*, Smithsonian Books, Washington, DC.
- Ernst, C.H. and Lovich, J.E. 2009. *Turtles of the United States and Canada*. Johns Hopkins University Press, Baltimore, MD.
- Ervin, E.L., Anderson, A.E., Cass, T.L., and Murcia, R.E. 2001. Natural history notes: *Spea hammondi*. Elevation record. *Herpetological Review* 32:36.
- Ervin, E.L., Beaman, K.R., and Fisher, R.N. 2013. Correction of locality records of the endangered arroyo toad (*Anaxyrus californicus*) in the desert regions of Southern California. *Bulletin of the Southern California Academy of Sciences* 112:197-205.
- Ervin, E.L. and Burkhardt, T.R. 2006. Natural history notes: *Ambystoma tigrinum mavortium*. Extralimital populations. *Herpetological Review* 37:435.
- Ervin, E.L. and Cass, T.L. 2007. Natural history notes: *Spea hammondi*. Reproductive pattern. *Herpetological Review* 38:196-197.
- Ervin, E.L. and Fisher, R.N. 2001. Natural history notes: *Thamnophis hammondi*. Prey. *Herpetological Review* 32:265-266.
- Ervin, E.L. and Fisher, R.N. 2007. Natural history notes: *Thamnophis hammondi*. Foraging behavior. *Herpetological Review* 38:345-346.
- Ervin, E.L., Kisner, D.A., and Fisher, R.N. 2006. Natural history notes: *Bufo californicus*. Mortality. *Herpetological Review* 37:199.
- Ervin, E.L., Smith, C.D., and Christopher, S.V. 2005. Natural history notes: *Spea hammondi*. Reproduction. *Herpetological Review* 36:309-310.
- Eschscholtz, J.F.V. 1833. *Zoologischer Atlas, enthaltend Abbildungen und Beschreibungen neuer Thierarten, während des Flottcapitains von Kotzebue zweiter Reise um die Welt, auf Russisch-Kaiserlich Kriegsschupp Predpriaetië in den Jahren 1823-1826 herausgegeben von D. Martin Heinrich Rathke. Fünftes Heft. G. Reimer, Berlin.* (In German)
- Ewert, M.A. 1991. Cold torpor, diapause, delayed hatching and aestivation in reptiles and birds, in Deeming, D.C. and Ferguson, M.W.J. (Eds): *Egg Incubation: Its Effects on Embryonic Development in Birds and Reptiles*, Cambridge University Press, Cambridge, UK, pp.173-191.
- Fahrig, L., Pedlar, J.H., Popc, S.E., Taylor, P.D., and Wegner, J.F. 1995. Effect of road traffic on amphibian density. *Biological Conservation* 73:177-182.
- Feaver, P.E. 1971. *Breeding Pool Selection and Larval Mortality of Three California Amphibians: Ambystoma tigrinum californiense Gray Hyla regilla Baird and Girard and Scaphiopus hammondi hammondi Girard*. Master's Thesis. Fresno State College, Fresno, CA.
- Feldman, C.R. and Parham, J.F. 2002. Molecular phylogenetics of emydid turtles: taxonomic revision and the evolution of shell kinesis. *Molecular Phylogenetics and Evolution* 22:388-398.
- Feldman, C.R. and Spicer, G.S. 2006. Comparative phylogeography of woodland reptiles in California: repeated patterns of cladogenesis and population expansion. *Molecular Ecology* 15:2201-2222.
- Fellers, G.M. 2005a. *Rana boylei*, in Lannoo, M.J. (Ed.): *Amphibian Declines: The Conservation Status of United States Species*, University of California Press, Berkeley, pp.534-536.
- Fellers, G.M. 2005b. *Rana draytonii*, in Lannoo, M.J. (Ed.): *Amphibian Declines: The Conservation Status of United States Species*. University of California Press, Berkeley, pp.552-554.
- Fellers, G.M. and Drost, C.A. 1993. Disappearance of the cascades frog *Rana cascadae* at the southern end of its range, California, U.S.A. *Biological Conservation* 65:177-181.
- Fellers, G.M. and Kleeman, P.M. 2007. California red-legged frog (*Rana draytonii*) movement and habitat use: implications for conservation. *Journal of Herpetology* 41:276-286.

- Fellers, G.M., Lanner, A.E., Rathbun, G., Bobzien, S., Alvarez, J., Sterner, D., Seymour, R.B., and Westphal, M. 2001. Overwintering tadpoles in the California red-legged frog (*Rana aurora draytonii*). *Herpetological Review* 32:156-157.
- Fellers, G.M., Pope, K.L., Stead, J.E., Koo, M.S., and Welsh, H.H., Jr. 2008. Turning population trend monitoring into active conservation: can we save the Cascades frog (*Rana cascadae*) in the Lassen region of California? *Herpetological Conservation and Biology* 3:28-39.
- Fellers, G.M., Wood, L.L., Carlisle, S., and Pratt, D. 2010. Unusual subterranean aggregations of the California giant salamander, *Dicamptodon ensatus*. *Herpetological Conservation and Biology* 5:149-154.
- Ferguson, D.E. 1961. The geographic variation of *Ambystoma macrodactylum* Baird, with the description of two new subspecies. *The American Midland Naturalist* 65:311-338.
- Ficetola, G.F., Miaud, C., Pompanon, F., and Taberlet, P. 2008. Species detection using environmental DNA from water samples. *Biology Letters* 4:423-425.
- Fisher, R.N. 2004. Life on the "Island": animals, in Houk, R. (Ed.): *Understanding the Life of Point Loma*, Cabrillo National Monument Foundation, San Diego, CA, pp.112-131.
- Fisher, R.N. and Case, T.J. 2000. Distribution of the herpetofauna of coastal Southern California with reference to elevation effects. In Keeley, J., Baer-Keeley, M., and Fotheringham, C.J. (Eds): *2nd Interface Between Ecology and Land Development in California, United States Geological Survey Open-File Report 00-62*, pp.137-143.
- Fisher, R.N. and Shaffer, H.B. 1996. The decline of amphibians in California's Great Central Valley. *Conservation Biology* 10:1387-1397.
- Fisher, R.N., Suarez, A.V., and Case, T.J. 2002. Spatial patterns in the abundance of the coastal horned lizard. *Conservation Biology* 16:205-215.
- Fitch, H.S. 1936. Amphibians and reptiles of the Rogue River Basin, Oregon. *The American Midland Naturalist* 17:634-652.
- Fitch, H.S. 1938. *Rana boylei* in Oregon. *Copeia* 1938:148-148.
- Fitch, H.S. 1940. A biogeographical study of the *ordinoides* Artenkreis of garter snakes (genus *Thamnophis*). *University of California Publications in Zoology* 44:1-150.
- Fitch, H.S. 1941. The feeding habits of California garter snakes. *California Fish and Game* 27:2-32.
- Fitch, H.S. 1948. Further remarks concerning *Thamnophis ordinoides* and its relatives. *Copeia* 1948:121-126.
- Fitch, H.S. 1970. Reproductive cycles in lizards and snakes. *University of Kansas Museum of Natural History Miscellaneous Publications* 52:1-247.
- Fitch, H.S. 1975. A demographic study of the ringneck snake (*Diadophis punctatus*) in Kansas. *University of Kansas Museum of Natural History Miscellaneous Publications* 62:1-53.
- Fite, K.V., Blaustein, A., Bengston, L., and Hewitt, H.E. 1998. Evidence of retinal light damage in *Rana cascadae*: a declining amphibian species. *Copeia* 1998:906-914.
- Fontanella, F.M., Feldman, C.R., Siddall, M.E., and Burbrink, F.T. 2008. Phylogeography of *Diadophis punctatus*: extensive lineage diversity and repeated patterns of historical demography in a trans-continental snake. *Molecular Phylogenetics and Evolution* 46:1049-1070.
- Ford, R.S. 1981. Geographic distribution: *Heterodermia suspectum cinctum*. *Herpetological Review* 12:64.
- Ford, R.S. 1983. Reptiles and Amphibians, in Stone, R.D. and Sumida, V.A. (Eds): *The Kingston Range of California: A Resource Survey*. Vol. 10. Environmental Field Program, University of California, Santa Cruz, pp.123-130.
- Foreman, L.D. (Ed.). 1997. Flat-Tailed Horned Lizard Rangeland Management Strategy. Report of Interagency Working Group.
- Forsman, E.D. and Swingle, J.K. 2007. Use of arboreal nests of tree voles (*Arborimus spp.*) by amphibians. *Herpetological Conservation and Biology* 2:113-118.
- Forson, D. and Storfer, A. 2006. Effects of atrazine and iridovirus infection on survival and life-history traits of the long-toed salamander (*Ambystoma macrodactylum*). *Environmental Toxicology and Chemistry* 25:168-173.
- Fouquette, M.J., Jr. 1968. Remarks on the type specimen of *Bufo alvarius* Girard. *The Great Basin Naturalist* 28:70-72.
- Fouquette, M.J., Jr., Painter, C.W., and Nanjappa, P., Jr. 2005. *Bufo alvarius* Girard, 1859: Colorado River toad, in Lannoo, M.J. (Ed.): *Amphibian Declines: The Conservation Status of United States Species*. University of California Press, Berkeley, pp.384-386.
- Fox, W. 1951. Relationships among the garter snakes of the *Thamnophis elegans* rassenkreis. *University of California Publications in Zoology* 50:485-530.
- Fox, W. 1956. Seminal receptacles of snakes. *The Anatomical Record* 124:519-539.

- Franco, G., Cayan, D.R., Moser, S., Hancmann, M., and Jones, M. 2011. Second California assessment: integrated climate change impacts assessment of natural and managed systems. *Climatic Change* 2011: S1–S19.
- Fried, J.S., Torn, M.S., and Mills, E. 2004. The impact of climate change on wildfire severity: a regional forecast for northern California. *Climatic Change* 64:169–191.
- Fromer, P.S., Doder, M., and Patterson, C. 1983. A Population Study of the Mojave Fringe-Toed Lizard (*Uma scoparia*) on the Twentynine Palms MCAGGC. Recon Number R 1397. Natural Resources Office, Marine Corps Air Ground Combat Center, Twentynine Palms, CA.
- Frost, D.R., Grant, T., Faiyovich, J., Bain, R., Haas, A., Iñárrada, C., De Sá, R. et al. 2006a. The amphibian tree of life. *Bulletin of the American Museum of Natural History* 297:1–291.
- Frost, D.R., Grant, T., and Mendelson, J.R. 2006b. *Ollotis* Cope, 1875 is the oldest name for the genus currently referred to as *Craugastis* Cope, 1875 (Anura: Hyloidea: Bufonidae). *Copeia* 2006:558–558.
- Frost, D.R. and Hillis, D.M. 1990. Species in concept and practice: herpetological applications. *Herpetologica* 46:86–104.
- Frost, D.R., McDiarmid, R.W., and Mendelson, J.R. 2009a. Response to the point of view of Gregory B. Pauly, David M. Hillis, and David C. Cannatella, by the Anuran Subcommittee of the SSAR/HI/ASIH Scientific and Standard English Names List. *Herpetologica* 65:136–153.
- Frost, D.R., Mendelson, J.R., and Pramuk, J. 2009b. Further notes on the nomenclature of Middle American toads (Bufonidae). *Copeia* 2009:418–418.
- Frost, J.S. and Platz, J.E. 1983. Comparative assessment of modes of reproductive isolation among four species of leopard frogs (*Rana pipiens* complex). *Evolution* 37:66–78.
- FTHL ICC (Flat-Tailed Horned Lizard Interagency Coordinating Committee). 2003. Flat-Tailed Horned Lizard Rangewide Management Strategy, 2003 Revision: An Arizona-California Conservation Strategy.
- Fuhrman, F.A. 1967. Tetrodotoxin. *Scientific American*. 217:60–71.
- Fuller, D.D. and Lind, A.J. 1992. Implications of fish habitat improvement structures for other stream vertebrates, in Harris, R. and Erman, D. (Eds): *Proceedings of the Symposium on Biodiversity of Northwestern California*, Santa Rosa, pp.96–104.
- Fuller, T.E. 2008. The Spatial Ecology of the Exotic Bullfrog (*Rana catesbeiana*) and Its Relationship to the Distribution of the Native Herpetofauna in a Managed River System. Master's Thesis. Humboldt State University, Arcata, CA.
- Fuller, T.E., Pope, K.L., Ashton, D.T., and Welsh, H.H., Jr. 2010. Linking the distribution of an invasive amphibian (*Rana catesbeiana*) to habitat conditions in a managed river system in northern California. *Restoration Ecology* 19:204–213.
- Funk, R.S. 1966. Notes about *Heloderma suspectum* along the western extremity of its range. *Herpetologica* 22:254–258.
- Funk, R.S. 1974. Geographic distribution: *Kinosternon sonoriense*. *Herpetological Review* 5:20.
- Funk, W.C. and Dunlap, W.W. 1999. Colonization of high-elevation lakes by long-toed salamanders (*Ambystoma macrodactylum*) after the extinction of introduced trout populations. *Canadian Journal of Zoology* 77:1759–1767.
- Funk, W.C., Pearl, C.A., Draheim, H.M., Adams, M.J., Mullins, T.D., and Haig, S.M. 2008. Range-wide phylogeographic analysis of the spotted frog complex (*Rana luteiventris* and *Rana pretiosa*) in northwestern North America. *Molecular Phylogenetics and Evolution* 49:198–210.
- Gamradt, S. and Kats, L. 1996. Effect of introduced crayfish and mosquitofish on California newts. *Conservation Biology* 10:1155–1162.
- Gamradt, S. and Kats, L. 1997. Impact of chaparral wildfire-induced sedimentation on oviposition of stream-breeding California newts (*Taricha torosa*). *Oecologia* 110:546–549.
- Gamradt, S., Kats, L., and Anzalone, C. 1997. Aggression by non-native crayfish deters breeding in California newts. *Conservation Biology* 11:793–796.
- Garcia and Associates. 2005. Results of 2004 Surveys and Monitoring for Foothill Yellow-Legged Frogs (*Rana boylei*) within the Rock Creek-Cresta Project Area, North Fork Feather River and 2002-2004 Recreation and Pulse Flow Biological Evaluation Summary. Prepared for Pacific Gas and Electric Company, San Ramon, California, Job 332/80, 89pp.
- Garcia, T.S., Romansic, J.M., and Blaustein, A.R. 2006. Survival of three species of anuran metamorphs exposed to UV-B radiation and the pathogenic fungus *Batrachochytrium dendrobatidis*. *Diseases of Aquatic Organisms*. 72:163.
- Garcia-Paris, M., Buchholz, D.R., and Parra-Olea, G. 2003. Phylogenetic relationships of Pelobatoida re-examined using mtDNA. *Molecular Phylogenetics and Evolution*. 28:12–23.

- Gardner, T.J., Foley, D.H., Brodie, E.D., and Young, K.V. 2004. Barrier fences prevent road mortalities in the flat-tailed horned lizard (*Phrynosoma mcallii*). *Herpetological Review* 35:250–250.
- Garman, S. 1884. The reptiles and batrachians of North America. *Memoirs of the Museum of Comparative Zoology* 8:1–185.
- Garwood, J.M. 2009. Spatial Ecology of the Cascades Frog: Identifying Dispersal, Migration, and Resource Uses at Multiple Spatial Scales. Master's Thesis. Humboldt State University, Arcata, CA.
- Garwood, J.M. and Welsh, H.H., Jr. 2007. Ecology of the Cascades Frog (*Rana cascadae*) and Interactions with Garter Snakes and Nonnative Trout in the Trinity Alps Wilderness, California. Final Report to California Department of Fish and Game and National Fish and Wildlife Foundation. 75pp.
- Gehlbach, F.R. 1965. Herpetology of the Zuni Mountains region, northwestern New Mexico. *Proceedings of the US National History Museum* 116:243–332.
- Gehlbach, F.R. 1974. Evolutionary relations of southwestern ringneck snakes (*Diadophis punctatus*). *Herpetologica* 30:140–148.
- Gergus, E.W.A. 1998. Systematics of the *Bufo microscaphus* complex: allozyme evidence. *Herpetologica* 54:317–325.
- Gergus, E.W.A., Grismer, L.L., and Beaman, K.R. 1997. Geographic distribution: *Bufo californicus*. *Herpetological Review* 28:47.
- Germano, D.J. 2010. Ecology of western pond turtles (*Actinemys marmorata*) at sewage-treatment facilities in the San Joaquin Valley, California. *The Southwestern Naturalist* 55:89–97.
- Germano, D.J. and Bury, R.B. 2001. Western pond turtles (*Clemmys marmorata*) in the Central Valley of California: status and population structure. *Transactions-Western Section of the Wildlife Society* 37:22–36.
- Germano, D.J. and Bury, R.B. 2009. Variation in body size, growth, and population structure of *Actinemys marmorata* from lentic and lotic habitats in southern Oregon. *Journal of Herpetology* 43:510–520.
- Germano, D.J. and Rathbun, G.B. 2008. Growth, population structure, and reproduction of western pond turtles (*Actinemys marmorata*) on the central coast of California. *Chelonian Conservation and Biology* 7:188–194.
- Germano, D.J., Rathbun, G.B., and Saslaw, L.R. 2001. Managing exotic grasses and conserving declining species. *Wildlife Society Bulletin* 29:551–559.
- Germano, J.M. and Bishop, P.J. 2009. Suitability of amphibians and reptiles for translocation. *Conservation Biology* 23:7–15.
- Gerson, M. 2011. Population status and habitat affinities of the Blainville's horned lizard (*Phrynosoma blainvillii*) at a site in the northern San Joaquin Valley, California, USA. *Herpetological Conservation and Biology* 6(2):228–236.
- Gibbs, J.P. and Shriver, W.G. 2002. Estimating the effects of road mortality on turtle populations. *Conservation Biology* 16:1647–1652.
- Gibbs, J.P. and Shriver, W.G. 2005. Can road mortality limit populations of pool-breeding amphibians? *Wetlands Ecology and Management* 13: 281–289.
- Gilbertson, M.K., Haffner, D.G., Drouillard, K.G., Albert, A., and Dixon, B. 2003. Immunosuppression in the northern leopard frog (*Rana pipiens*) induced by pesticide exposure. *Environmental Toxicology and Chemistry* 22:101–110.
- Gilhen, J. 1970. An unusual Nova Scotian population of the northern ringneck snake, *Diadophis punctatus edwardsi* (Merrem). *Occasional Papers of the Nova Scotia Museum, Halifax* 9:1–13.
- Girard C. 1859. in Baird. Reptiles of the Boundary, with Notes by the Naturalists of the Survey. Department of the Interior, Washington, DC.
- Giuliani, D. 1996. Resurvey of Eastern Sierra Nevada Salamanders. Unpublished Report. California Department of Fish and Game, Sacramento.
- Goebel, A.M., Ranker, T.A., Corn, P.S., and Olmstead, R.G. 2009. Mitochondrial DNA evolution in the *Anaxyrus boreas* species group. *Molecular Phylogenetics and Evolution* 50:209–225.
- Goldberg, C.S. and Waits, L.P. 2010. Comparative landscape genetics of two pond-breeding amphibian species in a highly modified agricultural landscape. *Molecular Ecology* 19:3650–3663.
- Goldberg, S.R. 1972. Reproduction in the southern alligator lizard *Gerrhonotus multicarinatus*. *Herpetologica* 28:267–273.
- Goldberg, S.R. 1983. Reproduction of the coast horned lizard, *Phrynosoma coronatum*, in southern California. *The Southwestern Naturalist* 28:478–479.
- Goldberg, S.R. 1999. Reproduction in the red diamond rattlesnake in California. *California Fish and Game* 85:177–180.
- Goldberg, S.R. 2000. Reproduction in the glossy snake, *Arizona elegans* (Serpentes: Colubridae) from California. *Bulletin of the Southern California Academy of Sciences* 99:105–109.

- Goldberg, S.R. and Beaman, K.R. 2003. Natural history notes: *Elgaria panamintina*. *Reproduction*. *Herpetological Review* 34: 143.
- Goldberg, S.R. and Müller, C. 1985. Reproduction of the silvery legless lizard, *Anniella pulchra pulchra* (Anniellidae), in Southern California. *The Southwestern Naturalist* 30: 617–619.
- Goldberg, S.R., Mahrtdt, C.R., and Beaman, K.R. 2010. Reproduction in Cope's leopard lizard, *Gambelia copeii* (Squamata: Grotaphytidae). *Bulletin of the Southern California Academy of Sciences* 109: 15–17.
- Gomez, D.M. and Anthony, R.G. 1996. Amphibian and reptile abundance in riparian and upslope areas of five forest types in western Oregon. *Northwest Science* 70: 109–119.
- Gonsolin, T.E. 2010. Ecology of Foothill Yellow-Legged Frogs in Upper Coyote Creek, Santa Clara County, CA. Master's Thesis. San Jose State University, San Jose, CA.
- Good, D.A. 1988. Allozyme variation and phylogenetic relationships among the species of *Elgaria* (Squamata: Anguillidae). *Herpetologica* 44: 154–162.
- Good, D.A. 1989. Hybridization and cryptic species in *Dicamptodon* (Caudata: Dicamptodontidae). *Evolution* 43: 728–744.
- Good, D.A. and Wake, D.B. 1992. Geographic variation and speciation in the torrent salamanders of the genus *Rhyacotriton* (Caudata: Rhyacotritonidae). *University of California Publications in Zoology* 126: 1–91.
- Goodman, R.H., Jr. 1997. Occurrence of double clutching in the southwestern pond turtle, *Clemmys marmorata pallida*, in the Los Angeles Basin. *Chelonian Conservation and Biology* 2: 419–420.
- Goodman, R.H., Jr., Watanabe, S.K., Condon, K.P., Pires, M.P., and Benton, M.S. 1998. Geographic distribution: *Batrachoseps gabrieli*. *Herpetological Review* 29: 171.
- Gorman, J. 1957. Recent collections of the California limbless lizard, *Anniella pulchra*. *Copeia* 1957: 148–150.
- Graeter, G.J., Buhlmann, K.A., Wilkinson, L.R., and Gibbons, J.W. (Eds). 2013. *Inventory and Monitoring: Recommended Techniques for Reptiles and Amphibians, with Application to the United States and Canada*. Paratners in Amphibian and Reptile Conservation, Birmingham, AL.
- Grant, D., Anderson, O., and Twitty, V. 1968. Homing orientation by olfaction in newts (*Taricha rivularis*). *Science* 160: 1354–1356.
- Grant, T.J. and Doherty, P.F. 2006. Natural history notes: *Phrynosoma mcallii*. Hibernation. *Herpetological Review* 37: 346–347.
- Grant, T.J. and Doherty, P.F. 2007. Monitoring of the flat-tailed horned lizard with methods incorporating detection probability. *Journal of Wildlife Management* 71: 1050–1056.
- Grant, T.J. and Doherty, P.F. 2009. Potential mortality effects of off-highway vehicles on the flat-tailed horned lizard (*Phrynosoma mcallii*): a manipulative experiment. *Environmental Management* 43: 508–513.
- Grasso, R.L. 2005. Palatability and Antipredator Response of Yosemite Toad (*Bufo canorus*) to Nonnative Brook Trout (*Salvelinus fontinalis*) in the Sierra Nevada Mountains of California. Master's Thesis. California State University, Sacramento.
- Grasso, R.L., Coleman, R.M., and Davidson, C. 2010. Palatability and antipredator response of Yosemite toads (*Anaxyrus canorus*) to nonnative brook trout (*Salvelinus fontinalis*) in the Sierra Nevada Mountains of California. *Copeia* 2010: 457–462.
- Gray, J.E. 1839. *Phrynosoma blainvillii*, in *The Zoology of Captain Beechey's voyage: Compiled from the collections and notes made by Captain Beechey, the officers and naturalist of the expedition, during a voyage to the Pacific and Behring's Straits performed in his majesty's ship Blossom, under the command of Captain FW Beechey, in the years 1825, 26, 27 and 28*, HIC Bohn, London, p.96.
- Gray, J.E. 1852. Descriptions of several new genera of reptiles, principally from the collection of HMS Herald. *Annual Magazine of Natural History* 10: 437–440.
- Graybeal, A. 1993. The phylogenetic utility of cytochrome b: lessons from bufonid frogs. *Molecular Phylogenetics and Evolution* 2: 256–269.
- Green, D.E. and Kagarise Sherman, C. 2001. Diagnostic histological findings in Yosemite toads (*Bufo canorus*) from a die-off in the 1970s. *Journal of Herpetology* 35: 92–103.
- Green, D.M. 1985. Differentiation in amount of centromeric hetero-chromatin between subspecies of the red-legged frog, *Rana aurora*. *Copeia*, 1985: 1071–1074.
- Green, D.M. 1986a. Systematics and evolution of western North American frogs allied to *Rana aurora* and *Rana boylei*: electrophoretic evidence. *Systematic Zoology* 35: 283–296.
- Green, D.M. 1986b. Systematics and evolution of western North American frogs allied to *Rana aurora* and *Rana boylei*: karyological evidence. *Systematic Zoology* 35: 273–282.
- Green, D.M., Kaiser, H., Sharbel, T.F., Kearsley, J., and McAllister, K.R. 1997. Cryptic species of

- spotted frogs, *Rana pretiosa* complex, in western North America. *Copeia* 1997:1–8.
- Green, D.M., Sharbel, T.F., Kearsley, J., and Kaiser, H. 1996. Postglacial range fluctuation, genetic subdivision and speciation in the western North American spotted frog complex, *Rana pretiosa*. *Evolution* 50:374–390.
- Greenberg, D.B. 2002. The Ecology of Movement and Site Selection in Desert Rattlesnakes (*Crotalus Mitchelli* and *Crotalus Ruber*) of the Southwestern United States. PhD Dissertation. University of California, Santa Barbara.
- Greenberg, D.B. and McClintock, W.J. 2008. Remember the third dimension: terrain modeling improves estimates of snake home range size. *Copeia* 2008:801–806.
- Gregory, P.T. 1977. Life-history parameters of the red-sided garter snake (*Thamnophis sirtalis parietalis*) in an extreme environment, the Interlake region of Manitoba. *National Museum of Canada Publications in Zoology* 13:1–44.
- Gregory, P.T. and Larsen, K.W. 1993. Geographic variation in reproductive characteristics among Canadian populations of the common garter snake (*Thamnophis sirtalis*). *Copeia* 1993:946–958.
- Griffin, P.C. and Case, T.J. 2001. Terrestrial habitat preferences of adult arroyo southwestern toads. *Journal of Wildlife Management* 65:633–644.
- Grinnell, J. 1908. The Biota of the San Bernardino Mountains. University of California Press, Berkeley.
- Grinnell, J. and Camp, C.L. 1917. A distributional list of the amphibians and reptiles of California. University of California Publications in Zoology 17:127–208.
- Grinnell, J., and Grinnell, H.W. 1907. Reptiles of Los Angeles County, California. *Throop Institute Bulletin* 35:1–64.
- Grinnell, J. and Storer, T.I. 1924. Animal life in the Yosemite. University of California Press, Berkeley.
- Grismer, L.L. 1994. The origin and evolution of the peninsular herpetofauna of Baja California, México. *Herpetological Natural History* 2:51–106.
- Grismer, L.L. 2002. Amphibians and reptiles of Baja California, including its Pacific islands, and the islands in the Sea of Cortés. University of California Press, Berkeley.
- Grismer, L.L. and Galvan, M.A. 1986. A new night lizard (*Xantusia henshawi*) from a sandstone habitat in San Diego County, California. *Transactions of the San Diego Society of Natural History* 21:155–165.
- Graff, L.A. 2011. A Species Distribution Model for Guiding Oregon Spotted Frog (*Rana pretiosa*) Surveys Near the Southern Extent of Its Geographic Range. Master's Thesis. Humboldt State University, Arcata, CA.
- Hager, S.B. 1996. Natural history notes: *Phrynosoma coronatum blainvillii*. *Growth. Herpetological Review* 27:80.
- Hager, S.B. and Brattstrom, B.H. 1997. Surface activity of the San Diego horned lizard *Phrynosoma coronatum blainvillii*. *The Southwestern Naturalist* 42:339–344.
- Haggard, J.A.G. 2000. A Radio Telemetric Study of the Movement Patterns of Adult Northern Red-Legged Frogs (*Rana aurora aurora*) at Freshwater Lagoon. Humboldt County, California. Humboldt State University, Arcata, CA.
- Haggarty, M. 2006. Habitat Differentiation and Resource Use among Different Age Classes of Post Metamorphic *Rana boylei* on Red Bank Creek, Tehama County, California. Master's Thesis. Humboldt State University, Arcata, CA.
- Halama, K.J., Malisch, A.J., Aspell, M., Rotenberry, J.T., and Allen, M.F. 2008. Modeling the landscape niche characteristics of red diamond rattlesnakes (*Crotalus ruber*): implications for biology and conservation, in Hayes, W.K., Beaman, K.R., Cardwell, M.D., and Bush, S.P. (Eds): *The Biology of Rattlesnakes Symposium*, Loma Linda University Press, Loma Linda, CA, pp.463–472.
- Hall, D.H. and Steidl, R.J. 2007. Movements, activity, and spacing of Sonoran mud turtles (*Kinosternon sonoriense*) in interrupted mountain streams. *Copeia* 2007:403–412.
- Hallowell, E. 1852. Descriptions of new species of reptiles inhabiting North America. *Proceedings of the Academy of Natural Sciences of Philadelphia* 6:177–182.
- Hammerson, G.A. 1977. Head body temperature differences monitored by telemetry in the snake *Masticophis flagellum piceus*. *Comparative Biochemistry and Physiology* 57:399–402.
- Hancock, J.P. 2009. Arroyo Toad (*Anaxyrus californicus*) Life History, Population Status, Population Threats, and Habitat Assessment of Conditions at Fort Hunter Liggett, Monterey County, California. Master's Thesis. California Polytechnic State University, San Luis Obispo.
- Hansen, R.W. 1988. Kern Canyon Slender Salamander (*Batrachoseps simatus*). California Department of Transportation, Biological Survey Report, District 6, Fresno.
- Hansen, R.W. and Wake, D.B. 2005a. *Batrachoseps campi*, in Lannoo, M.J. (Ed.): *Amphibian*

- Declines: The Conservation Status of United States Species, University of California Press, Berkeley, pp.669–671.
- Hansen, R.W. and Wake, D.B. 2005b. *Batrachoseps minor*, in Lannoo, M.J. (Ed.): Amphibian Declines: The Conservation Status of United States Species, University of California Press, Berkeley, pp.682–683.
- Hansen, R.W. and Wake, D.B. 2005c. *Batrachoseps relictus*, in Lannoo, M.J. (Ed.): Amphibian Declines: The Conservation Status of United States Species, University of California Press, Berkeley, pp.688–690.
- Hansen, R.W., Goodman, R.H. and Wake, D.B. 2005d. *Batrachoseps gabrieli*, in M.J. Lannoo, (Ed.): Amphibian Declines: The Conservation Status of United States Species, University of California Press, Berkeley, pp.672–673.
- Hanson, K., Snyder, J., and Kats, L. 1994. Natural history notes: *Taricha torosa*. Diet. Herpetological Review 25:62.
- Harding, J.H. 1997. Amphibians and Reptiles of the Great Lakes Region. University of Michigan Press, Ann Arbor.
- Harte, J., Ostling, A., Green, J.L., and Kinzig, A. 2004. Climate change and extinction risk. Nature 430:33.
- Hatch, A.C. and Blaustein, A.R. 2000. Combined effects of UV-b, nitrate, and low pH reduce the survival and activity level of larval cascades frogs (*Rana cascadae*). Archives of Environmental Contamination and Toxicology 39:494–499.
- Hawkins, C.P., Gottschalk, L.J., and Brown, S.S. 1988. Densities and habitat of tailed frog tadpoles in small streams near Mount St. Helens Washington U.S.A. following the 1980 eruption. Journal of the North American Benthological Society 7:246–252.
- Hayes, M.P. and Cliff, F.S. 1982. A checklist of the herpetofauna of Butte County, the Butte Sink, and Sutter Buttes, California. Herpetological Review 13:85–87.
- Hayes, M.P. and Jennings, M.R. 1986. Decline of ranid frog species in western North America: are bullfrogs (*Rana catesbeiana*) responsible? Journal of Herpetology 20:490–509.
- Hayes, M.P. and Jennings, M.R. 1988. Habitat correlates of distribution of the California red-legged frog (*Rana aurora draytonii*) and the foothill yellow-legged frog (*Rana boylei*): implications for management. in Szaro, R.C., Severson, K.E., and Patton, D.R. (Eds): Management of Amphibians, Reptiles, and Small Mammals in North America, General Technical Report RM-166. United States Department of Agriculture Forest Service, Rocky Mountain Forest and Range Experiment Station, Fort Collins, CO, pp.144–158.
- Hayes, M.P. and Krempels, D.M. 1986. Vocal sac variation among frogs of the genus *Rana* from western North America. Copeia 1986:927–936.
- Hayes, M.P. and Miyamoto, M.M. 1984. Biochemical, behavioral and body size differences between *Rana aurora aurora* and *R. a. draytonii*. Copeia 1984:1018–1022.
- Hayes, M.P., Pearl, C.A., and Rombough, C.J. 2001. *Rana aurora aurora*. Movement. Herpetological Review 32:35–36.
- Hayes, M.P., Quinn, T., Dugger, D.J., Hicks, T.L., Melchior, M.A., and Runde, D.E. 2006. Dispersion of coastal tailed frog (*Ascaphus truei*): an hypothesis relating occurrence of frogs in non-fish-bearing headwater basins to their seasonal movements. Journal of Herpetology 40:531–543.
- Hayes, M.P., Rombough, C.J., and Hayes, C.P. 2007. Natural History: *Rana aurora*. Movement. Herpetological Review 38:192–193.
- Hayes, M.P., Rombough, C.J., Padgett-Flohr, G.E., Hallock, L.A., Johnson, J.E., Wagner, R.S., and Engler, J.D. 2009. Amphibian chytridiomycosis in the Oregon spotted frog (*Rana pretiosa*) in Washington State, U.S.A. Northwestern Naturalist 90:148–151.
- Hayes, M.P. and Tennant, M.R. 1985. Diet and feeding behavior of the California red-legged frog, *Rana aurora draytonii* (Ranidae). The Southwestern Naturalist 30:601–605.
- Hayhoe, K., Cayan, D., Field, C.B., Frumhoff, P.C., Maurer, E.P., Miller, N.L., Moser, S.C. et al. 2004. Emissions pathways, climate change, and impacts on California. Proceedings of the National Academy of Sciences of the United States of America 101:12422–12427.
- Heath, J.E. 1965. Temperature regulation and diurnal activity in horned lizards. University of California Publications in Zoology 64:97–136.
- Hedgecock, D. 1978. Population subdivision and genetic divergence in red-bellied newt, *Taricha rivularis*. Evolution 32:271–286.
- Hedgecock, D. and Ayala, F.J. 1974. Evolutionary divergence in the genus *Taricha* (Salamandridae). Copeia 1974:738–747.
- Heifetz, W. 1941. A review of the lizards of the genus *Uma*. Copeia 1941:99–111.
- Hels, T. and Buchwald, E. 2001. The effect of road kills on amphibian populations. Biological Conservation 99:331–340.
- Hemphill, D.V. 1952. The Vertebrate Fauna of the Boreal Areas of the Southern Yolla Bolly

- Mountains, California. PhD Dissertation. Oregon State University, Corvallis.
- Henry, W.V. and Twitty, V.C. 1940. Contributions to the life histories of *Dicamptodon ensatus* and *Ambystoma gracile*. *Copeia* 1940:247-250.
- Hensley, F.R., Jones, T.R., Maxwell, M.S., Adams, L.J., and Nedella, N.S. 2010. Demography, terrestrial behavior, and growth of Sonora mud turtles (*Kinosternon sonoriense*) in an extreme habitat. *Herpetological Monographs* 24:174-193.
- Hensley, M.M. 1949. Mammal diet of *Heloderma*. *Herpetologica* 5:152.
- Heyer, W.R., Donnelly, M.A., McDiarmid, R.W., Hayek, L.A.C., and Foster, M.S. (Eds). 1994. Measuring and Monitoring Biological Diversity: Standard Methods for Amphibians. Smithsonian Institution Press, Washington, DC.
- Highton, R. 2000. Detecting cryptic species using allozyme data, in Bruce, J.C., Jaeger, R.G., and Houck, L.D. (Eds): *The Biology of Plethodontid Salamanders*. Kluwer Academic/Plenum Publishing, New York, pp.215-241.
- Hill, R.E. and Mackessy, S.P. 2000. Characterization of venom (Duvernoy's secretion) from twelve species of colubrid snakes and partial sequence of four venom proteins. *Toxicon* 38:1663-1687.
- Hillis, D.M. 1988. Systematics of the *Rana pipiens* complex: puzzle and paradigm. *Annual Review of Ecology and Systematics* 19:39-63.
- Hillis, D.M., Frost, J.S., and Wright, D.A. 1983. Phylogeny and biogeography of the *Rana pipiens* complex: a biochemical evaluation. *Systematic Zoology* 32:132-143.
- Hillis, D.M. and Wilcox, T.P. 2005. Phylogeny of the New World true frogs (*Rana*). *Molecular Phylogenetics and Evolution* 34:299-314.
- HilleRisLambers, J., Yelenik, S.G., Colman, B.P., and Levine, J.M. 2010. California annual grass invaders: the drivers or passengers of change? *Journal of Ecology* 98:1147-1156.
- Hilton, W.A. 1948. Salamander notes from the Northwest. *Herpetologica* 4:120.
- Hodge, R.P. 2004. Geographic distribution: *Rana aurora*. *Herpetological Review* 35:79.
- Holcomb, C. 2012. Examining the Relationship between Landscape Connectivity and the Breeding Effort of the Red-Legged Frog (*Rana aurora*) in Western Washington Wetlands. Master's Thesis. Evergreen State College, Olympia, WA.
- Holland, D.C. 1989. A Synopsis of the Ecology and Current Status of the Western Pond Turtle (*Clemmys marmorata*). Report. United States Fish and Wildlife Service National Ecology Research Center, Fort Collins, CO.
- Holland, D.C. 1992a. Level and Pattern in Morphological Variation: A Phylogeographic Study of the Western Pond Turtle (*Clemmys marmorata*). PhD Dissertation. University of Southwestern Louisiana, Lafayette.
- Holland, D.C. 1992b. A Synopsis of the Ecology and Status of the Western Pond Turtle (*Clemmys marmorata*) in 1991. United States Fish and Wildlife Service, San Simeon, CA.
- Holland, D.C. 1994. The Western Pond Turtle: Habitat and History. Report. United States Department of Energy, Bonneville Power Administration Environment, Portland, OR.
- Holman, J.A. and Fritz, U. 2001. A new emydid species from the Middle Miocene (Barstovian) of Nebraska, U.S.A. with a new generic arrangement for the species of *Clemmys* sensu McDowell (1964) (Reptilia: Testudines: Emydidae). *Zoologische Abhandlungen-Staatliches Museum Fur Tierkunde in Dresden* 51:331-353.
- Holte, D.L. 1998. Nest Site Characteristics of the Western Pond Turtle, *Clemmys marmorata*, at Fern Ridge Reservoir, in West Central Oregon. Master's Thesis. Oregon State University, Corvallis.
- Holway, D.A. 1995. Distribution of the argentine ant (*Linepithema humile*) in Northern California. *Conservation Biology* 9:1634-1637.
- Holway, D.A. 1998. Effect of argentine ant invasions on ground-dwelling arthropods in northern California riparian woodlands. *Oecologia* 116:252-258.
- Hossack, B.R., Corn, P.S., and Fagre, D.B. 2006. Divergent patterns of abundance and age-class structure of headwater stream tadpoles in burned and unburned watersheds. *Canadian Journal of Zoology* 84:1482-1488.
- Hothorn, R.L., Meckstroth, A.M., Wegner, K.E., Jennings, M.R., and Crayon, J.J. 2009. Diets of three species of anurans from the Cache Creek watershed, California, U.S.A. *Journal of Herpetology* 43:275-283.
- Howard, C.W. 1974. Comparative reproductive ecology of horned lizards (genus *Phrynosoma*) in southwestern United States and northern Mexico. *Journal of the Arizona Academy of Sciences* 9:108-116.
- Howard, J.H., and Wallace, R.L. 1985. Life history characteristics of populations of the long-toed salamander (*Ambystoma macrodactylum*) from different altitudes. *The American Midland Naturalist* 113:361-373.
- Hughes, L. 2000. Biological consequences of global warming: is the signal already apparent? *Trends in Ecology and Evolution* 15:56-61.

- Hulse, A.C. 1974. Food habits and feeding behavior in *Kinosternon sonoriense* (Chelonina; Kinosternidae). *Journal of Herpetology* 8:195-199.
- Hulse, A.C. 1982. Reproduction and population structure in the turtle, *Kinosternon sonoriense*. *The Southwestern Naturalist* 27:447-456.
- Hunt, L.E. 1983. A nomenclatural rearrangement of the genus *Anniella* (Sauria: Anniellidae). *Copeia* 1983:79-89.
- Hunt, L.E. 1984. Morphological Variation in the Fossorial Lizard *Anniella*. Master's Thesis. University of Kansas, Lawrence.
- ICF Jones and Stokes. 2010. Hatchery and Stocking Program Environmental Impact Report/ Environmental Impact Statement. Final. January. (ICF J&S 00264.08) (SCH #2008082025). Prepared for the California Department of Fish and Game and US Fish and Wildlife Service, Sacramento.
- ICZN (International Commission on Zoological Nomenclature). 2000. Opinion 1960 *Crotalus ruber* Cope 1892 (Reptilia, Serpentes). specific name given precedence over that of *Crotalus exsul* Garman, 1884. *Bulletin of Zoological Nomenclature* 57:189-190.
- Iverson, J.B. 1976. *Kinosternon sonoriense*. *Catalogue of American Amphibians and Reptiles* 176:1-2.
- Jacobson, E.R. 1993. Implications of infectious diseases for captive propagation and introduction programs of threatened/endangered reptiles. *Journal of Zoo and Wildlife Medicine* 24:245-255.
- Jacobson, E.R. and Whitford, W.G. 1971. Physiological responses to temperature in the patch-nosed snake, *Salvadora hexalepis*. *Herpetologica* 27:289-295.
- Jaeger, J.R., Riddle, B.R., Jennings, R.D., and Bradford, D.F. 2001. Rediscovering *Rana onca*: evidence for phylogenetically distinct leopard frogs from the border region of Nevada, Utah, and Arizona. *Copeia* 2001:339-354.
- Jancowski, K. and Orchard, S. 2013. Stomach contents from invasive American bullfrogs *Rana catesbeiana* (= *Lithobates catesbeianus*) on southern Vancouver Island, British Columbia, Canada. *NeoBiota* 16:17-37.
- Janzen, F.J., Krenz, J.G., Haselkorn, T.S., and Brodie, E.D. 2002. Molecular phylogeography of common garter snakes (*Thamnophis sirtalis*) in western North America: implications for regional historical forces. *Molecular Ecology* 11:1739-1751.
- Jayne, B.C. and Bennett, A.F. 1990. Selection on locomotor performance capacity in a natural population of garter snakes. *Evolution* 44:1204-1229.
- Jennings, M.R. 1984. Longevity records for lizards of the family Helodermatidae. *Bulletin of the Maryland Herpetological Society* 20:22-23.
- Jennings, M.R. 1987. Impact of the curio trade for San Diego horned lizards (*Phrynosoma coronatum blainvillii*) in the Los Angeles basin, California: 1885-1930. *Journal of Herpetology* 21:356-358.
- Jennings, M.R. 1988a. Natural history and decline of native ranids in California, in De Lisle, H.F., Brown, P.R., Kanfman, B., and McGurty, B.M. (Eds): *Proceedings of the Conference on California Herpetology*. Special Publication No 4. Southwestern Herpetologists Society, Van Nuys, CA, pp.61-72.
- Jennings, M.R. 1988b. Origin of the population of *Rana aurora draytonii* on Santa Cruz Island, California. *Herpetological Review* 19:76.
- Jennings, M.R. 1988c. *Phrynosoma coronatum*. *Catalogue of American Amphibians and Reptiles* 428:1-5.
- Jennings, M.R. and Fuller, M.M. 2004. Origin and distribution of leopard frogs, *Rana pipiens* complex, in California. *California Fish and Game* 90:119-139.
- Jennings, M.R. and Hayes, M.P. 1985. Pre-1900 overharvest of California red-legged frogs (*Rana aurora draytonii*): the inducement for bullfrog (*Rana catesbeiana*) introduction. *Herpetologica* 41:94-103.
- Jennings, M.R. and Hayes, M.P. 1994a. Amphibian and Reptile Species of Special Concern in California. California Department of Fish and Game, Inland Fisheries Division, Rancho Cordova.
- Jennings, M.R. and Hayes, M.P. 1994b. Decline of native ranid frogs in the desert Southwest, in Brown, P.R. and Wright, J.W. (Eds): *Herpetology of the North American Deserts*, *Proceedings of a Symposium*. Special Publication No 5. Southwestern Herpetologists Society, Van Nuys, CA, pp.183-211.
- Jennings, M.R. and Hayes, M.P. 2005. Natural history notes: *Rana boylei*. Coloration. *Herpetological Review* 36:438.
- Jockusch, E.L., Martinez-Solano, J., Hansen, R.W., and Wake, D.B. 2012. Morphological and molecular diversification of slender salamanders (Caudata: Plethodontidae: *Batrachoseps*) in the southern Sierra Nevada of California with descriptions of two new species. *Zootaxa* 3190:1-30.
- Jockusch, E.L. and Wake, D.B. 2002. Falling apart and merging: diversification of slender salamanders (Plethodontidae: *Batrachoseps*) in the American West. *Biological Journal of the Linnean Society* 76:361-391.

- Jockusch, E.L., Wake, D.B., and Yanev, K.P. 1998. New species of *Batrachoseps* (Caudata: Plethodontidae) from the Sierra Nevada, California. Los Angeles County Museum Contributions in Science 472:1-17.
- Jockusch, E.L., Yanev, K.P., and Wake, D.B. 2001. Molecular phylogenetic analysis of slender salamanders, genus *Batrachoseps* (Amphibia: Plethodontidae), from central coastal California with descriptions of four new species. Herpetological Monographs 15:54-99.
- Johnson, J., Thomson, R.C., Micheletti, S., and Shaffer, H.B. 2010. The origin of tiger salamander populations in California: introductions or relicts? Conservation Genetics 12:35-370.
- Jones, K.B. 1983. Movement patterns and foraging ecology of Gila monsters (*Heloderma suspectum* Cope) in northwestern Arizona. Herpetologica 39:247-253.
- Joseph, M.B., Piovia-Scott, J., Lawler, S.P., Pope, K.L. 2011. Indirect effects of introduced trout on Cascades frogs (*Rana cascadae*) via shared aquatic prey. Freshwater Biology 56:828-838.
- Kagarise Sherman, C. 1980. A Comparison of the Natural History and Mating System of Two Anurans: Yosemite Toads (*Bufo canorus*) and Black Toads (*Bufo exsul*). PhD Dissertation. University of Michigan, Ann Arbor.
- Kagarise Sherman, C. and Morton, M.L. 1984. The toad that stays on its toes. Natural History 93:72-78.
- Kagarise Sherman, C. and Morton, M.L. 1993. Population declines of Yosemite toads in the eastern Sierra Nevada of California. Journal of Herpetology 27:186-198.
- Karlstrom, E.L. 1962. The toad genus *Bufo* in the Sierra Nevada of California: ecological and systematic relationships. University of California Publications in Zoology 62: 1-104.
- Karlstrom, E.L. and Livezey, R.L. 1955. The eggs and larvae of the Yosemite toad *Bufo canorus* Camp. Herpetologica 11:221-227.
- Karraker, N.E. 1999. Natural history notes: *Rhyacotriton variegatus*. Nest site. Herpetological Review 30:160-161.
- Karraker, N.E., Pilliod, D.S., Adams, M.J., Bull, E.L., Coru, P.S., Diller, L.V., Dupuis, L.A. et al. 2006. Taxonomic variation in oviposition by tailed frogs (*Ascaphus* spp.). Northwestern Naturalist 87:87-97.
- Kats, L.B., Breeding, J.A., Hanson, K.M., and Smith, P. 1994. Ontogenetic changes in California newts (*Taricha torosa*) in response to chemical cues from conspecific predators. Journal of the North American Benthological Society 13:321-325.
- Kats, L.B., Elliott, S.A., and Currens, J. 1992. Intraspecific oophagy in stream-breeding California newts (*Taricha torosa*). Herpetological Review 23:7-8.
- Kelly, P.A., Phillips, S.E., and Williams, D.F. 2005. Documenting ecological change in time and space: the San Joaquin Valley of California, in Lacey, E.A. and Myers, P. (Eds): Mammalian Diversification: from Chromosomes to Phylogeography (a Celebration of the Career of James L. Patton), University of California Publications in Zoology, Berkeley, pp.57-78.
- Kennicott, R. 1860. Descriptions of new species of North American serpents in the Museum of the Smithsonian Institution, Washington. Proceedings of the Academy of Natural Sciences of Philadelphia 12:328-338.
- Kephart, D.G. and Arnold, S.J. 1982. Garter snake diets in a fluctuating environment: a seven-year study. Ecology 63:1232-1236.
- Kerby, J.L. and Kats, L. 1998. Modified interactions between salamander life stages caused by wildfire-induced sedimentation. Ecology 79:740-745.
- Kerby, J.L., Richards-Hrdlicka, K.L., Storfer, A., and Skelly, D.K. 2010. An examination of amphibian sensitivity to environmental contaminants: are amphibians poor canaries? Ecology Letters 13:60-67.
- Kessel, E.L. and Kessel, B.B. 1943a. The rate of growth of the young larvae of the Pacific giant salamander, *Dicamptodon ensatus* (Eschscholtz). The Wasmann Collector 5:108-111.
- Kessel, E.L. and Kessel, B.B. 1943b. The rate of growth of the older larvae of the Pacific giant salamander, *Dicamptodon ensatus* (Eschscholtz). The Wasmann Collector 5:141-142.
- Kessel, E.L. and Kessel, B.B. 1944. Metamorphosis of the Pacific giant salamander, *Dicamptodon ensatus* (Eschscholtz). The Wasmann Collector 6:38-48.
- Kiesecker, J.M. and Blaustein, A.R. 1998. Effects of introduced bullfrogs and smallmouth bass on microhabitat use, growth, and survival of native red-legged frogs (*Rana aurora*). Conservation Biology 12:776-787.
- Kiesecker, J.M. and Blaustein, A.R. 1999. Pathogen reverses competition between larval amphibians. Ecology 80:2442-2448.
- Kimball, S. and Schiffman, P.A. 2003. Differing effects of cattle grazing on native and alien plants. Conservation Biology 17:1681-1693.

- King, K.A., Martinez, C.T., and Rosen, P.C. 1996. Contaminants in Sonoran mud turtles from Quitobaquito Springs, Organ Pipe Cactus National Monument, Arizona. United States Fish and Wildlife Service, Region 2, Phoenix, AZ.
- King, T. and Robbins, M. 1991. A status survey of *Kinosternon sonoriense* and *Bufo alvarius* along the California side of the lower Colorado River Basin. California Department of Fish and Game, Inland Fisheries, Rancho Cordova.
- Kingsbury, B.A. 1989. Factors influencing activity in *Coleonyx variegatus*. Journal of Herpetology 23:399-404.
- Kirk, J.J. 1988. Western spotted frog (*Rana pretiosa*) mortality following forest spraying of DDT. Herpetological Review 19:51-53.
- Klauber, L.M. 1924. Notes on the distribution of snakes in San Diego County, California. Bulletin of the Zoological Society of San Diego 1:1-23.
- Klauber, L.M. 1928. A list of the amphibians and reptiles of San Diego County, California. Bulletin of the Zoological Society of San Diego 4:1-8.
- Klauber, L.M. 1929. Range extensions in California. Copeia 1929:15-22.
- Klauber, L.M. 1930. A list of the amphibians and reptiles of San Diego County, California. Bulletin of the Zoological Society of San Diego 5:1-8.
- Klauber, L.M. 1931. A statistical survey of the snakes of the southern border of California. Bulletin of the Zoological Society of San Diego 8:1-93.
- Klauber, L.M. 1934. An annotated list of the amphibians and reptiles of the southern border of California. Bulletin of the Zoological Society of San Diego 11:1-28.
- Klauber, L.M. 1936. The horned lizards of the coronatum group. Copeia 1936:103-110.
- Klauber, L.M. 1939. Studies of reptile life in the arid southwest. Bulletin of the Zoological Society of San Diego 14:1-100.
- Klauber, L.M. 1945. The geckos of the genus *Coleonyx* with descriptions of new subspecies. Transactions of the San Diego Society of Natural History 10:133-216.
- Klauber, L.M. 1946. The glossy snake. *Arizona*, with descriptions of new subspecies. Transactions of the San Diego Society of Natural History 10:311-398.
- Klauber, L.M. 1956. Rattlesnakes: Their Habits, Life Histories and Influence on Mankind. Vols. 1-2, University of California Press, Los Angeles.
- Kluge, A.G. 1966. A new pelobatine frog from the lower Miocene of South Dakota with a discussion of the evolution of the *Scaphiopus-Spea* complex. Los Angeles County Museum of Natural History 113:1-26.
- Knapp, R.A. 2005. Effects of nonnative fish and habitat characteristics on lentic herpetofauna in Yosemite National Park, U.S.A. Biological Conservation 121:265-279.
- Knapp, R.A. and Matthews, K.R. 2000. Non-native fish introductions and the decline of the mountain yellow-legged frog from within protected areas. Conservation Biology 14:428-438.
- Knowles, N. and Cayan, D.R. 2002. Potential effects of global warming on the Sacramento/San Joaquin watershed and the San Francisco estuary. Geophysical Research Letters 29:1891-1895.
- Knowles, N. and Cayan, D.R. 2004. Elevational dependence of projected changes in the San Francisco estuary and watershed. Climatic Change 62:319-336.
- Knowlton, G.F. 1944. Some insect food of *Rana pipiens*. Copeia 1944:119.
- Kuchta, S.R. 2005. *Taricha torosa*, in Lannoo, M.J. (Ed.): Amphibian Declines: The Conservation Status of United States Species, University of California Press, Berkeley, pp.904-908.
- Kuchta, S.R. 2007. Contact zones and species limits: hybridization between lineages of the California newt, *Taricha torosa*, in the southern Sierra Nevada. Herpetologica 63:332-350.
- Kuchta, S.R. and Tan, A.M. 2006a. Limited genetic variation across the range of the red-bellied newt, *Taricha rivularis*. Journal of Herpetology 40:561-565.
- Kuchta, S.R. and Tan, A.M. 2006b. Lineage diversification on an evolving landscape: phylogeography of the California newt, *Taricha torosa* (Caudata: Salamandridae). Biological Journal of the Linnean Society 89:213-239.
- Kuhnz, L. 2000. Microhabitats and Home Range of the California Legless Lizard Using Biotelemetry. Master's Thesis. California State University, San Jose.
- Kuhnz, L., Burton, R., Slattery, P., and Oakden, J. 2005. Microhabitats and population densities of California legless lizards, with comments on effectiveness of various techniques for estimating numbers of fossorial reptiles. Journal of Herpetology 39:395-402.
- Kupferberg, S.J. 1996a. Hydrologic and geomorphic factors affecting conservation of a river-breeding frog (*Rana boylei*). Ecological Applications 6:1332-1344.
- Kupferberg, S.J. 1996b. The Ecology of Native Tadpoles (*Rana boylei* and *Hyla regilla*) and the

- Impact of Invading Bullfrogs (*Rana catesbeiana*) in a Northern California River. PhD Dissertation. University of California, Berkeley.
- Kupferberg, S.J. 1997. Bullfrog (*Rana catesbeiana*) Invasion of a California River: The Role of Larval Competition. *Ecology* 78:1736–1751.
- Kupferberg, S.J., Catenazzi, A., Lunde, K., Lind, A.J., and Palen, W.J. 2009a. Parasitic copepod (*Lernaea cyprinacea*) outbreaks in foothill yellow-legged frogs (*Rana boylei*) linked to unusually warm summers and amphibian malformations in northern California. *Copeia* 2009:529–537.
- Kupferberg, S.J., Lind, A.J., Mount, J., and Yarnell, S. 2009b. Pulsed Flow Effects on Foothill Yellow-Legged Frog (*Rana boylei*): Integration of Empirical, Experimental, and Hydrodynamic Modeling Approaches. Report. California Energy Commission, PIER, Sacramento.
- Kupferberg, S.J., Lind, A.J., and Palen, W.J. 2009c. Pulsed Flow Effects on the Foothill Yellow-legged frog (*Rana boylei*): Population Modeling. Report. California Energy Commission, PIER, Sacramento.
- Kupferberg, S.J., Lind, A.J., Thill, V., and Yarnell, S.M. 2011. Water velocity tolerance in tadpoles of the foothill yellow-legged frog (*Rana boylei*): swimming performance, growth and survival. *Copeia* 2011:141–152.
- Kupferberg, S.J., Palen, W.J., Lind, A.J., Bobzien, S., Catenazzi, A., Drennan, J., and Power, M.E. 2012. Effects of flow regimes altered by dams on survival, population declines, and range-wide losses of California river-breeding frogs. *Conservation Biology* 26:513–524.
- La Berteaux, D.L. and Garlinger, B.H. 1998. Inyo California Towhee (*Pipilo crissalis eremophilus*) Census in the Argus and Coso Mountain Ranges, Inyo County, California. Contract N2474-90-M-3113. Naval Air Weapons Station, China Lake, CA.
- La Rivers, I. 1942. Some new amphibian and reptile records for Nevada. *Journal of Entomology and Zoology* 34:53–68.
- Lacan, I., Matthews, K., and Feldman, K. 2008. Interaction of an introduced predator with future effects of climate change in the recruitment dynamics of the imperiled Sierra Nevada yellow-legged frog (*Rana sierrae*). *Herpetological Conservation and Biology* 3:211–223.
- Lande, R. 1988. Genetics and demography in biological conservation. *Science* 241:1455.
- Langen, T.A., Ogden, K.M., and Schwarting, L.L. 2009. Predicting hot spots of herpetofauna road mortality along highway networks. *Journal of Wildlife Management* 73:104–114.
- Lannoo, M.J., ed. 2005. *Amphibian Declines: The Conservation Status of United States Species*. University of California Press, Berkeley.
- Lappin, A.K. and Swinney, E.J. 1999. Sexual dimorphism as it relates to natural history of leopard lizards (Crotaphytidae: *Gambelia*). *Copeia* 1999:649–660.
- Larsen, K.W., Gregory, P.T., and Antoniak, R. 1993. Reproductive ecology of the common garter snake *Thamnophis sirtalis* at the northern limit of its range. *The American Midland Naturalist* 129:336–345.
- Larson, A. 1980. Paedomorphosis in relation to rates of morphological and molecular evolution in the salamander *Aneides flavipunctatus* (Amphibia, Plethodontidae). *Evolution* 34:1–17.
- Larson, M.D. 2012. Diet of the Cascades Frog (*Rana cascadae*) as It Relates to Prey Availability in the Klamath Mountains of Northwest California. Master's Thesis. Humboldt State University, Arcata.
- Larson, N.M. 1984. Geographic Variation in the Two-Striped Garter Snake, *Thamnophis hammondi*. Master's Thesis. California State Polytechnic University, Pomona.
- Lawler, J.J., Shafer, S.L., Bancroft, B.A., and Blaustein, A.R. 2010. Projected climate impacts for the amphibians of the western hemisphere. *Conservation Biology* 24:38–50.
- Lawler, S.P., Dritz, D., Strange, T., and Holyoak, M. 1999. Effects of introduced mosquitofish and bullfrogs on the threatened California red-legged frog. *Conservation Biology* 13:613–622.
- Lawson, R. and Dessauer, H.C. 1979. Biochemical genetics and systematics of garter snakes of the *Thamnophis elegans*, *Thamnophis couchii* and *Thamnophis ordinoides* complex. Occasional Papers of the Museum of Zoology, Louisiana State University 56:1–24.
- Lazaroff, D.W., Rosen, P.C., and Lowe, C.H. 2006. *Amphibians, Reptiles, and Their Habitats at Sabino Canyon*. University of Arizona Press, Tucson.
- Le Conte, J. 1854. Description of four new species of *Kinosternum*. *Proceedings of the Academy of Natural Sciences of Philadelphia* 7:180–190.
- Leaché, A.D., Koo, M.S., Spencer, C.L., Papenfuss, T.J., Fisher, R.N., and McGuire, J.A. 2009. Quantifying ecological, morphological, and genetic variation to delimit species in the coast horned lizard species complex (*Phrynosoma*). *Proceedings of the National Academy of Sciences of the United States of America* 106:12418–12423.

- Leaché, A.D. and Mulcahy, D.G. 2007. Phylogeny, divergence times and species limits of spiny lizards (*Sceloporus magister* species group) in western North American deserts and Baja California. *Molecular Ecology* 16:5216–5233.
- Leatherman, B. 1996. Natural history notes: *Phrynosoma coronatum blainvillii*. *Conservation. Herpetological Review* 27:80.
- Leavitt, D.H., Bezy, R.L., Crandall, K.A., and Sites, J. 2007. Multi-locus DNA sequence data reveal a history of deep cryptic vicariance and habitat-driven convergence in the desert night lizard *Xantusia vigilis* species complex (Squamata: Xantusiidae). *Molecular Ecology* 16:4455–4481.
- Lebassi, B., Gonzalez, J., Fabris, D., Maurer, E., Miller, N., Milesi, C., Switzer, P., and Bornstein, R. 2009. Observed 1970–2005 cooling of summer daytime temperatures in coastal California. *Journal of Climate* 22:3558–3573.
- Lee, J.C. 1975. The autecology of *Xantusia henshawi henshawi* (Sauria: Xantusiidae). *Transactions of the San Diego Society of Natural History* 17:259–278.
- Leidy, R.A., Gonsolin, E., and Leidy, G.A. 2009. Late-summer aggregation of the foothill yellow-legged frog (*Rana boylei*) in central California. *The Southwestern Naturalist* 54:367–368.
- Lemm, J.M. 2006. *Field Guide to Amphibians and Reptiles of the San Diego Region*. University of California Press, Berkeley.
- Lenihan, J.M., Bachelet, D., Neilson, R.P., and Drapek, R. 2008. Response of vegetation distribution, ecosystem productivity, and fire to climate change scenarios for California. *Climatic Change* 87:5215–5230.
- Leviton, A.E. and Banta, B.H. 1964. Mid-winter reconnaissance of the herpetofauna of the Cape Region of Baja California, Mexico. *Proceedings of the California Academy of Sciences* 4:127–156.
- Leyse, K. 2005. *Intentional Introductions and Biodiversity in Fishless Waters: The Effects of Introduced Fish on Native Aquatic Species*. PhD Dissertation. University of California, Davis.
- Licht, L.E. 1969. Comparative breeding behavior of the red-legged frog (*Rana aurora aurora*) and the western spotted frog (*Rana pretiosa pretiosa*) in southwestern British Columbia. *Canadian Journal of Zoology* 47:1287–1299.
- Licht, L.E. 1971. Breeding habits and embryonic thermal requirements of the frogs, *Rana aurora aurora* and *Rana pretiosa pretiosa*, in the Pacific Northwest. *Ecology* 52:116–124.
- Licht, L.E. 1974. Survival of embryos, tadpoles, and adults of the frogs *Rana aurora aurora* and *Rana pretiosa pretiosa* sympatric in southwestern British Columbia. *Canadian Journal of Zoology* 52:613–627.
- Licht, L.E. 1986a. Comparative escape behavior of sympatric *Rana aurora* and *Rana pretiosa*. *The American Midland Naturalist* 115:239–247.
- Licht, L.E. 1986b. Food and feeding behavior of sympatric red-legged frogs, *Rana aurora*, and spotted frogs, *Rana pretiosa*, in southwestern British Columbia. *Canadian Field-Naturalist* 100:22–31.
- Licht, L.E. 1991. Habitat selection of *Rana pipiens* and *Rana sylvatica* during exposure to warm and cold temperatures. *The American Midland Naturalist* 125:259–268.
- Licht, P. and Brown, A. 1967. Behavioral thermoregulation and its role in ecology of red-bellied newt, *Taricha rivularis*. *Ecology* 48:598–611.
- Ligon, D.B. and Peterson, C.C. 2002. Physiological and behavioral variation in estivation among mud turtles (*Kinosternon* spp.). *Physiological and Biochemical Zoology* 75:283–293.
- Ligon, D.B. and Stone, P. 2003. Radiotelemetry reveals terrestrial estivation in Sonoran mud turtles (*Kinosternon sonoriense*). *Journal of Herpetology* 37:750–754.
- Lind, A.J. 2005. *Reintroduction of a Declining Amphibian: Determining an Ecologically Feasible Approach for the Foothill Yellow-Legged Frog (*Rana boylei*) through Analysis of Decline Factors, Genetic Structure, and Habitat Associations*. PhD Dissertation. University of California, Davis.
- Lind, A.J., Bettaso, J.B., and Yarnell, S.M. 2003. Natural history notes: *Rana boylei* and *Rana catesbeiana*. Reproductive behavior. *Herpetological Review* 34:235.
- Lind, A.J., Spinks, P.Q., Fellers, G.M., and Shaffer, H.B. 2011. Rangelwide phylogeography and landscape genetics of the western U.S. endemic frog *Rana boylei* (Ranidae): implications for the conservation of frogs and rivers. *Conservation Genetics* 12:269–284.
- Lind, A.J., Welsh, H.H., Jr., and Wilson, R.A. 1996. The effects of a dam on breeding habitat and egg survival of the foothill yellow-legged frog (*Rana boylei*) in northwestern California. *Herpetological Review* 27:62–67.
- Linnaeus, C. 1758. *Systema Naturae*, Edition 10. Impensis Direct. Laurentii Salvii, Stockholm. (In Swedish)
- Linsdale, J.M. 1932. *Amphibians and reptiles from Lower California*. University of California Publications in Zoology 38:345–386.
- Linzey, D.W. 1967. Food of the leopard frog, *Rana pipiens*, in central New York. *Herpetologica* 23:11–17.

- Liss, W.J. and Larson, G.L. 1991. Ecological effects of stocked trout on north Cascades naturally fishless lakes. *Park Science* 11:22-23.
- Liss, W.J., Larson, G.L., Deimling, F., Ganio, L.M., Gresswell, R., Hoffman, R., Kiss, M. et al. 1995. Ecological effects of stocked trout in naturally fishless high mountain lakes. Technical Report NPS/PNROSU/NRTR-95-03. North Cascades National Park Service Complex, United States National Park Service, Seattle, WA.
- Livezey, R.L. 1955. A northward range extension for *Bufo canorus* Camp. *Herpetologica* 11:212.
- Lockington, W.N. 1879. Notes on some reptiles and Batrachia of the Pacific Coast. *American Naturalist* 13:780-783.
- Loomis, R.B. 1965. The yellow-legged frog, *Rana boylei*, from the Sierra San Pedro Martir, Baja California Norte, Mexico. *Herpetologica* 21:78-80.
- Lovich, J. and Meyer, K. 2002. The western pond turtle (*Clemmys marmorata*) in the Mojave River, California, U.S.A.: highly adapted survivor or tenuous relict? *Journal of Zoology* 256:537-545.
- Lovich, J.E. and Beaman, K.R. 2007. A history of Gila monster (*Heloderma suspectum cinctum*) records from California with comments on factors affecting their distribution. *Bulletin of the Southern California Academy of Sciences* 106:39-58.
- Lovich, J.E. and Beaman, K.R. 2008. Distribution of native turtles in the arid southwestern United States with comments on *Kinosternon sonoriense*: a species presumed to be lost from California's herpetofauna, in Reynolds, R.E. (Ed.): The 2008 Desert Symposium Field Guide and Proceedings. California State University, Desert Studies Consortium and LSA Associates, Soda Springs, pp.127-134.
- Lovich, J.E., Drost, C., Monatesti, A.J., Casper, D., Wood, D.A., and Girard, M. 2010. Reptilian prey of the Sonora mud Turtle (*Kinosternon sonoriense*) with comments on saurophagy and ophiophagy in North American turtles. *The Southwestern Naturalist* 55:135-138.
- Lovich, J.E. and Haxel, G. 2011. A previously unreported locality record for the Gila Monster (*Heloderma suspectum*). *Bulletin of the Southern California Academy of Sciences* 110:59-62.
- Lovich, J.E., Madrak, S.V., Drost, C.A., Monatesti, A.J., Casper, D., and Znari, M. 2012. Optimal egg size in a suboptimal environment; reproductive ecology of female Sonora mud turtles (*Kinosternon sonoriense*) in central Arizona, USA. *Amphibia-Reptilia* 33:161-170.
- Lovich, R.E. 2001. Phylogeography of the night lizard, *Xantusia henshawi*, in southern California: evolution across fault zones. *Herpetologica* 57:470-487.
- Lovich, R.E. 2009a. Phylogeography and Conservation of the Arroyo Toad (*Bufo californicus*). PhD Dissertation. Loma Linda University, Loma Linda, CA.
- Lovich, R.E. 2009b. Sandstone night lizard, in Jones, L.L.C. and Lovich, R.E. (Eds): *Lizards of the American Southwest*. Rio Nuevo, Tucson, AZ, pp.420-423.
- Lovich, R.E. and Grismer, L.L. 2001. *Xantusia gracilis*. *Catalogue of American Amphibians and Reptiles* 772:1-2.
- Lubcke, G.M. and Wilson, D.S. 2007. Variation in shell morphology of the western pond turtle (*Actinemys marmorata* Baird and Girard) from three aquatic habitats in northern California. *Journal of Herpetology* 41:107-114.
- Luckenbach, R.A. and Bury, R.B. 1983. Effects of off-road vehicles on the biota of the Algodones Dunes, Imperial County, California. *The Journal of Applied Ecology* 20:265-286.
- Lynch, J.F. 1974. Ontogenetic and Geographic Variation in the Morphology and Ecology of the Black Salamander (*Aneides flavipunctatus*). PhD Dissertation. University of California, Berkeley.
- Lynch, J.F. 1981. Patterns of ontogenetic and geographic variation in the black salamander, *Aneides flavipunctatus* (Caudata: Plethodontidae). *Smithsonian Contributions to Zoology* 324:1-53.
- Lynch, J.F. 1985. The feeding ecology of *Aneides flavipunctatus* and sympatric plethodontid salamanders in northwestern California. *Journal of Herpetology* 19:328-352.
- Macey, J.R. and Papenfuss, T.J. 1991a. Amphibians, in Hall, C.A. (Ed.): *Natural History of the White Inyo Range*. University of California Press, Berkeley, pp.277-290.
- Macey, J.R. and Papenfuss, T.J. 1991b. Reptiles, Hall, C.A. (Ed.): *Natural History of the White Inyo Range*. University of California Press, Berkeley, pp.291-360.
- Macey, J.R., Strasburg, J., Brisson, J., Vredenburg, V., Jennings, M., and Larson, A. 2001. Molecular phylogenetics of western North American frogs of the *Rana boylei* species group. *Molecular Phylogenetics and Evolution* 19:131-143.
- MacTague, L. and Northen, P.T. 1993. Underwater vocalization by the foothill yellow-legged frog (*Rana boylei*). *Transactions of the Western Section of the Wildlife Society* 29:1-7.
- Mahrtdt, C.R., Lovich, R., and Zimmitti, S.J. 2002. Natural history notes: *Bufo californicus*. *Habitat*

- and population status. *Herpetological Review* 33:123-125.
- Mahrtdt, C.R., McGuire, J.A., and Beaman, K.R. 2010. *Gambelia copeii*. Catalogue of American Amphibians and Reptiles 871:1-8.
- Marco, A. and Blaustein, A. 1999. The effects of nitrite on behavior and metamorphosis in Cascades frogs (*Rana cascadae*). *Environmental Toxicology and Chemistry* 18:946-949.
- Marco, A., Quilchano, C., and Blaustein, A. 1999. Sensitivity to nitrate and nitrite in pond-breeding amphibians from the Pacific Northwest, U.S.A. *Environmental Toxicology and Chemistry* 18:2836-2839.
- Marks, S.B. and Doyle, D. 2005. *Taricha rivularis*, in Lannoo, M.J. (Ed.): *Amphibian Declines: The Conservation Status of United States Species*. University of California Press, Berkeley, pp.901-908.
- Marlow, R.W., Brode, J.M., and Wake, D.B. 1979. A new salamander, genus *Batrachoseps*, from the Inyo Mountains of California; with a discussion of relationships in the genus. *Contributions in Science of the Natural History Museum of Los Angeles County* 308:1-17.
- Martin, D.L. 2008. *Decline, Movement and Habitat Utilization of the Yosemite Toad (Bufo canorus): An Endangered Anuran Endemic to the Sierra Nevada of California*. PhD Dissertation. University of California, Santa Barbara.
- Martin K.J. and McComb, B.C. 2003. Amphibian habitat associations at patch and landscape scales in the central Oregon Coast Range. *Journal of Wildlife Management* 67:672-683.
- Martin, R.F. 1972. Evidence from osteology, in Blair, W.F. (Ed.): *Evolution in the Genus Bufo*. University of Texas Press, Austin, pp.37-70.
- Marty, J.T. 2005. Effects of cattle grazing on diversity in ephemeral wetlands. *Conservation Biology* 19:1626-1632.
- Maslin, T.P. and Secoy, D.M. 1986. A checklist of the lizard genus *Cnemidophorus*. *Contributions in Zoology, University of Colorado Museum* 1:1-60.
- Matsuda, B.M. and Richardson, J.S. 2005. Movement patterns and relative abundance of coastal tailed frogs in clearcuts and mature forest stands. *Canadian Journal of Forest Research* 35:1131-1138.
- Maurer, E.P. 2007. Uncertainty in hydrologic impacts of climate change in the Sierra Nevada, California, under two emissions scenarios. *Climatic Change* 82:309-325.
- Maurer, E.P. and Duffy, P.B. 2005. Uncertainty in projections of streamflow changes due to climate change in California. *Geophysical Research Letters* 32:L03704.
- Mautz, W.J. 1979. The metabolism of reclusive lizards, the Xantusiidae. *Copeia* 1979:577-584.
- Mautz, W.J. and Case, T.J. 1974. A diurnal activity cycle in the granite night lizard, *Xantusia henshawi*. *Copeia* 1974:243-251.
- Mayhew, W.W. 1962. *Scaphiopus couchii* in California's Colorado Desert. *Herpetologica* 18:153-161.
- Mayhew, W.W. 1964a. Photoperiodic responses in three species of the lizard genus *Uma*. *Herpetologica* 20:95-113.
- Mayhew, W.W. 1964b. Taxonomic status of California populations of the lizard genus *Uma*. *Herpetologica* 20:170-183.
- Mayhew, W.W. 1965. Adaptations of the amphibian, *Scaphiopus couchii*, to desert conditions. *The American Midland Naturalist* 74:95-109.
- Mayhew, W.W. 1966. Reproduction in the psammophilous lizard *Uma scoparia*. *Copeia* 1966:114-122.
- Mayhew, W.W. 1968. Biology of desert amphibians and reptiles, in Brown, G.W.J. (Ed.): *Desert Biology*, Vol. 1. Academic Press, New York, pp.195-356.
- Mazerolle, M.J. 2004. Amphibian road mortality in response to nightly variations in traffic intensity. *Herpetologica* 60:45-53.
- Mazerolle, M.J., Bailey, L.L., Kendall, W.L., Andrew, R.J., Converse, S.J., and Nichols, J.D. 2007. Making great leaps forward: accounting for detectability in herpetological field studies. *Journal of Herpetology* 41:672-689.
- McAllister, K.R. and Leonard, W.P. 1997. Washington State status report for the Oregon spotted frog. Washington Department of Fish and Wildlife, Olympia. 38pp.
- McAllister, K.R., Leonard, W.P., and Storm, R.M. 1993. Spotted frog (*Rana pretiosa*) surveys in the Puget trough of Washington, 1989-1991. *Northwestern Naturalist* 74:10-15.
- McCauley, R.H. 1945. *The Reptiles of Maryland and the District of Columbia*. Free Press Printing Co., Burlington, VT.
- McClanahan, L., Jr. 1967. Adaptations of the spadefoot toad *Scaphiopus couchii*, to desert environments. *Comparative Biochemistry and Physiology* 20:73-99.
- McComb, W.C., McGarigal, K., and Anthony, R.G. 1993. Small mammal and amphibian abundance in streamside and upslope habitats of mature Douglas-fir stands, western Oregon. *Northwest Science* 67:7-15.
- McGrann, M., Wright, G., Dial, R., and McGrann, A. 2006. Off-highway vehicle impact on the

- flat-tailed horned lizard, *Phrynosoma mcallii*, in the Colorado Desert of southern California. *California Fish and Game* 92:67–80.
- McGuire, J.A. 1996. Phylogenetic systematics of crotaphytid lizards (Reptilia: Iguania: Crotaphytidae). *Bulletin of Carnegie Museum of Natural History* 32:1–143.
- McGuire, J.A. and Grismer, L.L. 1993. The taxonomy and biogeography of *Thamnophis hammondi* and *T. digueti* (Reptilia: Squamata: Colubridae) in Baja California, Mexico. *Herpetologica* 49:354–365.
- McGuire, J.A., Linkem, C.W., Koo, M.S., Hutchison, D.W., Lappin, A.K., Orange, D.I., Lemos-Espinal, J., Riddle, B.R., and Jaeger, J.R. 2007. Mitochondrial introgression and incomplete lineage sorting through space and time: phylogenetics of Crotaphytid lizards. *Evolution* 61:2879–2897.
- Metter, D.E. 1964. A morphological and ecological comparison of 2 populations of *Ascaphus truei* Stejneger. *Copeia* 1964:181–195.
- Miller, A.H. and Stebbins, R.C. 1964. *The Lives of Desert Animals in Joshua Tree National Monument*. University of California Press, Berkeley.
- Miller, C.M. 1944. Ecologic relations and adaptations of the limbless lizards of the genus *Anniella*. *Ecological Monographs* 14:271–289.
- Miller, D.A.W., Brehme, C.S., Hines, J.E., Nichols, J.D., and Fisher, R.N. 2012. Joint estimation of habitat dynamics and species interactions: disturbance reduces co-occurrence of non-native predators with an endangered toad. *The Journal of Animal Ecology* 81:1288–1297.
- Miller, M.P., Haig, S.M., and Wagner, R.S. 2006. Phylogeography and spatial genetic structure of the southern torrent salamander: implications for conservation and management. *Journal of Heredity* 97:561–570.
- Minnich, J.E. and Shoemaker, V.H. 1972. Water and electrolyte turnover in a field population of the lizard, *Uma scoparia*. *Copeia* 1972:650–659.
- Mitchell, J.C. 1994. *Reptiles of Virginia*. Smithsonian Books, Washington, DC.
- Mitrovich, M.J. 2006. A Case Study in Conservation Science: The Spatial Ecology and Evolutionary History of the Coachwhip Snake (*Masticophis flagellum*) and Striped Racer (*Masticophis lateralis*). PhD Dissertation. San Diego State University, San Diego, CA.
- Mitrovich, M.J., Diffendorfer, J.E., and Fisher, R.N. 2009. Behavioral response of the coachwhip (*Masticophis flagellum*) to habitat fragment size and isolation in an urban landscape. *Journal of Herpetology* 43:646–656.
- Mitrovich, M.J., Gallegos, E.A., Lyren, L.M., Lovich, R.E., and Fisher, R.N. 2011. Habitat use and movement of the endangered arroyo toad (*Anaxyrus californicus*) in Coastal Southern California. *Journal of Herpetology* 45:319–328.
- Mittleman, M.B. and Myers, G.S. 1949. Geographic variation in the ribbed frog, *Ascaphus truei*. *Proceedings of the Biological Society of Washington* 62:57–68.
- Monsen, K.J. and Blouin, M.S. 2003. Genetic structure in a montane ranid frog: restricted gene flow and nuclear-mitochondrial discordance. *Molecular Ecology* 12:3275–3286.
- Monsen, K.J. and Blouin, M.S. 2004. Extreme isolation by distance in a montane frog *Rana cascadae*. *Conservation Genetics* 5:827–835.
- Montanucci, R.R. 1965. Observations on the San Joaquin leopard lizard, *Crotaphytus wislizenii silus*. *Herpetologica* 21:270–283.
- Montanucci, R.R. 1968. Notes on the distribution and ecology of some lizards in the San Joaquin Valley, California. *Herpetologica* 24:316–320.
- Montanucci, R.R. 2004. Geographic variation in *Phrynosoma coronatum* (Lacertilia, Phrynosomatidae): further evidence for a peninsular archipelago. *Herpetologica* 60:117–139.
- Morafka, D.J. and Banta, B.H. 1976. Ecological relationships of the recent herpetofauna of Pinnacles National Monument, Monterey, and San Benito counties, California. *The Wasmann Journal of Biology* 34:304–324.
- Morey, S.R. 1998. Pool duration influences age and body mass at metamorphosis in the western spadefoot toad: implications for vernal pool conservation, in *Ecology, Conservation, and Management of Vernal Pool Ecosystems: Proceedings from a 1996 Conference*. California Native Plant Society, Sacramento, pp.86–91.
- Morey, S.R. 2005. *Spea hammondi*. in Lannoo, M.J. (Ed.): *Amphibian Declines: The Conservation Status of United States Species*. University of California Press, Berkeley, pp.514–517.
- Morey, S.R. and Guinn, D.A. 1992. Activity patterns, food habits, and changing abundance in a community of vernal pool amphibians. in Williams, D.F., Byrne, S., and Rado, T.A. (Eds): *Endangered and Sensitive Species of the San Joaquin Valley, California: Their Biology, Management, and Conservation*. California Energy Commission and the Wildlife Society, Western Section, Sacramento, pp.149–158.
- Morey, S.R. and Janes, D.N. 1994. Variation in larval habitat duration influences metamorphosis in *Scaphiopus couchii*, in Brown, P.R. and Wright, J.W. (Eds): *Herpetology of the North American*

- Deserts. Special Publication No. 5, Southwestern Herpetologists Society, Van Nuys, CA.
- Morey, S.R. and Reznick, D. 2001. Effects of larval density on postmetamorphic spadefoot toads (*Spea hammondi*). *Ecology* 82:510–522.
- Morey, S.R. and Reznick, D.N. 2004. The relationship between habitat permanence and larval development in California spadefoot toads: field and laboratory comparisons of developmental plasticity. *Oikos* 104:172–190.
- Morgan, J.A.T., Vredenburg, V.T., Rachowicz, L.J., Knapp, R.A., Stice, M.J., Tinninstall, T., Bingham, R.E. et al. 2007. Population genetics of the frog-killing fungus *Batrachochytrium dendrobatidis*. *Proceedings of the National Academy of Sciences of the United States of America* 104:13845–13850.
- Moritz, C. 1994. Defining “Evolutionarily Significant Units” for conservation. *Trends in Ecology and Evolution* 9:373–375.
- Morton, M.L. and Sokoloski, K.N. 1978. Sympatry in *Bufo boreas* and *Bufo canorus* and additional evidence of natural hybridization. *Bulletin of the Southern California Academy of Sciences* 77:52–55.
- Mosher, H.S., Fuhrman, F.A., Buchwald, H.D., and Fischer, H.G. 1964. Tarichatoxin-tetrodotoxin: a potent neurotoxin. *Science* 144:1100–1110.
- Moyle, P.B. 1973. Effects of introduced bullfrogs, *Rana catesbeiana*, on the native frogs of the San Joaquin Valley, California. *Copeia* 1973:18–22.
- Moyle, P.B. 2002. *Inland Fishes of California*. University of California Press, Berkeley.
- Moyle, P.B., Williams, J.E., and Wikramanayake, E.D. 1989. *Fish Species of Special Concern of California*. California Department of Fish and Game, Inland Fisheries Division, Rancho Cordova.
- Moyle, P.B., Yoshiyama, R.M., Williams, J.E., and Wikramanayake, E.D. 1995. *Fish species of special concern in California*. 2nd ed. California Department of Fish and Game, Inland Fisheries Division, Rancho Cordova.
- Mulcahy, D., Spaulding, A., Mendelson, J., and Brodie, E. 2006. Phylogeography of the flat-tailed horned lizard (*Phrynosoma mcallii*) and systematics of the *P. mcallii-platyrrhinus* mtDNA complex. *Molecular Ecology* 15:1807–1826.
- Mullally, D.P. 1953. Observations on the ecology of the toad *Bufo canorus*. *Copeia* 1953:182–183.
- Mullally, D.P. 1956. The relationships of the Yosemite and western toads. *Herpetologica* 12:133–135.
- Mullen, L.B., Woods, H.A., Schwartz, M.K., Sepulveda, A.J., and Lowe, W.H. 2010. Scale-dependent genetic structure of the Idaho giant salamander (*Dicamptodon aterrimus*) in stream networks. *Molecular Ecology* 19:898–909.
- Mullin, S.J., Imbert, H., Fish, J.M., Ervin, E.L., and Fisher, R.N. 2004. Snake (Colubridae: *Thamnophis*) predatory responses to chemical cues from native and introduced prey species. *The Southwestern Naturalist* 49:449–456.
- Murphy, R.W., Kovar, V., Haddrath, O., Allen, G., Fishbein, A., and Mandrak, N. 1995. mtDNA gene sequence, allozyme, and morphological uniformity among red diamond rattlesnakes, *Crotalus ruber* and *Crotalus exsul*. *Canadian Journal of Zoology* 73:270–281.
- Murphy, R.W., Trépanier, T.L., and Morafka, D.J. 2006. Conservation genetics, evolution and distinct population segments of the Mojave fringe-toed lizard, *Uma scoparia*. *Journal of Arid Environments* 67:226–247.
- Musgrave, M.E. 1930. *Bufo alvarius*, a poisonous toad. *Copeia* 1930:96–99.
- Muth, A. and Fisher, M. 1992. Development of Baseline Data and Procedures for Monitoring Populations of the Flat-Tailed Horned Lizard, *Phrynosoma mcallii*. Contract Report No. FG9268 to California Department of Fish and Game, Sacramento.
- Myers, C.W. 1965. Biology of the ringneck snake, *Diadophis punctatus*. Florida. *Bulletin of the Florida State Museum, Biological Sciences* 10:43–90.
- Myers, E.A., Rodríguez-Robles, J.A., Denardo, D.F., Staub, R.E., Stropoli, A., Ruane, S., and Burhink, F.T. 2013. Multilocus phylogeographic assessment of the California Mountain King-snake (*Lampropeltis zonata*) suggests alternative patterns of diversification for the California Floristic Province. *Molecular Ecology* 22:5418–5429.
- Myers, G.S. 1930. The status of the southern California toad, *Bufo californicus* Camp. *Proceedings of the Biological Society of Washington* 43:73–78.
- Myers, G.S. and Maslin, T.P., Jr. 1948. The California plethodont salamander, *Aneides flavipunctatus* (Strauch), with a description of a new subspecies and notes on other western *Aneides*. *Proceedings of the Biological Society of Washington* 61:127–138.
- Myers, N., Mittermeier, R.A., Mittermeier, C.G., da Fonseca, G.A.B., and Kent, J. 2000. Biodiversity hotspots for conservation priorities. *Nature* 403:853–858.
- Nagy, Z.T., Lawson, R., Joger, U., and Wink, M. 2004. Molecular systematics of racers, whip-

- snakes and relatives (Reptilia : Colubridae) using mitochondrial and nuclear markers. *Journal of Zoological Systematics and Evolutionary Research* 42:223–233.
- National Audubon Society. 2011. *The Christmas Bird Count Historical Results*. Available from <http://www.audubon.org>.
- Newbold, T.A.S. 2005. Desert horned lizard (*Phrynosoma platyrhinos*) locomotor performance: the influence of cheatgrass (*Bromus tectorum*). *The Southwestern Naturalist* 50:17–23.
- Newman, R.A. 1989. Developmental plasticity of *Scaphiopus couchii* tadpoles in an unpredictable environment. *Ecology* 70:1775–1787.
- Nielson, M., Lohman, K., Daugherty, C.H., Allendorf, F.W., Knudsen, K.L., and Sullivan, J. 2006. Allozyme and mitochondrial DNA variation in the tailed frog (Anura: *Ascaphus*): the influence of geography and gene flow. *Herpetologica* 62:235–258.
- Nielson, M., Lohman, K., and Sullivan, J. 2007. Phylogeography of the tailed frog (*Ascaphus truei*): implications for the biogeography of the Pacific Northwest. *Evolution* 55:147–160.
- Nieto, N.C. 2004. Prevalence of Chytridiomycosis in Red-Legged Frog (*Rana aurora aurora*) Larvae in and around Redwood National Park, Northern Coastal California, with Notes on Other Parasites. Master's Thesis. Humboldt State University, Arcata, CA.
- Noble, G.K. and Clausen, H.J. 1936. The aggregation behavior of *Storeria dekayi* and other snakes, with especial reference to the sense organs involved. *Ecological Monographs* 6:269–316.
- Norris, K.S. 1949. Observations on the habits of the horned lizard *Phrynosoma mcallii*. *Copeia* 1949:176–180.
- Norris, K.S. 1958. The evolution and systematics of the iguanid genus *Uma* and its relation to the evolution of other North American desert reptiles. *Bulletin of the American Museum of Natural History* 114:253–326.
- Northwest Forest Plan. 1994. Record of Decision for Amendments to Forest Service and Bureau of Land Management Planning Documents within the Range of the Northern Spotted Owl. USDA Forest Service and USDI Bureau of Land Management, Washington, DC, 74pp.
- Nussbaum, R.A. 1969. A nest site of the Olympic salamander, *Rhyacotriton olympicus* (Gaige). *Herpetologica* 25:277–278.
- Nussbaum, R.A. 1976. Geographic variation and systematics of salamanders of the genus *Dicamptodon* Strauch (Amphystomatidae). *Miscellaneous Publications of the Museum of Zoology* 149:1–94.
- Nussbaum, R.A., Brodie, E., and Storm, R. 1983. *Amphibians and Reptiles of the Pacific Northwest*. University of Idaho Press, Moscow, ID.
- Nussbaum, R.A. and Tait, C.K. 1977. Aspects of the life history and ecology of the Olympic salamander, *Rhyacotriton olympicus* (Gaige). *The American Midland Naturalist* 98:176–199.
- O'Donnell, R.P., Staniland, K., and Mason, R.T. 2007. Experimental evidence that oral secretions of northwestern ring-necked snakes (*Diadophis punctatus occidentalis*) are toxic to their prey. *Toxicon* 50:810–815.
- O'Hara, R.K. and Blaustein, A.R. 1981. An investigation of sibling recognition in *Rana cascadae* tadpoles. *Animal Behaviour* 29:1121–1126.
- O'Hara, R.K. and Blaustein, A.R. 1985. *Rana cascadae* tadpoles aggregate with siblings: an experimental field study. *Oecologia* 67:44–51.
- Ohmart, R.D., Anderson, B.W., and Hunter, W.C. 1988. The ecology of the lower Colorado River from Davis Dam to the Mexico-United States international boundary: a community profile. United States Fish and Wildlife Service Biological Report 85(7.19):1–296.
- Olson, D.H., Anderson, P.D., Frissell, C.A., Welsh, H.H., Jr., and Bradford, D.F. 2007. Biodiversity management approaches for stream-riparian areas: perspectives for Pacific Northwest headwater forests, microclimates, and amphibians. *Forest Ecology and Management* 246:81–107.
- Olson, D.H. and Burnett, K.M. 2009. Design and management of linkage areas across headwater drainages to conserve biodiversity in forest ecosystems. *Forest Ecology and Management* 258S:S117–S126.
- O'Regan, S.M., Palen, W.J., and Anderson, S.C. 2014. Climate warming mediates negative impacts of rapid pond drying for three amphibian species. *Ecology* 95:845–855.
- Ouren, D.S., Haas, C., Melcher, C.P., Stewart, S.C., Ponds, P.D., Sexton, N.R., Burris, L., Fancher, T., and Bowen, Z.H. 2007. Environmental effects of off-highway vehicles on Bureau of Land Management lands: A Literature Synthesis, Annotated Bibliographies, Extensive Bibliographies, and Internet Resources. Open-File Report 2007-1353. US Geological Survey, Reston, VA, pp.1–225.
- Ovaska, K., Hyatt, L., and Sopuck, L. 2002. Geographic distribution: *Rana aurora*. *Herpetological Review* 33:318.
- Pace, A. 1974. Systematic and biological studies of the leopard frogs (*Rana pipiens* complex) of the

- United States. Miscellaneous publications of the Museum of Zoology, University of Michigan 184:1-140.
- Packer, W.C. 1960. Bioclimatic influences on the breeding migration of *Taricha rivularis*. *Ecology* 41:509-517.
- Packer, W.C. 1961. Feeding behavior in adult *Taricha*. *Copeia* 1961:351-352.
- Packer, W.C. 1962. Aquatic homing behavior in *Taricha rivularis*. *Copeia* 1962:207-208.
- Packer, W.C. 1963. Observations on the breeding migration of *Taricha rivularis*. *Copeia* 1963:378-382.
- Padgett-Flohr, G.E. 2008. Pathogenicity of *Batrachochytrium dendrobatidis* in two threatened California amphibians: *Rana draytonii* and *Ambystoma californiense*. *Herpetological Conservation and Biology* 3:182-191.
- Padgett-Flohr, G.E. 2010. Landscape epidemiology of *Batrachochytrium dendrobatidis* in central California. *Ecography* 33:688-697.
- Padgett-Flohr, G.E. and Hayes, M.P. 2011. Assessment of the vulnerability of the Oregon spotted frog (*Rana pretiosa*) to the amphibian chytrid fungus (*Batrachochytrium dendrobatidis*). *Herpetological Conservation and Biology* 6:99-106.
- Padgett-Flohr, G.E. and Hopkins, R.L. 2009. *Batrachochytrium dendrobatidis*, a novel pathogen approaching endemism in central California. *Diseases of Aquatic Organisms* 83:1-9.
- Padgett-Flohr, G.E. and Longcore, J.E. 2007. Natural history notes: *Taricha torosa*. Fungal infection. *Herpetological Review* 78:176-177.
- Palen, W.J. and Schindler, D.E. 2010. Water clarity, maternal behavior, and physiology combine to eliminate UV radiation risk to amphibians in a montane landscape. *Proceedings of the National Academy of Sciences of the United States of America* 107:9701-9706.
- Palen, W.J., Schindler, D.E., Adams, M.J., Pearl, C.A., Bury, R.B., and Diamond, S.A. 2002. Optical characteristics of natural waters protect amphibians from UV-B in the U.S. Pacific northwest. *Ecology* 83:2951-2957.
- Papenfuss, T.J. and Macey, J.R. 1986. A Review of the Population Status of the Inyo Mountains Salamander (*Batrachoseps campi*). Final Report under Order 10188-5671-5 for the United States Fish and Wildlife Service. Endangered Species Office, Sacramento, CA.
- Papenfuss, T.J. and Parham, J.F. 2013. Four new species of California legless lizards (*Anniella*). *Breviora* 536:1-17.
- Parham, J. and Papenfuss, T.J. 2009. High genetic diversity among fossorial lizard populations (*Anniella pulchra*) in a rapidly developing landscape (Central California). *Conservation Genetics* 10:169-176.
- Parisi, M. (Ed.). 2003. Atlas of the Biodiversity of California. California Department of Fish and Game, Sacramento.
- Parker, M.S. 1994. Feeding ecology of stream-dwelling Pacific giant salamander larvae (*Dicamptodon tenebrosus*). *Copeia* 1994:705-718.
- Parker, W.S. and Brown, W.S. 1974. Notes on the ecology of regal ringneck snakes (*Diadophis punctatus regalis*) in northern Utah. *Journal of Herpetology* 8:262-263.
- Parmesan, C. 2006. Ecological and evolutionary responses to recent climate change. *Annual Review of Ecology, Evolution and Systematics* 37:637-669.
- Parmesan, C. and Yohe, G. 2003. A globally coherent fingerprint of climate change impacts across natural systems. *Nature* 421:37-42.
- Patten, M.A. and Myers, S.J. 1992. Geographic distribution: *Bufo californicus*. *Herpetological Review* 23:122.
- Patten, R.B. and Banta, B.H. 1980. A rattlesnake, *Crotalus ruber*, feeds on a road-killed animal. *Journal of Herpetology* 14:111-112.
- Pauly, G.B., Hillis, D.M., and Cannatella, D.C. 2004. The history of a nearctic colonization: molecular phylogenetics and biogeography of the nearctic toads. *Evolution* 58:2517-2535.
- Pauly, G.B., Hillis, D.M., and Cannatella, D.C. 2009. Taxonomic freedom and the role of official lists of species names. *Herpetologica* 65:115-128.
- Pearl, C.A. 2005. *Rana aurora*, in Lannoo, M.J. (Ed.): *Amphibian Declines: The Conservation Status of United States Species*. University of California Press, Berkeley, pp.58-530.
- Pearl, C.A. and Adams, M.J. 2005. *Rana cascadae*, in Lannoo, M.J. (Ed.): *Amphibian Declines: The Conservation Status of United States Species*. University of California Press, Berkeley, pp. 538-540.
- Pearl, C.A., Adams, M.J., Bury, R.B., and McCreary, B. 2004. Asymmetrical effects of introduced bullfrogs (*Rana catesbeiana*) on native ranid frogs in Oregon. *Copeia* 2004:11-20.
- Pearl, C.A., Adams, M.J., and Leuthold, N. 2009. Breeding habitat and local population size of the Oregon spotted frog (*Rana pretiosa*) in Oregon, U.S.A. *Northwestern Naturalist* 90:136-147.
- Pearl, C.A., Adams, M.J., Leuthold, N., and Bury, R. 2005a. Amphibian occurrence and aquatic invaders in a changing landscape: implications for wetland mitigation in the Willamette Valley, Oregon, USA. *Wetlands* 25:76-88.

- Pearl, C.A., Bowerman, J. and Knight, D. 2005b. Feeding behavior and aquatic habitat use by Oregon spotted frogs (*Rana pretiosa*) in central Oregon. *Northwestern Naturalist* 86:36–38.
- Pearl, C.A., Bull, E.L., Green, D.E., Bowerman, J., Adams, M.J., Hyatt, A., and Wente, W.H. 2007. Occurrence of the amphibian pathogen *Batrachochytrium dendrobatidis* in the Pacific Northwest. *Journal of Herpetology* 41:145–149.
- Pearl, C.A. and Hayes, M.P. 2002. Predation by Oregon spotted frogs (*Rana pretiosa*) on western toads (*Bufo boreas*) in Oregon. *The American Midland Naturalist* 147:145–152.
- Pearl, C.A. and Hayes, M.P. 2005. *Rana pretiosa*. In Lannoo, M.J. (Ed.): *Amphibian Declines: The Conservation Status of United States Species*. University of California Press, Berkeley, pp.577–580.
- Pearl, C.A., Hayes, M.P., Haycock, R., Engler, J.D., and Bowerman, J. 2005c. Observations of interspecific amplexus between western North American ranid frogs and the introduced American bullfrog (*Rana catesbeiana*) and an hypothesis concerning breeding interference. *The American Midland Naturalist* 154:126–134.
- Pearse, D. and Pogson, G. 2000. Parallel evolution of the melanistic form of the California legless lizard, *Anniella pulchra*, inferred from mitochondrial DNA sequence variation. *Evolution* 54:1041–1046.
- Peek, R. 2010. Landscape Genetics of Foothill Yellow-Legged Frogs (*Rana boylei*) in Regulated and Unregulated Rivers: Assessing Connectivity and Genetic Fragmentation. Master's Thesis. University of San Francisco, San Francisco, CA.
- Perkins, C.B. 1938. The snakes of San Diego County with descriptions and key. *Bulletin of the Zoological Society of San Diego* 13:1–66.
- Perry, G., Buchanan, B.W., Fisher, R.N., Salmon, M., and Wise, S.E. 2008. Effects of artificial night lighting on amphibians and reptiles in urban environments, in Mitchell, J.C., Brown, J., and Bartholomew, B. (Eds): *Urban Herpetology*. Society for the Study of Amphibians and Reptiles. Salt Lake City, UT, pp.239–256.
- Perry, G. and Fisher, R.N. 2006. Night lights and reptiles: observed and potential effects, in Rich, C. and Longcore, T. (Eds): *Ecological Consequences of Artificial Night Lighting*. Island Press, Washington, DC, pp.169–191.
- Petranka, J.W. 1958. *Salamanders of the United States and Canada*. Smithsonian Institution Press, Washington, DC.
- Pfennig, D. 1990. The adaptive significance of an environmentally-cued developmental switch in an anuran tadpole. *Oecologia* 85:101–107.
- Phillipsen, I.C., Funk, W.C., Hoffman, E.A., Monsen, K.J., and Blouin, M.S. 2011. Comparative analyses of effective population size within and among species: ranid frogs as a case study. *Evolution* 65:2927–2945.
- Pianka, E.R. and Parker, W.S. 1975. Ecology of horned lizards: a review with special reference to *Phrynosoma platyrhinos*. *Copeia* 1975:141–162.
- Picco, A.M., Brunner, J.L., and Collins, J.P. 2007. Susceptibility of the endangered California tiger salamander, *Ambystoma californiense*, to Ranavirus infection. *Journal of Wildlife Diseases* 43:286–290.
- Pilliod, D.S., Bury, R.B., Hyde, E.J., Pearl, C.A., and Corn, P.S. 2003. Fire and amphibians in North America. *Forest Ecology and Management* 178:163–181.
- Pilliod, D.S. and Fronzuto, J.A. 2005. *Ambystoma macrodactylum*, in Lannoo, M.J. (Ed.): *Amphibian Declines: The Conservation Status of United States Species*. University of California Press, Berkeley, pp.617–621.
- Pilliod, D.S., and Peterson, C.R. 2001. Local and landscape effects of introduced trout on amphibians in historically fishless watersheds. *Ecosystems* 4:322–333.
- Piovia-Scott, J., Pope, K.L., Lawler, S.P., Cole, E.M., and Foley, J.E. 2011. Factors related to the distribution and prevalence of the fungal pathogen *Batrachochytrium dendrobatidis* in *Rana cascadae* and other amphibians in the Klamath Mountains. *Biological Conservation* 144:2913–2921.
- Piry, S., Luikart, G., and Cornuet, J.M. 1999. BOTTLENECK: a computer program for detecting recent reductions in the effective size using allele frequency data. *Journal of Heredity* 90:502–503.
- Platz, J.E. 1976. Biochemical and morphological variation of leopard frogs in Arizona. *Copeia* 1976:660–672.
- Platz, J.E. 1988. *Rana yavapaiensis*. *Catalogue of American Amphibians and Reptiles* 418:1–2.
- Platz, J.E., Clarkson, R.W., Rorabaugh, J.C., and Hillis, D.M. 1990. *Rana berlandieri*: recently introduced populations in Arizona and southeastern California. *Copeia* 1990:324–333.
- Platz, J.E. and Frost, D. 1984. *Rana yavapaiensis*, a new species of leopard frog (*Rana pipiens* complex). *Copeia* 1984:940–948.
- Platz, J.E. and Mecham, J.S. 1979. *Rana chiricahuensis*, a new species of leopard frog (*Rana*

- pipiens complex) from Arizona. *Copeia* 1979:383–390.
- Platz, J.E. and Platz, A.L. 1973. *Rana pipiens* complex: hemoglobin phenotypes of sympatric and allopatric populations in Arizona. *Science* 179:1334–1336.
- Pollett, K.L., MacCracken, J.G. and MacMahon, J.A. 2010. Stream buffers ameliorate the effects of timber harvest on amphibians in the Cascade Range of Southern Washington, USA. *Forest Ecology and Management* 260:1083–1087.
- Pomeroy, L.V. 1981. Developmental Polymorphism in the Tadpoles of the Spadefoot Toad *Scaphiopus multiplicatus*. PhD Dissertation. University of California, Riverside.
- Pope, C.H. 1939. *Turtles of the United States and Canada*. AA Knopf, New York.
- Pope, K., Brown, C., Hayes, M., Green, G., and Macfarlane, D. 2014. Cascades Frog Conservation Assessment. United States Department of Agriculture, Forest Service, General Technical Report PSW-GTR-244 32. 116pp.
- Pope, K.L. 2008. Assessing changes in amphibian population dynamics following experimental manipulations of introduced fish. *Conservation Biology* 22:1572–1581.
- Pope, K.L., Garwood, J.M., Welsh, H.H., Jr., and Lawler, S.P. 2008. Evidence of indirect impacts of introduced trout on native amphibians via facilitation of a shared predator. *Biological Conservation* 141:1321–1331.
- Pope, K.L. and Larson, M.D. 2013. Status of the Cascades Frog (*Rana cascadae*) in the Southern Cascade Mountains of California. Final Report to the United States Forest Service for the years 2008–2012. ISA #12-05-06-02.
- Pope, K.L., Larson, M.D., and Piovia-Scott, J. 2011. Status of Remnant Populations of Cascades Frogs (*Rana cascadae*) in the Lassen Area of California. Draft Final Report to the United States Forest Service, Lassen National Forest.
- Pope, S.E., Fahrig, L. and Merriam, H.G. 2000. Landscape complementation and metapopulation effects on leopard frog populations. *Ecology* 81:2498–2508.
- Pough, F.H. 1974. *Uma scoparia*. *Catalogue of American Amphibians and Reptiles* 155:1–2.
- Pounds, J.A., Bustamante, M.R., Coloma, L.A., Consuegra, J.A., Fogden, M.P.L., Foster, P.N., La Marca, E. et al. 2006. Widespread amphibian extinctions from epidemic disease driven by global warming. *Nature* 439:161–167.
- PRBO (PRBO Conservation Science). 2011. Projected effects of climate change in California: ecoregional summaries emphasizing consequences for wildlife. Version 1.0. Accessed January 30, 2012. <http://data.prbo.org/apps/fbssc/climatechange>.
- Price, A.H. and Sullivan, B.K. 1988. *Bufo microscaphus*. *Catalogue of American Amphibians and Reptiles* 415:1–3.
- Prugh, L.R., Hodges, K.E., Sinclair, A.R.E., and Brashares, J.S. 2008. Effect of habitat area and isolation on fragmented animal populations. *Proceedings of the National Academy of Sciences of the United States of America* 105:20770–20775.
- Pyke, C.R. 2005. Interactions between habitat loss and climate change: implications for fairy shrimp in the Central Valley ecoregion of California, U.S.A. *Climatic Change* 68:199–218.
- Pyron, R.A. and Burbrink, F.T. 2009. Neogene diversification and taxonomic stability in the snake tribe Lamproleptini (Serpentes: Colubridae). *Molecular Phylogenetics and Evolution* 52:524–529.
- Rainey, W.E. 1985. Genetic Variation in *Anniella nigra*. Attachment in Bury. Status Report, *Anniella nigra*, in Central California. Office of Endangered Species, United States Fish and Wildlife Service, Portland, OR.
- Rathbun, G.B., Jennings, M.R., Murphey, T.G., and Sicpel, N.R. 1993. Status and Ecology of Sensitive Aquatic Vertebrates in Lower San Simeon and Pico Creeks, San Luis Obispo County, California. Final Report under Cooperative Agreement 14-16-0009-91-1909 between U.S. Fish and Wildlife Service and California Department of Parks and Recreation. Publication No. PB93-230779, National Technical Information Service, Springfield, VA.
- Rathke, M.H. 1833. *Triton torosus*, in *Zoologischer Atlas, enthaltend Abbildungen und Beschreibungen neuer Thierarten, während des Flottcapitains von Kotzebue zweiter Reise um die Welt, auf Russisch-Kaiserlich Kriegsschupp Predpriaetië in den Jahren 1823-1826 herausgegeben von D. Martin Heinrich Rathke. Fünftes Heft*. G. Reimer, Berlin, pp.12–14. (In German)
- Ray, C. 1958. Vital limits and rates of desiccation in salamanders. *Ecology* 39:75–83.
- Recuero, R., Martinez-Solano, I., Parra-Olea, G., and Garcia-Paris, M. 2006a. Phylogeography of *Pseudacris regilla* (Anura: Hylidae) in western North America, with a proposal for a new taxonomic rearrangement. *Molecular Phylogenetics and Evolution* 39:293–304.
- Recuero, R., Martinez-Solano, I., Parra-Olea, G., and Garcia-Paris, M. 2006b. Corrigendum to “Phylogeography of *Pseudacris regilla* (Anura:

- Hylidae) in western North America, with a proposal for a new taxonomic rearrangement" [Mol. Phylogenet. Evol. 39 (2006) 293–304]. Molecular Phylogenetics and Evolution 41:511.
- Reeder, T.W., Cole, C.J., and Dessauer, H.C. 2002. Phylogenetic relationships of whiptail lizards of the genus *Cnemidophorus* (Squamata: Teiidae): a test of monophyly, reevaluation of karyotypic evolution, and review of hybrid origins. American Museum Novitates 3365:1–61.
- Reese, D.A. 1996. Comparative Demography and Habitat Use of Western Pond Turtles in Northern California: The Effects of Damming and Related Alterations. PhD Dissertation. University of California, Berkeley.
- Reese, D.A. and Welsh, H.H., Jr. 1997. Use of terrestrial habitat by western pond turtles, *Clemmys marmorata*: implications for management. in Proceedings of an International Conference on Conservation, Restoration, and Management of Tortoises and Turtles. State University of New York, Purchase, pp.352–357.
- Reese, D.A. and Welsh, H.H., Jr. 1998. Comparative demography of *Clemmys marmorata* populations in the Trinity River of California in the context of dam-induced alterations. Journal of Herpetology 32:505–515.
- Reeve, W.L. 1952. Taxonomy and distribution of the horned lizard genus *Phrynosoma* (family Iguanidae). University of Kansas Science Bulletin 34:817–960.
- Reilly, S.B., Portik, D.M., Koo, M.S., and Wake, D.B. In Press. Discovery of a new, disjunct population of a narrowly distributed salamander (*Taricha rivularis*) in California presents conservation challenges. Journal of Herpetology 48.
- Relyea, R.A. 2005a. The lethal impacts of Roundup and predatory stress on six species of North American tadpoles. Archives of Environmental Contamination and Toxicology 48:351–357.
- Relyea, R.A. 2005b. The lethal impact of Roundup on aquatic and terrestrial amphibians. Ecological Applications 15:1118–1124.
- Relyea, R.A. 2008. A cocktail of contaminants: how mixtures of pesticides at low concentrations affect aquatic communities. Oecologia 159:363–376.
- Relyea, R.A. and Jones, D.K. 2009. The toxicity of Roundup Original MAX(R) to thirteen species of larval amphibians. Environmental Toxicology and Chemistry 28:2004–2008.
- Remsen, J.V., Jr. 1978. Bird species of special concern in California. California Department of Fish and Game, Wildlife Management Branch Report 78(1):1–63pp.
- Reynolds, F.A. 1943. Notes on the western glossy snake in captivity. Copeia 1943:196.
- Rhodin, A.G.J., van Dijk, P.P., Iverson, J.B., and Shaffer, H.B. 2010. Turtles of the world, 2010 update: annotated checklist of taxonomy, synonymy, distribution, and conservation status. Chelonian Research Monographs 5:85–164.
- Richter, K.O. and Azous, A.L. 1995. Amphibian occurrence and wetland characteristics in the Puget Sound Basin. Wetlands 15:305–312.
- Rieder, J.P., Newbold, T.A.S., and Ostojka, S.M. 2010. Structural changes in vegetation coincident with annual grass invasion negatively impacts sprint velocity of small vertebrates. Biological Invasions 12:2429–2439.
- Riemer, W.J. 1958. Variation and systematic relationships within the salamander genus *Taricha*. University of California Publications in Zoology 56:301–390.
- Riley, S.P.D., Busteed, G., Kats, L., Vandergon, T., Lee, L., Dagit, R., Kerby, J., Fisher, R., and Sauvajot, R. 2005. Effects of urbanization on the distribution and abundance of amphibians and invasive species in southern California streams. Conservation Biology 19:1894–1907.
- Riley, S.P.D., Shaffer, H.B., Voss, S.R. and Fitzpatrick, B.M. 2003. Hybridization between a rare, native tiger salamander (*Ambystoma californiense*) and its introduced congener. Ecological Applications 13:1263–1275.
- Rissler, L.J. and Apodaca, J.J. 2007. Adding more ecology into species delimitation: ecological niche models and phylogeography help define cryptic species in the black salamander (*Aneides flavipunctatus*). Systematic Biology 56:924–942.
- Ritter, W.E. 1897. The life-history and habits of the Pacific coast newt (*Diemyctylus torosus* Esch.). Proceedings of the California Academy of Sciences, 3rd Series 1:73–114.
- Rochester, C.J., Brehme, C.S., Clark, D.R., Stokes, D.C., Hathaway, S.A., and Fisher, R.N. 2010. Reptile and amphibian responses to large-scale wildfires in southern California. Journal of Herpetology 44:333–351.
- Rodríguez-Robles, J.A., Bell, C.J., and Greene, H.W. 1999a. Food habits of the glossy snake, *Arizona elegans*, with comparisons to the diet of sympatric long-nosed snakes, *Rhincheilus lecontei*. Journal of Herpetology 33:87–92.
- Rodríguez-Robles, J.A., DeNardo, D.F., and Staub, R.E. 1999b. Phylogeography of the California mountain kingsnake, *Lampropeltis zonata* (Colubridae). Molecular Ecology 8:1923–1934.

- Rodríguez-Robles, J.A. and Galina-Tessaro P. 2006. Natural history notes: *Thamnophis hammondi*. Diet. Herpetological Review 37:355.
- Roelants, K., Gower, D.J., Wilkinson, M., Loader, S.P., Biju, S.D., Guillaume, K., Moriau, L., and Bossuyt, F. 2007. Global patterns of diversification in the history of modern amphibians. Proceedings of the National Academy of Sciences, USA 104:887–892.
- Rohr, J.R., Schotthoefer, A.M., Raffel, T.R., Garrick, H.J., Halstead, N., Hoverman, J.T., Johnson, C.M. et al. 2008. Agrochemicals increase trematode infections in a declining amphibian species. Nature 455:1235–1239.
- Romansic, J.M., Diez, K.A., Higashi, E.M., Johnson, J.E., and Blaustein, A.R. 2009a. Effects of the pathogenic water mold *Saprolegnia ferax* on survival of amphibian larvae. Diseases of Aquatic Organisms 83:187–193.
- Romansic, J.M., Higashi, E.M., Diez, K.A., and Blaustein, A.R. 2007. Susceptibility of newly-metamorphosed frogs to a pathogenic water mould (*Saprolegnia* sp.). Herpetological Journal 17:161–166.
- Romansic, J.M., Waggener, A.A., Bancroft, B.A., and Blaustein, A.R. 2009b. Influence of ultraviolet-B radiation on growth, prevalence of deformities, and susceptibility to predation in Cascades frog (*Rana cascadae*) larvae. Hydrobiologia 624:219–233.
- Roos, M. 1992. The Hydrology of the 1987–1992 California Drought. Technical Paper. State of California Resources Agency, Department of Water Resources, Division of Flood Management.
- Root, T.L., MacMynowski, D.P., Mastrandrea, M.D., and Schneider, S.H. 2005. Human-modified temperatures induce species changes: joint attribution. Proceedings of the National Academy of Sciences of the United States of America 102:7465–7469.
- Root, T.L., Price, J.T., Hall, K.R., Schneider, S.H., Rosenzweig, C., and Pounds, J.A. 2003. Fingerprints of global warming on wild animals and plants. Nature 421:57–60.
- Rorabaugh, J.C. 2005. *Rana pipiens*, in Lannoo, M.J. (Ed.): Amphibian Declines: The Conservation Status of United States Species. University of California Press, Berkeley, pp.570–577.
- Rorabaugh, J.C., Palermo, C.L., and Dunn, S.C. 1987. Distribution and relative abundance of the flat-tailed horned lizard (*Phrynosoma mcallii*) in Arizona. The Southwestern Naturalist 32:103–109.
- Rorabaugh, J.C., Sredl, M.J., Miera, V., and Drost, C.A. 2002. Continued invasion by an introduced frog (*Rana berlandieri*): southwestern Arizona, southeastern California, and Rio Colorado, Mexico. The Southwestern Naturalist 47:12–20.
- Rorabaugh, J.C. and Young, K.V. 2009. Flat-tailed horned lizard, *Phrynosoma mcallii* (Hallowell, 1852), in Jones, L.L.C. and Lovich, R.E. (Eds.): Lizards of the American Southwest: A Photographic Field Guide. Rio Nuevo Publishers, Tucson, AZ, pp.182–185.
- Rossman, D.A. 1979. Morphological evidence for taxonomic partitioning of the *Thamnophis elegans* complex (Serpentes, Colubridae). Occasional Papers of the Museum of Zoology, Louisiana State University 55:1–12.
- Rossman, D.A., Ford, N.B., and Seigel, R.A. 1996. The Garter Snakes: Evolution and Ecology. University of Oklahoma Press, Norman.
- Rossman, D.A. and Stewart, G.R. 1987. Taxonomic reevaluation of *Thamnophis couchii* (Serpentes: Colubridae). Occasional Papers of the Museum of Zoology, Louisiana State University 63:1–25.
- Rovito, S.M. 2010. Lineage divergence and speciation in the web-toed salamanders (Plethodontidae: *Hydromantes*) of the Sierra Nevada, California. Molecular Ecology 19:4554–4571.
- Rowe, J.C. and Garcia, T.S. 2013. Impacts of wetland restoration efforts on an amphibian assemblage in a multi-invader community. Wetlands 34:141–153.
- Royle, J.A. and Young, K.V. 2008. A hierarchical model for spatial capture-recapture data. Ecology 89:2281–2289.
- Ruibal, R. 1959. The ecology of a brackish water population of *Rana pipiens*. Copeia 1959:315–322.
- Rundio, D.E. and Olson, D.H. 2001. Palatability of southern torrent salamander (*Rhyacotriton variegatus*) larvae to Pacific giant salamander (*Dicamptodon tenebrosus*) larvae. Journal of Herpetology 35:133–136.
- Ruppert, R.M. 2010a. A recent sighting of the banded Gila monster, (*Heloderma suspectum cinctum*) in Mojave National Preserve, California. Mojave National Preserve Science Newsletter 2010:1–3.
- Ruppert, R.M. 2010b. Geographic distribution: *Heloderma suspectum cinctum*. Herpetological Review 41:107.
- Russell, R.W. and Anderson, J.D. 1956. A disjunct population of the long-nosed salamander from the coast of California. Herpetologica 12:137–140.
- Rüthling, P.D. 1915. Hibernation of reptiles in southern California. Copeia 1915:10–11.
- Ruthven, A.G. 1907. A collection of reptiles and amphibians from southern New Mexico and

- Arizona. Bulletin of the American Museum of Natural History 23:483–604.
- Ryan, M.E., Johnson, J.R., and Fitzpatrick, B.M. 2009. Invasive hybrid tiger salamander genotypes impact native amphibians. Proceedings of the National Academy of Sciences of the United States of America 106:11166–11171.
- Salt, G.S. 1952. The bell toad, *Ascaphus truei*, in Mendocino County, California. Copeia 1952:193–194.
- Sartorius, S.S. and Rosen, P.C. 2000. Breeding phenology of the lowland leopard frog (*Rana yavapaiensis*): implications for conservation and ecology. The Southwestern Naturalist 45:267–273.
- Sattler, P.W. 1980. Genetic relationships among selected species of North American *Scaphiopus*. Copeia. 1980:605–610.
- Sauer, J.R., Hines, J.E., Fallon, J.E., Pardieck, K.L., Ziolkowski, D.J., Jr., and Link, W.A. 2011. The North American Breeding Bird Survey, results and analysis 1966–2009. Version 3.23. 2011 USGS Patuxent Wildlife Research Center, Laurel, MD.
- Savage, W.K. 2008. Landscape Genetics and Phylogenetic Relationships in the *Ambystoma macrodactylum* species complex. PhD Dissertation. University of California, Davis.
- Savage, W.K., Premier, A.K., and Shaffer, H.B. 2010. Landscape genetics of alpine Sierra Nevada salamanders reveal extreme population subdivision in space and time. Molecular Ecology 19:3301–3314.
- Schloegel, L.M., Picco, A.M., Kilpatrick, A.M., Davies, A.J., Hyatt, A.D., and Daszak, D. 2009. Magnitude of the U.S. trade in amphibians and presence of *Batrachochytrium dendrobatidis* and Ranavirus infection in imported North American bullfrogs (*Rana catesbeiana*). Biological Conservation 142:1420–1426.
- Schmidt, B.R. and Zumbach, S. 2008. Amphibian road mortality and how to prevent it: a review, in Mitchell, J.C., Jung Brown, R.E. and Bartholomew, B. (Eds.): Urban Herpetology. Society for the Study of Amphibians and Reptiles, Salt Lake City, UT, pp.157–167.
- Schmieder, R.R. and Nauman, R.S. 1994. Effects of Non-Native Aquatic Predators on Premetamorphic California Red-Legged Frogs (*Rana aurora draytonii*). University of California, Santa Cruz, 12pp.
- Schoenherr, A.A. 1976. The Herpetofauna of the San Gabriel Mountains, Los Angeles County, California, including Distribution and Biogeography. Special Publication No. 1. Southwestern Herpetologists Society, Van Nuys, CA.
- Schreber, H. 1782. Der Naturforscher. (Johann Jacob Gebaur, Halle. (In German)
- Schulte, J.A., Macey, J.R., and Papenfuss, T.J. 2006. A genetic perspective on the geographic association of taxa among arid North American lizards of the *Sceloporus magister* (Squamata: Iguanidae: Phrynosomatidae). Molecular Phylogenetics and Evolution 39:873–880.
- Scott, N.J., Rathbun, G.B., Murphey, T.G., and Harker, M.B. 2008. Reproduction of Pacific pond turtles (*Actinemys marmorata*) in coastal streams of central California. Herpetological Conservation and Biology 3:143–148.
- Searcy, C.A. and Shaffer, H.B. 2008. Calculating biologically accurate mitigation credits: insights from the California tiger salamander. Conservation Biology 22:997–1005.
- Searcy, C.A. and Shaffer, H.B. 2011. Determining the migration distance of a vagile vernal pool specialist: how much land is required for conservation of California tiger salamanders?, in Alexander, D.G. and Schlising, R.A. (Eds). Research and Recovery in Vernal Pool Landscapes, Studies from the Herbarium, No. 16. California State University, Chico. pp.73–87.
- Secor, S.M. 1995. Ecological aspects of foraging mode for the snakes *Crotalus cerastes* and *Masticophis flagellum*. Herpetological Monographs 9:169–186.
- Seeliger, L.M. 1945. Variation in the Pacific mud turtle. Copeia 1945:150–159.
- Setser, K. 2004. Natural History, Demography, and Home Range Characteristics of a Southern California Population of *Phrynosoma mcallii* Inhabiting Atypical Habitat. Master's Thesis. Utah State University, Logan.
- Sever, D.M., Moriarty, E.C., Rania, L.C., and Hamlett, W.C. 2001. Sperm storage in the oviduct of the internal fertilizing frog *Ascaphus truei*. Journal of Morphology 248:1–21.
- Shaffer, H.B., Cook, D., Fitzpatrick, B.M., Leyse, K., Picco, A., and Trenham, P.C. 2009. Guidelines for the relocation of California tiger salamanders (*Ambystoma californiense*). Report. United States Fish and Wildlife Service, Sacramento, CA.
- Shaffer, H.B., Fellers, G.M., Magee, A., and Voss, S.R. 2000. The genetics of amphibian declines: population substructure and molecular differentiation in the Yosemite Toad, *Bufo canorus* (Anura, Bufonidae) based on single-strand conformation polymorphism analysis (SSCP) and mitochondrial DNA sequence data. Molecular Ecology 9:245–257.
- Shaffer, H.B., Fellers, G.M., Randal, V., Oliver, J.C., and Pauly, G.B. 2004. Species boundaries,

- phylogeography and conservation genetics of the red-legged frog (*Rana aurora/draytonii*) complex. *Molecular Ecology* 13:2667–2677.
- Shedd, J., Bogiatto, R.J. and Kirn, S.A. 2011. Natural history notes: *Phrynosoma blainvillii*. *Commenalism. Herpetological Review* 42:94–95
- Shoemaker, V.H., McClanahan, L., and Ruibal, R. 1969. Seasonal changes in body fluids in a field population of spadefoot toads. *Copeia* 1969:585–591.
- Shuford, W.D. and Gardali, T. (Eds). 2008. *California Bird Species of Special Concern: A Ranked Assessment of Species, Subspecies, and Distinct Populations of Birds of Immediate Conservation Concern in California*. *Studies of Western Birds 1*. Western Field Ornithologists, Camarillo, CA, and California Department of Fish and Game, Sacramento.
- Sinclair, E.A., Bezy, R.L., Bolles, K., Camarillo, R.J.L., Crandall, K.A., and Sites, J. 2004. Testing species boundaries in an ancient species complex with deep phylogeographic history: genus *Xantusia* (Squamata: Xantusiidae). *The American Naturalist* 164:396–414.
- Slater, J.R. 1939. Description and life-history of a new *Rana* from Washington. *Herpetologica* 1:145–149.
- Slevin, J.R. 1928. The amphibians of western North America: an account of the species known to inhabit California, Alaska, British Columbia, Washington, Oregon, Idaho, Utah, Nevada, Arizona, Sonora, and Lower California. *Occasional Papers of the California Academy of Sciences* 16:1–152
- Smith, H.M. 1946. *Handbook of Lizards: Lizards of the United States and Canada*. Comstock Publishing, Ithaca, NY.
- Smith, H.M., Brown, L.E., Chiszar, D., Grismer, L.L., Allen, G.S., Fishbein, A., Hollingsworth, B.D. et al. 1998. *Crotalus ruber* Cope, 1892 (Reptilia, Serpentes): proposed precedence of the specific name over that of *Crotalus exsul* Garman, 1884. *Bulletin of Zoological Nomenclature* 55:229–232.
- Sneed, D. 14 May 2009. Cuesta group makes monster find on field trip. *The Tribune*, San Luis Obispo, B1.
- Snyder, M.A. and Sloan, L.C. 2005. Transient future climate over the western United States using a regional climate model. *Earth Interactions* 9:1–21
- Snyder, M.A., Sloan, L.C., and Bell, J.L. 2004. Modeled regional climate change in the hydrologic regions of California: a CO₂ sensitivity study. *Journal of the American Water Resources Association* 40:591–601.
- Snyder, M.A., Sloan, L.C., Dillenbaugh, N.S., and Bell, J.L. 2003. Future climate change and upwelling in the California Current. *Geophysical Research Letters* 30:1–4
- Sparling, D.W., Fellers, G.M., and McConnell, L.L. 2001. Pesticides and amphibian population declines in California, U.S.A. *Environmental Toxicology and Chemistry* 20:1591–1595.
- Spear, S.F. and Storfer, A. 2008. Landscape genetic structure of coastal tailed frogs (*Ascaphus truei*) in protected vs. managed forests. *Molecular Ecology* 17:4642–4656.
- Spear, S.F. and Storfer, A. 2010. Anthropogenic and natural disturbance lead to differing patterns of gene flow in the Rocky Mountain tailed frog, *Ascaphus montanus*. *Biological Conservation* 143:778–786.
- Spencer, W.D., Beier, P., Penrod, K., Winters, K., Paulman, C., Rustigian-Romsos, H., Stritholt, J., Parisi, M., and Pettler, A. 2010. *California Essential Habitat Connectivity Project: A Strategy for Conserving a Connected California*. Prepared for California Department of Transportation, California Department of Fish and Game, and Federal Highways Administration.
- Spinks, P.Q., Pauly, G.B., Crayon, J.J., and Shaffer, H.B. 2003. Survival of the western pond turtle (*Emys marmorata*) in an urban California environment. *Biological Conservation* 113:257–267.
- Spinks, P.Q. and Shaffer, H.B. 2005. Range-wide molecular analysis of the western pond turtle (*Emys marmorata*): cryptic variation, isolation by distance, and their conservation implications. *Molecular Ecology* 14:2047–2064.
- Spinks, P.Q. and Shaffer, H.B. 2009. Conflicting mitochondrial and nuclear phylogenies for the widely disjunct *Emys* (Testudines: Emydidae) species complex, and what they tell us about biogeography and hybridization. *Systematic Biology* 58:1–20.
- Spinks, P.Q., Thomson, R.C., Lovely, G.A., and Shaffer, H.B. 2009. Assessing what is needed to resolve a molecular phylogeny: simulations and empirical data from emydid turtles. *BMC Evolutionary Biology* 9:56
- Spinks, P.Q., Thomson, R.C., and Shaffer, H.B. 2010. Nuclear gene phylogeography reveals the historical legacy of an ancient inland sea on lineages of the western pond turtle, *Emys marmorata* in California. *Molecular Ecology* 19:542–56.
- Spinks, P.Q., Thomson, R.C., and Shaffer, H.B. 2014. The advantages of going large: genome-

- wide SNPs clarify the complex population history and systematics of the threatened western pond turtle. *Molecular Ecology* 23:2228–2241
- Sredl, M.J., Collins, E.P., and Howland, J.M. 1997. Mark-Recapture Studies of Arizona Leopard Frogs. Arizona Game and Fish Department, Phoenix.
- Stanila, B.D. 2009. Morphology and Demography of Sonoran Mud Turtles (*Kinosternon sonoriense*) along an Aquatic Habitat Permanence Gradient. Master's Thesis. University of Central Oklahoma, Edmond.
- Staub, N. and Wake, D. 2005. *Aneides flavipunctatus*. in Lannoo, M.J. (Ed.): Amphibian Declines: The Conservation Status of United States Species. University of California Press, Berkeley, pp.660–661.
- Staub, R.E. and Mulks, M.F. 2009. Preliminary Assessment of Habitat Damage in the Laguna Mountains and Impacts on the San Diego Mountain Kingsnake. Unpublished report.
- Stebbins, R.C. 1943. Adaptations in the nasal passages for sand burrowing in the saurian genus *Uma*. *The American Naturalist* 77: 38–52.
- Stebbins, R.C. 1944. Some aspects of the ecology of the Iguanid genus *Uma*. *Ecological Monographs* 14:311–332.
- Stebbins, R.C. 1948. Nasal structure in lizards with reference to olfaction and conditioning of the inspired air. *American Journal of Anatomy* 83:183–221.
- Stebbins, R.C. 1951. Amphibians of Western North America. University of California Press, Berkeley.
- Stebbins, R.C. 1954. Amphibians and Reptiles of Western North America. McGraw-Hill Book Company, New York.
- Stebbins, R.C. 1955. Southern occurrence of the Olympic salamander, *Rhyacotriton olympicus*. *Herpetologica* 11:238–239.
- Stebbins, R.C. 1958. A new alligator lizard from the Panamint Mountains, Inyo County, California. *American Museum Novitates* 1883:27.
- Stebbins, R.C. 1959. Reptiles and Amphibians of the San Francisco Bay Region. University of California Press, Berkeley.
- Stebbins, R.C. 1966. Field Guide to Western Reptiles and Amphibians. Houghton Mifflin Company, Boston, MA.
- Stebbins, R.C. 1972. California Amphibians and Reptiles. University of California Press, Berkeley.
- Stebbins, R.C. 1985. A Field Guide to Western Amphibians and Reptiles. Houghton Mifflin Company, Boston, MA.
- Stebbins, R.C. 2003. A Field Guide to Western Reptiles and Amphibians. Houghton Mifflin Company, Boston, MA.
- Stebbins, R.C. and Lowe, C.H. 1951. Subspecific differentiation in the Olympic salamander *Rhyacotriton olympicus*. University of California Publications in Zoology 50:465–484.
- Stejneger, L. 1899. Description of a new genus and species of discoglossoid toad from North America. *Proceedings of the United States National Museum* 21:899–901.
- Stewart, G.R. 1972. An unusual record of sperm storage in a female garter snake (genus *Thamnophis*). *Herpetologica* 28:346–347.
- Stewart, I.T., Cayan, D.R., and Dettinger, M. 2005. Changes toward earlier streamflow timing across western North America. *Journal of Climate*. 18:1136–1155.
- Stoddard, M.A. and Hayes, J.P. 2005. The influence of forest management on headwater stream amphibians at multiple spatial scales. *Ecological Applications* 15:811–823.
- Stone, P.A. 2001. Movements and demography of the Sonoran mud turtle, *Kinosternon sonoriense*. *The Southwestern Naturalist* 46:41–53.
- Stone, P.A., Babb, M.E., Stanila, B.D., Kersey, G., and Stone, Z. 2005. Natural history notes: *Kinosternon sonoriense*. Diet. *Herpetological Review* 36:167–168.
- Stone, P.A., Stone, M.E.B., Stanila, B.D., and Lacey, K.J. 2011. Terrestrial flight response: a new context for terrestrial activity in Sonoran mud turtles. *The American Midland Naturalist* 165:128–136.
- Storer, T.I. 1925. A synopsis of the amphibia of California. University of California Publications in Zoology 27:1–342.
- Storer, T.I. 1930. Notes on the range and life-history of the Pacific fresh-water turtle, *Clemmys marmorata*. University of California Publications in Zoology 35:421–441.
- Storm, R.M. 1960. Notes on the breeding biology of the red-legged frog (*Rana aurora aurora*). *Herpetologica* 16:251–259.
- Storm, R.M. and Pimentel, R.A. 1954. A method for studying amphibian breeding populations. *Herpetologica* 10:161–166.
- Stralberg, D., Jongsomjit, D., Howell, C.A., Snyder, M.A., Alexander, J.D., Wiens, J.A., and Root, T.L. 2009. Re-shuffling of species with climate disruption: a no-analog future for California birds? *PLOS ONE* 4:e6825.
- Stromberg, M.R. 1997. Natural history notes: *Taricha torosa*. Response to fire. *Herpetological Review* 28:82–83.

- Stuart, S.N., Chanson, J.S., Cox, N.A., Young, B.E., Rodrigues, A.S.L., Fischman, D.L., and Waller, R.W. 2004. Status and trends of amphibian declines and extinctions worldwide. *Science* 306:1783–1786.
- Suarez, A. and Case, T. 2002. Bottom-up effects on persistence of a specialist predator: ant invasions and horned lizards. *Ecological Applications* 12:291–298.
- Suarez, A., Richmond, J., and Case, T. 2000. Prey selection in horned lizards following the invasion of Argentine ants in southern California. *Ecological Applications* 10:711–725.
- Sullivan, B.K. 1981. Distribution and relative abundance of snakes along a transect in California. *Journal of Herpetology* 15:247–248.
- Sullivan, B.K. and Fernandez, P.J. 1999. Breeding activity, estimated age-structure, and growth in Sonoran Desert anurans. *Herpetologica* 55:334–343.
- Sullivan, B.K. and Malmos, K.B. 1994. Call variation in the Colorado River toad (*Bufo alvarius*): behavioral and phylogenetic implications. *Herpetologica* 50:146–156.
- Sun, M.C. 2012. *Batrachochytrium dendrobatidis* Prevalence in Northern Red-Legged Frogs (*Rana aurora*): 10 Years Later. Master's Thesis. Humboldt State University, Arcata, CA.
- Survey and manage program. 2010. Annual Species Review. USDI Bureau of Land Management. Available from <http://www.blm.gov/or/plans/surveyandmanage/species.php>
- Sutherland, R.W., Dunning, P.R., and Baker, W.M. 2010. Amphibian encounter rates on roads with different amounts of traffic and urbanization. *Conservation Biology* 24:1626–1635.
- Sweet, S.S. 1992. Initial Report on the Ecology and Status of the Arroyo Toad (*Bufo microscaphus californicus*) on the Los Padres National Forest of Southern California, with Management Recommendations. Los Padres National Forest, Goleta, CA.
- Sweet, S.S. 1993. Second Report on the Biology and Status of the Arroyo Toad (*Bufo microscaphus californicus*) on the Los Padres National Forest of southern California. Los Padres National Forest, Goleta, CA.
- Sweet, S.S. and Leviton, A.E. 1983. Geographic distribution: *Rana aurora draytoni*. *Herpetological Review* 14:27.
- Sweet, S.S. and Sullivan, B.K. 2005. *Bufo californicus*. in *Amphibian declines: the conservation status of United States species*. Lannoo, M.J. (Ed.): University of California Press, Berkeley, pp.396–400.
- Syde, W.E. 1975. Breeding Habits, Embryonic Thermal Requirements and Embryonic and Larval Development of the Cascade Frog, *Rana cascadae* Slater. PhD Dissertation. Oregon State University, Corvallis.
- Tait, C.K. and Diller, L.V. 2006. Life history of the southern torrent salamander (*Rhyacotriton variegatus*) in coastal northern California. *Journal of Herpetology* 40:43–54.
- Tan, A.M. and Wake, D. 1995. MtDNA phylogeography of the California newt, *Taricha torosa* (Caudata, Salamandridae). *Molecular Phylogenetics and Evolution* 4:383–394.
- Tejon Ranch Conservancy. 2008. Tejon Ranch Conservation and Land Use Agreement. Available from: <http://www.tejonconservancy.org/>.
- Tevis, L., Jr. 1943. Field notes on a red rattlesnake in Lower California. *Copeia* 1943:242–245.
- Thomas, C.D., Cameron, A., Green, R.E., Bakkenes, M., Beaumont, L.J., Collingham, Y.C., Erasmus, B.F.N., de Siqueira, M.F., Grainger, A., and Hannah, L. 2004. Extinction risk from climate change. *Nature* 427:145–148.
- Thompson, C., Sweitzer, R., Gabriel, M., Purcell, K., Barrett, R. and Poppenga, R. 2014. Impacts of rodenticide and insecticide toxicants from marijuana cultivation sites on fisher survival rates in the Sierra National Forest, California. *Conservation Letters* 7:91–102.
- Thomson, R.C., Spinks, P.Q., and Shaffer, H.B. 2010. Distribution and abundance of invasive red-eared sliders (*Trachemys scripta elegans*) in California's Sacramento river basin and possible impacts on native western pond turtles (*Emys marmorata*). *Chelonian Conservation and Biology* 9:297–302.
- Tihen, J.A. 1962. Osteological observations on New World *Bufo*. *The American Midland Naturalist* 67:157–183.
- Tinkham, E.R. 1962. Notes on the occurrence of *Scaphiopus couchii* in California. *Herpetologica* 18:204.
- Tinkham, E.R. 1971. The biology of the Gila monster, in Bucherl, W. and Buckley, E.E. (Eds): *Venomous Animals and Their Venoms: Venomous Vertebrates*, Vol. 2. Academic Press, New York, pp.387–413.
- Todd, B.D., Scott, D.E., Pechmann, J.H.K., and Gibbons, J.W. 2011. Climate change correlates with rapid delays and advancements in reproductive timing in an amphibian community. *Proceedings of the Royal Society Series B* 278:2191–2197.
- Tollestrup, K. 1979. The Ecology, Social Structure, and Foraging Behavior of Two Closely Related

- Species of Leopard Lizards, *Gambelia silus* and *Gambelia wislizenii*. PhD Dissertation. University of California, Berkeley.
- Tollestrup, K. 1981. The social behavior and displays of 2 species of horned lizards, *Phrynosoma platyrhinos* and *Phrynosoma coronatum*. *Herpetologica* 37:130–141.
- Tracey, J.A. 2000. Movement of Red Diamond Rattlesnakes (*Crotalus ruber*) in Heterogeneous Landscapes in Coastal Southern California. Master's Thesis. University of California, San Diego.
- Tracey, J.A., Zhu, J., and Crooks, K. 2005. A set of nonlinear regression models for animal movement in response to a single landscape feature. *Journal of Agricultural Biological and Environmental Statistics* 10:1–18.
- Trenham, P.C. 1998. Demography, migration, and metapopulation structure of pond breeding salamanders. PhD Dissertation. University of California, Davis, California.
- Trenham, P.C. and Shaffer, H.B. 2005. Amphibian npland habitat use and its consequences for population viability. *Ecological Applications* 15:1158–1168.
- Trépanier, T.L. and Murphy, R.W. 2001. The Coachella Valley fringe-toed lizard (*Uma inornata*): genetic diversity and phylogenetic relationships of an endangered species. *Molecular Phylogenetics and Evolution* 18:327–334.
- Turner, F.B. and Medica, P.A. 1982. The distribution and abundance of the flat-tailed horned lizard (*Phrynosoma mcallii*). *Copeia* 1982:815–823.
- Turner, F.B., Rorabaugh, J.C., Nelson, E.C., and Jorgensen, M.C. 1980. A Survey of the Occurrence and Abundance of the Flat-Tailed Horned Lizard (*Phrynosoma mcallii*) in California. Contract YA-512-CT8-58. Unpublished Report.
- Turner, F.B., Weaver, D.C., and Rorabaugh, J.C. 1984. Effects of reduction in windblown sand on the abundance of the fringe-toed lizard (*Uma inornata*) in the Coachella Valley, California. *Copeia* 1984:370–378.
- Twining, H. and Hensley, A. 1943. Distribution of muskrats in California. *California Fish and Game* 29:64–78.
- Twitty, V.C. 1935. Two new species of *Triturus* from California. *Copeia* 1935:73–80.
- Twitty, V.C. 1942. The species of Californian *Triturus*. *Copeia* 1942:65–76.
- Twitty, V.C. 1955. Field experiments on the biology and genetic relationships of the Californian species of *Triturus*. *Journal of Experimental Zoology* 129:129–147.
- Twitty, V.C. 1959. Migration and speciation in newts. *Science* 130:1735–1743.
- Twitty, V.C. 1961. Experiments on homing behavior and speciation in *Taricha*, in Blair, W.F. (Ed.): *Vertebrate Speciation*. University of Texas Press, Austin, pp.415–459.
- Twitty, V.C. 1964. *Taricha rivularis* (Twitty), red-bellied newt. *Catalogue of American Amphibians and Reptiles* 9:1–2.
- Twitty, V.C. 1966. *Of Scientists and Salamanders*. WH Freeman & Company, San Francisco, CA.
- Twitty, V.C., Grant, D., and Anderson, O. 1964. Long distance homing in newt *Taricha rivularis*. *Proceedings of the National Academy of Sciences of the United States of America* 51:51–58.
- Twitty, V.C., Grant, D., and Anderson, O. 1966. Course and timing of homing migration in newt *Taricha rivularis*. *Proceedings of the National Academy of Sciences of the United States of America* 56:864–871.
- Twitty, V.C., Grant, D., and Anderson, O. 1967a. Initial homeward orientation after long-distance displacements in newt *Taricha rivularis*. *Proceedings of the National Academy of Sciences of the United States of America* 57:342–348.
- Twitty, V.C., Grant, D., and Anderson, O. 1967b. Home range in relation to homing in newt *Taricha rivularis* (Amphibia: Caudata). *Copeia* 1967:649–653.
- Tyler, T., Liss, W.J., Ganio, L.M., Larson, G.L., Hoffman, R., Deimling, E., and Lomnický, G. 1998. Interaction between introduced trout and larval salamanders (*Ambystoma macrodactylum*) in high-elevation lakes. *Conservation Biology* 12:94–105.
- Ultsch, G.R. 2006. The ecology of overwintering among turtles: where turtles overwinter and its consequences. *Biological Reviews* 81:339–367.
- US Census Bureau. 2013. California Demographic Profile. Accessed August 27, 2014. census.gov.
- USDA (United States Department of Agriculture). 2007. Census of Agriculture. Accessed August 27, 2014. http://www.agcensus.usda.gov/Publications/2007/Online_Highlights/County_Profiles/California/index.asp.
- USFWS (United States Fish and Wildlife Service). 1996. Endangered and threatened wildlife and plants: determination of threatened status for the California red-legged frog. *Federal Register* 61:25813–25833.
- USFWS. 1997. Recovery Plan for the Threatened Marbled Murrelet (*Brachyramphus marmoratus*) in Washington, Oregon, and California. USFWS, Portland, OR.

- USFWS. 1999. Recovery Plan for the Arroyo Southwestern Toad. USFWS, Portland, OR.
- USFWS. 2002. Recovery Plan for the California Red-Legged Frog (*Rana aurora draytonii*). USFWS, Portland, OR.
- USFWS. 2008. Endangered and Threatened Wildlife and Plants; 90-Day Finding on Petition to List the Amargosa River Population of the Mojave Fringe-Toed Lizard (*Uma scoparia*) as Threatened or Endangered with Critical Habitat. Ventura, CA.
- USFWS. 2009. Arroyo Toad 5-Year Review: Summary and Evaluation. Ventura, CA.
- USFWS. 2011a. Endangered and threatened wildlife and plants; withdrawal of proposed rule to list the flat-tailed horned lizard as threatened. Federal Register 76:14210–14268.
- USFWS. 2011b. Endangered and threatened wildlife and plants; 12-month finding on a petition to list the Amargosa River population of the Mojave fringe-toed lizard as an endangered or threatened distinct population segment. Federal Register 76:61321–61330.
- USFWS. 2014. Endangered and threatened wildlife and plants; threatened status for the Oregon spotted frog. Federal Register 79:51658–51710.
- USNPS (United States National Park Service). 2010. Channel Islands National Park: Restoring Santa Cruz Island. Accessed August 27, 2014. <http://www.nps.gov/chis/naturescience/restoring-santa-cruz-island.htm>.
- Valentine, B.D. and Dennis, D.M. 1964. A comparison of the gill-arch system and fins of three genera of larval salamanders, *Rhyacotriton*, *Gyrinophilus*, and *Ambystoma*. Copeia 1964:196–201.
- Van Denburgh, J. 1894. Descriptions of three new lizards from California and lower California, with a note on *Phrynosoma blainvillii*. Proceedings of the California Academy of Sciences 4:296–301.
- Van Denburgh, J. 1895. Notes on the habits and distribution of *Autodax iecanus*. Proceedings of the California Academy of Sciences 5:776–778.
- Van Denburgh, J. 1897. The reptiles of the Pacific coast and Great Basin: An account of the species known to inhabit California, and Oregon, Washington, Idaho and Nevada. Occasional Papers of the California Academy of Sciences, San Francisco.
- Van Denburgh, J. 1922. The reptiles of western North America: an account of the species known to inhabit California and Oregon, Washington, Idaho, Utah, Nevada, Arizona, British Columbia, Sonora and lower California. Vol. 1. Lizards. Occasional Papers of the California Academy of Sciences 10:1–611.
- Van Denburgh, J. and Slevin, J.R. 1913. A list of the amphibians and reptiles of Arizona, with notes on the species in the collection of the Academy. Proceedings of the California Academy of Sciences 3:391–454.
- Van Loben Sels, R.C., Congdon, J.D., and Austin, J.T. 1997. Life history and ecology of the Sonoran mud turtle (*Kinosternon sonoriense*) in southeastern Arizona: a preliminary report. Chelonian Conservation and Biology 2:338–344.
- Van Wagner, T.J. 1996. Selected Life-History and Ecological Aspects of a Population of Foothill Yellow-Legged Frogs (*Rana boylei*) from Clear Creek, Nevada County, California. Master's Thesis. California State University, Chico.
- Vesely, D.G. 1996. Terrestrial Amphibian Abundance and Species Richness in Headwater Riparian Buffer Strips, Oregon Coast Range. Master's Thesis. Oregon State University, Corvallis, OR. 48pp.
- Vesely, D.G. and McComb, W.C. 2002. Salamander abundance and amphibian species richness in riparian buffer strips in the Oregon Coast Range. Forest Science 48:291–297.
- Vitt, L.J. 1975. Observations on reproduction in five species of Arizona snakes. Herpetologica 31:83–84.
- Vitt, L.J. and Ohmart, R.D. 1978. Herpetofauna of the lower Colorado River: Davis Dam to the Mexican border. Proceedings of the Western Foundation of Vertebrate Zoology 2:35–72.
- Von Bloeker, J.C.J. 1942. Fauna and flora of the El Segundo sand dunes: 13. Amphibians and reptiles of the dunes. Bulletin of the Southern California Academy of Sciences 41:29–38.
- Vredenburg, V.T. 2004. Reversing introduced species effects: experimental removal of introduced fish leads to rapid recovery of a declining frog. Proceedings of the National Academy of Sciences of the United States of America 101:7646–7650.
- Vredenburg, V.T., Bingham, R., Knapp, R., Morgan, J.A.T., Moritz, C., and Wake, D. 2007. Concordant molecular and phenotypic data delineate new taxonomy and conservation priorities for the endangered mountain yellow-legged frog. Journal of Zoology 271:361–374.
- Vredenburg, V.T., Knapp, R.A., Tunstall, T.S., and Briggs, C.J. 2010. Dynamics of an emerging disease drive large-scale amphibian population extinctions. Proceedings of the National Academy of Sciences of the United States of America 107:9695–9700.

- Wahbe, T.R. and Bunnell, F.L. 2003. Relations among larval tailed frogs, forest harvesting, stream microhabitat, and site parameters in southwestern British Columbia. *Canadian Journal of Forest Research* 33:1256–1266.
- Wahbe, T.R., Bunnell, F.L., and Bury, R.B. 2004. Terrestrial movements of juvenile and adult tailed frogs in relation to timber harvest in coastal British Columbia. *Canadian Journal of Forest Research* 34:2455–2466.
- Wake, D.B. 1996. A new species of *Batrachoseps* (Amphibia: Plethodontidae) from the San Gabriel Mountains, southern California. *Contributions in Science from the Natural History Museum of Los Angeles County* 463:1–12.
- Wake, D.B. and Jockusch, E.L. 2000. Detecting species borders using diverse data sets: plethodontid salamanders in California, in Bruce, R.C., Jaeger, R.G., and Houck, L.D. (Eds): *The Biology of the Plethodontidae*. Kluwer Academic/Plenum Publishers, New York, pp.95–119.
- Wake, D.B., and Papenfuss, T.J., 2005. *Hydromantes platycephalus*, in Lannoo, M.J. (Ed.): *Amphibian Declines: The Conservation Status of United States Species*, University of California Press, Berkeley, pp.83–784.
- Wake, D.B., Yanev, K.P., and Hansen, R.W. 2002. New species of slender salamander, genus *Batrachoseps*, from the southern Sierra Nevada of California. *Copeia* 2002:1016–1028.
- Wallace, R.L. and Diller, L.V. 1998. Length of the larval cycle of *Ascaphus truei* in coastal streams of the redwood regions, northern California. *Journal of Herpetology* 32:404–409.
- Walther, G.R., Post, E., Convey, P., Menzel, A., Parmesan, C., Beebee, T.J.C., Fromentin, J.M., Hoegh-Guldberg, O., and Bairlein, F. 2002. Ecological responses to recent climate change. *Nature* 416:389–395.
- Wang, I.J. 2009b. Fine-scale population structure in a desert amphibian: landscape genetics of the black toad (*Bufo exsul*). *Molecular Ecology* 18:3847–3856.
- Wang, I.J. 2012. Environmental and topographical variables shape patterns of genetic structure and effective population size in the Yosemite toad. *Diversity and Distributions* 18:1033–1041.
- Wang, I.J., Johnson, J.R., Johnson, B.B., and Shaffer, H.B. 2011. Effective population size is strongly correlated with breeding pond size in the endangered California tiger salamander, *Ambystoma californiense*. *Conservation Genetics* 12:911–920.
- Wang, I.J., Savage, W.K., and Shaffer, H.B. 2009. Landscape genetics and least-cost path analysis reveal unexpected dispersal routes in the California tiger salamander (*Ambystoma californiense*). *Molecular Ecology* 18:1365–1374.
- Wang, J. 2009a. A new method for estimating effective population sizes from a single sample of multilocus genotypes. *Molecular Ecology* 18:2148–2164.
- Ward, P.S. 1987. Distribution of the introduced Argentine ant (*Iridomyrmex humilis*) in natural habitats of the lower Sacramento Valley and its effects on the indigenous ant fauna. *Hilgardia* 55:1–16.
- Watson, J.W., McAllister, K.R., and Pierce, D.J. 2003. Home ranges, movements, and habitat selection of Oregon spotted frogs (*Rana pretiosa*). *Journal of Herpetology* 37:292–300.
- Watters, T.S. and Kats, L.B. 2006. Longevity and breeding site fidelity in the California newt (*Taricha torosa*): a long-term study showing the efficacy of pit tagging. *Herpetological Review* 37:151.
- Weinstein, S.B. 2009. An aquatic disease on a terrestrial salamander: individual and population level effects of the amphibian chytrid fungus, *Batrachochytrium dendrobatidis*, on *Batrachoseps attenuatus* (Plethodontidae). *Copeia* 2009:653–660.
- Wells, M.T. 1998. *Wildlife Management Plan for Torrey Pines State Reserve: Terrestrial Vertebrates*. *Wildlife Management Plan for Torrey Pines State Reserve*, Torrey Pines State Reserve, La Jolla, CA.
- Welsh, H.H., Jr. 1985. Geographic distribution: *Ascaphus truei*. *Herpetological Review* 16: 59.
- Welsh, H.H., Jr. 1988. An ecogeographic analysis of the herpetofauna of the Sierra San Pedro Martir region, Baja California: with a contribution to the biogeography of the Baja California herpetofauna. *Proceedings of the California Academy of Sciences* 46:1–72.
- Welsh, H.H., Jr. 1990. Relictual amphibians and old-growth forests. *Conservation Biology* 4:309–319.
- Welsh, H.H., Jr. 2011. Frogs, fish and forestry: an integrated watershed network paradigm conserves biodiversity and ecological services. *Diversity* 2011:503–530.
- Welsh, H.H., Jr., and Bury, R.B. 2005. *Plethodon elongates*, in Lannoo, M.J. (Ed.): *Amphibian Declines: The Conservation Status of United States Species*. University of California Press, Berkeley, pp.806–807.
- Welsh, H.H., Jr., Fellers, G., and Liud, A. 2007. Amphibian populations in the terrestrial

- environment: Is there evidence of declines of terrestrial forest amphibians in northwestern California? *Journal of Herpetology* 41: 469–482.
- Welsh, H.H., Jr., and Hodgson, G.R. 2008. Amphibians as metrics of critical biological thresholds in forested headwater streams of the Pacific Northwest, U.S.A. *Freshwater Biology* 53:1470–1488.
- Welsh, H.H., Jr., and Hodgson, G.R. 2011. Spatial relationships in a dendritic network: the herpetofaunal metacommunity of the Mattole River catchment of northwest California. *Ecography* 34:49–66.
- Welsh, H.H., Jr., Hodgson, G.R., Duda, J.J., and Emlen, J.M. 2010. Faunal assemblages and multi-scale habitat patterns in headwater tributaries of the South Fork Trinity River: an unregulated river embedded within a multiple-use landscape. *Animal Biodiversity and Conservation* 33:63–87.
- Welsh, H.H., Jr., Hodgson, G.R., and Lind, A.J. 2005. Ecogeography of the herpetofauna of a northern California watershed: linking species patterns to landscape processes. *Ecography* 28:521–536.
- Welsh, H.H., Jr., and Lind, A.J. 1988. Old-growth forests and the distribution of terrestrial herpetofauna, in Szaro, R.C., Severson, R.E., and Patton, D.R. (Eds): *Management of Amphibians, Reptiles, and Small Mammals in North America*. General Technical Report RM-166. US Department of Agriculture, Forest Service, Rocky Mountain Forest and Range Experiment Station, Fort Collins, CO, pp.439–459.
- Welsh, H.H., Jr., and Lind, A.J. 1991. The structure of the herpetofaunal assemblage in the Douglas-fir/hardwood forests of northwestern California and southwestern Oregon. US Department of Agriculture, Forest Service, General Technical Report PNW 285:395–413.
- Welsh, H.H., Jr., and Lind, A.J. 1992. Population ecology of two relictual salamanders from the Klamath Mountains of northwestern California, in McCullough, D. and Barret, R. (Eds): *Wildlife 2001: Populations*. Elsevier Science Publications Limited, London, pp.419–437.
- Welsh, H.H., Jr., and Lind, A.J. 1995. Habitat correlates of the Del Norte salamander, *Plethodon elongatus* (Caudata, Plethodontidae), in northwestern California. *Journal of Herpetology* 29:198–210.
- Welsh, H.H., Jr., and Lind, A.J. 1996. Habitat correlates of the southern torrent salamander, *Rhyacotriton variegatus* (Caudata: Rhyacotritonidae), in northwestern California. *Journal of Herpetology* 30:385–398.
- Welsh, H.H., Jr., and Lind, A.J. 2002. Multiscale habitat relationships of stream amphibians in the Klamath-Siskiyou Region of California and Oregon. *Journal of Wildlife Management* 66:581–602.
- Welsh, H.H., Jr., and Ollivier, L.M. 1998. Stream amphibians as indicators of ecosystem stress: a case study from California's redwoods. *Ecological Applications* 8:1118–1132.
- Welsh, H.H., Jr., Pope, K.L., and Boiano, D. 2006. Sub-alpine amphibian distributions related to species palatability to non-native salmonids in the Klamath mountains of northern California. *Diversity and Distributions* 12:298–309.
- Welsh, H.H., Jr., Roelofs, T.D., and Frissell, C.A. 2000. Aquatic ecosystems of the redwood region, in Noss, R. (Ed.): *The Redwood Forest*. Island Press, Washington, DC, pp.165–200.
- Westerling, A.L. and Bryant, B.P. 2008. Climate change and wildfire in California. *Climatic Change* 87:S231–S249.
- Westerling, A.L., Bryant, B.P., Preisler, H.K., Holmes, T.P., Hidalgo, H.G., Das, T., and Shrestha, S.R. 2011. Climate change and growth scenarios for California wildfire. *Climatic Change* 109:S445–S463.
- Westerling, A.L., Cayan, D.R., Brown, T.J., Hall, B.L., and Riddle, L.G. 2004. Climate, Santa Ana winds and autumn wildfires in southern California. *EOS, Transactions, American Geophysical Union* 85:289–296.
- Wheeler, C.A., Garwood, J.M., and Welsh, H.H., Jr. 2005. Natural history notes: *Rana boylei*. Physiological skin color transformation. *Herpetological Review* 36:164–165.
- Wheeler, C.A. and Welsh, H.H., Jr. 2008. Mating strategy and breeding patterns of the foothill yellow-legged frog (*Rana boylei*). *Herpetological Conservation and Biology* 3:128–142.
- Wheeler, C.A., Welsh, H.H., Jr., and Roelofs, T. 2006. Oviposition Site Selection, Movement, and Spatial Ecology of the Foothill Yellow-Legged Frog (*Rana boylei*). California Department of Fish and Game, Sacramento.
- White, M. and Kolb, J.A. 1974. A preliminary study of *Thamnophis* near Sagehen Creek, California. *Copeia* 1974:126–136.
- Wiens, J. and Titus, T. 1991. A phylogenetic analysis of *Spea* (Anura, Pelobatidae). *Herpetologica* 47:21–28.
- Wilgenbusch, J. and De Queiroz, K. 2000. Phylogenetic relationships among the phryno-

- somatid sand lizards inferred from mitochondrial DNA sequences generated by heterogeneous evolutionary processes. *Systematic Biology* 49:592–612.
- Wilkins, R.N. and Peterson, N.P. 2000. Factors related to amphibian occurrence and abundance in headwater streams draining second-growth Douglas-fir forests in southwestern Washington. *Forest Ecology and Management* 139:79–91.
- Williams, D.F. 1986. *Mammalian Species of Special Concern in California*. California Department of Fish and Game, Sacramento.
- Wilson, G.A. and Rannala, B. 2003. Bayesian inference of recent migration rates using multilocus genotypes. *Genetics* 163:1177–1191.
- Wilson, L.D. 1971. The coachwhip snake *Masticophis flagellum* (Shaw): taxonomy and distribution. *Tulane Studies in Zoology and Botany* 16: 31–99.
- Wilson, L.D. 1973. *Masticophis flagellum*. *Catalogue of American Amphibians and Reptiles* 145:1–4.
- Wiseman, K.D. and Bettaso, J. 2007. Natural history notes: *Rana boylei*. Cannibalism and predation. *Herpetological Review* 38:193.
- Wiseman, K.D., Marlow, K.R., Jackman, R.E., and Drennan, J.E. 2005. Natural history notes: *Rana boylei*. Predation. *Herpetological Review* 36:162–163.
- Wojtaszek, B.F., Staznik, B., Chartrand, D.T., Stephenson, G.R., and Thompson, D.G. 2004. Effects of Vision® herbicide on mortality, avoidance response, and growth of amphibian larvae in two forest wetlands. *Environmental Toxicology and Chemistry* 23:832–842.
- Wollmuth, L.P., Crawshaw, L.I., Forbes, R.B., and Grahn, D.A. 1987. Temperature selection during development in a montane anuran species, *Rana cascadae*. *Physiological Zoology* 60:472–480.
- Wong, B. and Beauchamp, B. 2003. Movement, home range, and activity patterns of the horned lizard, *Phrynosoma mcallii*. *Journal of Herpetology* 37:679–686.
- Wood, D.A., Fisher, R.N., and Reeder, T.W. 2008. Novel patterns of historical isolation, dispersal, and secondary contact across Baja California in the Rosy Boa (*Lichanura trivirgata*). *Molecular Phylogenetics and Evolution* 46:484–502.
- Wood, D.A. and Richmond, J.Q. 2003. Geographic distribution: *Diadophis punctatus*. *Herpetological Review* 34:169
- Woodhams, D.C., Bosch, J., Briggs, C.J., Cashins, S., Davis, L.R., Lauer, A., Muths et al. 2011. Mitigating amphibian disease: strategies to maintain wild populations and control chytridiomycosis. *Frontiers in Zoology* 8:8
- Woodson, W.D. 1949. Gila monster in California. *Herpetologica* 5:151.
- Woodward, B.D. 1982. Sexual selection and nonrandom mating patterns in desert anurans (*Bufo woodhousei*, *Scaphiopus couchi*, *S. multiplicatus* and *S. bombifrons*). *Copeia* 1982:351–355.
- Worthylake, K.M. and Hovingh, P. 1989. Mass mortality of salamanders (*Ambystoma tigrinum*) by bacteria (*Acinetobacter*) in an oligotrophic seepage mountain lake. *The Great Basin Naturalist* 49:364–372.
- Wright, A.H. and Wright, A.A. 1949. *Handbook of Frogs and Toads of the United States and Canada*. Comstock Publishing Co., Ithaca, NY.
- Wright, A.H., and Wright, A.A. 1957. *Handbook of Snakes of the United States and Canada*. Comstock Publishing Co., Ithaca, NY.
- Wright, A.N., Hijmans, R.J., Schwartz, M.W., and H.B. Shaffer. 2013. *California Amphibian and Reptile Species of Future Concern: Conservation and Climate Change*. Final Report to the California Department of Fish and Wildlife, Nongame Wildlife Program, Task 12, Contract No. P0685904.
- Wroble, J. and Waters, D. 1989. Summary of tailed frog (*Ascaphus truei*) and Olympic salamander (*Rhyacotriton olympicus variegatus*) stream surveys for the Pacific Lumber Company, October 1987 to September 1988. Pacific Lumber Co., Scotia, CA.
- Yanev, K.P. 1978. *Evolutionary Studies of the Plethodontid Salamander Genus Batrachoseps*. PhD Dissertation. University of California, Berkeley.
- Yanev, K.P. 1980. Biogeography and distribution of three parapatric salamander species in coastal and borderland California, in Power, D.M. (Ed.): *The California Islands: Proceedings of a Multidisciplinary Symposium*. Santa Barbara Museum of Natural History, Santa Barbara, California, pp.531–550.
- Yanev, K.P. and Wake, D.B. 1981. Genic differentiation in a relict desert salamander, *Batrachoseps campi*. *Herpetologica* 37:16–28.
- Yarnell, S.M. 2000. *The Influence of Sediment Supply and Transport Capacity on Foothill Yellow-Legged Frog Habitat, South Yuba River, California*. Master's Thesis. University of California, Davis.
- Yarnell, S.M. 2005. *Spatial Heterogeneity of Rana boylei Habitat: Physical Processes, Quantification and Ecological Meaningfulness*. PhD Dissertation. University of California, Davis.
- Yarrow, H.C. 1882. *Descriptions of new species of reptiles and amphibians in the United States*

- National Museum. Proceedings of the United States National Museum 5:438-443.
- Young, K.V. 2010. Comparative Ecology of Narrowly Sympatric Horned Lizards under Variable Climatic Conditions. PhD Dissertation. Utah State University, Logan.
- Young, K.V. and Royle, J. 2005. Abundance and Site Occupancy of Flat-Tailed Horned Lizard (*Phrynosoma mcallii*) Populations in Arizona and California. Final Report to US Bureau of Reclamation, US Navy, and Arizona Game and Fish Department.
- Young, K.V. and Young, A. 2000. Scientific study of the flat-tailed horned lizard, *Phrynosoma mcallii*. Final Report to US Department of the Navy.
- Young, K.V. and Young, A.T. 2005. Indirect Effects of Development on the Flat-Tailed Horned Lizard. Final Report to Arizona Game and Fish Department, Yuma, 11pp.
- Zalusky, S.B., Gaudin, A.J., and Swanson, J.R. 1980. A comparative study of cranial osteology in the North American sand lizards, genus *Uma* (Reptilia: Iguanidae). Copeia 1980:296-310.
- Zeiner, D.C., Laudenslayer, W.F., Jr., and Mayer, K.E. 1988. California's Wildlife. Vol. 1. Amphibians and Reptiles. California Department of Fish and Game, Sacramento.
- Zweifel, R.G. 1955. Ecology, distribution, and systematics of frogs of the *Rana boylei* group. University of California Publications in Zoology 54:207-292.

This page intentionally left blank

APPENDIX 1

List of Native Amphibian and Reptile Taxa Occurring in California

Taxon ¹	Common name	CDFG special animal	USFWS ²	CDFW ¹	IUCN ⁴	USFS ⁵	BLM
<i>Aquas</i>							
Ascaphidae							
<i>Ascaphus trueti</i>	Coastal tailed frog	X		SSC	LC		
Bufo							
<i>Bufo alvarius</i>	Sonoran Desert toad	X		SSC	LC		
<i>Bufo boreas boreas</i>	Western toad				NT		
<i>Bufo boreas halophilus</i>	California western toad				NT		
<i>Bufo californicus</i>	Arroyo toad	X	E	SSC	E		
<i>Bufo canorus</i>	Yosemite toad	X	T	SSC	E	\$	
<i>Bufo cognatus</i>	Great Plains toad				LC		
<i>Bufo exsul</i>	Black road	X			V	\$	\$
<i>Bufo punctatus</i>	Red-spotted toad				LC		
<i>Bufo woodhousii</i>	Woodhouse's toad				LC		
Hylidae							
<i>Pseudacris cadaverina</i>	California treefrog				LC		
<i>Pseudacris regilla?</i>	Pacific treefrog				LC		
Ranidae							
<i>Rana aurora</i>	Northern red-legged frog	X		SSC	LC	\$	
<i>Rana boylei</i>	Foothill yellow-legged frog	X		SSC	NT	\$	\$

(continued)

Taxon ¹	Common name	CDFG special animal	USFWS ²	CDFW ³	IUCN ⁴	USFS ⁵	BLM
Anura							
<i>Rana cascadae</i>	Cascades frog	X		SSC	NT	S	
<i>Rana draytonii</i>	California red-legged frog	X	T	SSC	V		
<i>Rana muscosa</i>	Southern Mountain yellow-legged frog	X	E	E	E	S	
<i>Rana pipiens</i> ^{6a}	Northern leopard frog	X		SSC	LC		
<i>Rana pretiosa</i> ^{6b}	Oregon spotted frog	X	T	SSC	V		S
<i>Rana sierrae</i>	Sierra Nevada yellow-legged frog	X	E	E	E	S	
<i>Rana yavapaiensis</i>	Lowland leopard frog	X		SSC	LC		S
Scaphiropodidae							
<i>Scaphiopus couchii</i>	Couch's spadefoot	X		SSC	LC		S
<i>Spea hammondi</i>	Western spadefoot	X		SSC	NT		S
<i>Spea intermontana</i>	Great basin spadefoot				LC		
Caudata							
Ambystomatidae							
<i>Ambystoma californiense</i>	California tiger salamander	X	T	T	V		
<i>Ambystoma californiense</i> "Santa Barbara"	Santa Barbara tiger salamander	X	E	T	V		
<i>Ambystoma californiense</i> "Sonoma"	Sonoma tiger salamander	X	E	T	V		
<i>Ambystoma gracile</i>	Northwestern salamander				LC		
<i>Ambystoma macrodactylum croceum</i>	Santa Cruz long-toed salamander	X	E	E, FP	LC		
<i>Ambystoma macrodactylum sigillatum</i>	Southern long-toed salamander			SSC	LC		
Dicamptodontidae							
<i>Dicamptodon ensatus</i>	California giant salamander			SSC	NT		
<i>Dicamptodon tenebrosus</i>	Pacific giant salamander				LC		

Plethodontidae

<i>Aneides ferreus</i>	Clouded salamander							NT
<i>Aneides flavipunctatus</i>	Black salamander							NT
<i>Aneides flavipunctatus niger</i>	Santa Cruz black salamander			SSC				NT
<i>Aneides flavipunctatus "shasta"</i> ¹¹	Shasta black salamander							NT
<i>Aneides lugubris</i>	Arboreal salamander							LC
<i>Aneides vagrans</i>	Wandering salamander							NT
<i>Batrachoseps alasierrae</i>	Greenhorn Mountains slender salamander							
<i>Batrachoseps attenuatus</i>	California slender salamander							LC
<i>Batrachoseps bramei</i>	Fairview slender salamander							S
<i>Batrachoseps campi</i>	Inyo Mountains salamander	X		SSC	E	S	S	S
<i>Batrachoseps diabolicus</i>	Hell Hollow slender salamander	X						DD
<i>Batrachoseps gabrieli</i>	San Gabriel Mountains slender salamander	X						DD S
<i>Batrachoseps gabilanensis</i>	Gabilan Mountains slender salamander							LC
<i>Batrachoseps gregarius</i>	Gregarius slender salamander	X						LC
<i>Batrachoseps incognitus</i>	San Simeon slender salamander	X						DD S
<i>Batrachoseps kawia</i>	Sequoia slender salamander	X						DD
<i>Batrachoseps luciae</i>	Santa Lucia Mountains slender salamander	X						LC
<i>Batrachoseps major aridus</i>	Desert slender salamander	X		E	E			LC
<i>Batrachoseps major major</i>	Garden slender salamander							LC
<i>Batrachoseps minor</i>	Lesser slender salamander	X		SSC				DD S
<i>Batrachoseps nigriventris</i>	Black-bellied slender salamander							LC

(continued)

Taxon ¹	Common name	CDFG special animal	USFWS ¹	CDPW ³	IUCN ⁴	USFS ⁵	BLM
Caudata							
<i>Batrachoseps pacificus</i>	Channel Islands slender salamander	X			LC		
<i>Batrachoseps regius</i>	Kings River slender salamander	X			V	S	
<i>Batrachoseps relictus</i>	Relictual slender salamander	X		SSC	DD	S	
<i>Batrachoseps robustus</i>	Kern Plateau salamander	X			NT		
<i>Batrachoseps simatus</i>	Kern Canyon slender salamander	X		T	V	S	
<i>Batrachoseps stebbinsi</i>	Tehachapi slender salamander	X		T	V		S
<i>Ensatina eschscholtzii croceator</i>	Yellow-blotched ensatina	X			LC	S	S
<i>Ensatina eschscholtzii eschscholtzii</i>	Monterey ensatina				LC		
<i>Ensatina eschscholtzii klauberi</i>	Large-blotched ensatina	X			LC	S	
<i>Ensatina eschscholtzii oregonensis</i>	Oregon ensatina				LC		
<i>Ensatina eschscholtzii picta</i>	Painted ensatina				LC		
<i>Ensatina eschscholtzii platensis</i>	Sierra Nevada ensatina				LC		
<i>Ensatina eschscholtzii xanthoptica</i>	Yellow-eyed ensatina				LC		
<i>Hydromantes brunus</i>	Limestone salamander	X		T, FP	V	S	S
<i>Hydromantes platycephalus</i> ¹¹	Mount Lyell salamander	X			LC		
<i>Hydromantes shastae</i>	Shasta salamander	X		T	V	S	S
<i>Methodon asupak</i>	Scott River salamander	X		T	V		
<i>Methodon dunni</i>	Dunn's salamander				LC		
<i>Methodon elongatus</i>	Del Norte salamander	X			NT		
<i>Methodon stormi</i>	Siskiyou Mountains salamander	X		T	E	S	
Rhyacotritonidae							
<i>Rhyacotriton variegatus</i>	Southern torrent salamander	X		SSC	LC	S	

Salamandridae							
<i>Taricha granulosa</i>	Rough-skinned newt						LC
<i>Taricha rivularis</i>	Red bellied newt			SSC			LC
<i>Taricha sierrae</i>	Sierra newt						LC
<i>Taricha torosa</i>	Coast Range newt	X		SSC			LC
Squamata—Lizards							
Anguillidae							
<i>Elgaria coerulea coerulea</i>	San Francisco alligator lizard						LC
<i>Elgaria coerulea palmeri</i>	Sierra Nevada alligator lizard						LC
<i>Elgaria coerulea principis</i>	Northwestern alligator lizard						LC
<i>Elgaria coerulea shastensis</i>	Shasta alligator lizard						LC
<i>Elgaria multicarinata multicarinata</i>	California alligator lizard						LC
<i>Elgaria multicarinata scincicauda</i>	Oregon alligator lizard						LC
<i>Elgaria multicarinata webbiai</i>	San Diego alligator lizard						LC
<i>Elgaria panamintina</i>	Panamint alligator lizard	X		SSC	Y	S	S
Anniellidae							
<i>Anniella pulchra pulchra</i> ^H	Silvery legless lizard	X		SSC			LC
<i>Anniella pulchra nigra</i>	Black legless lizard	X		SSC			LC
Crotaphytidae							
<i>Crotaphytus bicinctores</i>	Great Basin collared lizard						LC
<i>Crotaphytus vestigiium</i>	Baja California collared lizard						LC
<i>Gambelia copeii</i>	Cope's leopard lizard			SSC			LC
<i>Gambelia sila</i>	Blunt-nosed leopard lizard	X	E	E, FP			E
<i>Gambelia wislizenii</i>	Long-nosed leopard lizard						LC

Taxon ¹	Common name	CDFG special animal	USFWS ⁴	CDFW ³	IUCN ⁶	USFS ⁵	BLM
Squamata—Lizards							
Gekkonidae							
<i>Coleonyx variegatus</i>	Barefoot gecko	X		T	LC		\$
<i>Coleonyx variegatus abboti</i>	San Diego banded gecko	X		SSC	LC		
<i>Coleonyx variegatus variegatus</i>	Desert banded gecko				LC		
<i>Phyllodactylus nocticolus</i>	Peninsular leaf-toed gecko				LC		
Helodermatidae							
<i>Heloderma suspectum cinetum</i>	Banded Gila monster	X		SSC	NT		\$
Iguanidae							
<i>Dipsosaurus dorsalis</i>	Desert iguana				LC		
<i>Sauromalus ater</i>	Common chuckwalla				LC		
Phrynosomatidae							
<i>Callisaurus draconoides</i>	Zebra-tailed lizard				LC		
<i>Petrosaurus mearnsi</i>	Banded rock lizard				LC		
<i>Phrynosoma blainwillii</i> ²	Coast horned lizard	X		SSC	LC	\$	\$
<i>Phrynosoma douglasii</i>	Pigmy short-horned lizard				LC		
<i>Phrynosoma mullii</i>	Flat-tailed horned lizard	X		SSC	NT	\$	\$
<i>Phrynosoma platyrhinos calidiarum</i>	Southern desert horned lizard				LC		
<i>Phrynosoma platyrhinos platyrhinos</i>	Northern desert horned lizard				LC		
<i>Sceloporus graciosus gracilis</i>	Western sagebrush lizard				LC		
<i>Sceloporus graciosus graciosus</i>	Northern sagebrush lizard	X			LC		\$
<i>Sceloporus graciosus vandenburghianus</i>	Southern sagebrush lizard				LC		

<i>Sceloporus magister uniformis</i> ⁴	Yellow-backed desert spiny lizard					LC
<i>Sceloporus magister transversus</i>	Barred desert spiny lizard					LC
<i>Sceloporus occidentalis becki</i>	Island fence lizard					LC
<i>Sceloporus occidentalis biserialis</i>	San Joaquin fence lizard					LC
<i>Sceloporus occidentalis bocourti</i>	Coast Range fence lizard					LC
<i>Sceloporus occidentalis longipes</i>	Great Basin fence lizard					LC
<i>Sceloporus occidentalis occidentalis</i>	Northwestern fence lizard					LC
<i>Sceloporus occidentalis taylori</i>	Sierra fence lizard					LC
<i>Sceloporus orcutti</i>	Granite spiny lizard					LC
<i>Uma inornata</i>	Coachella Valley fringe-toed lizard	X	F	F	E	
<i>Uma notata</i>	Colorado Desert fringe-toed lizard	X			SSC	NT S
<i>Uma scoparia</i>	Mojave fringe-toed lizard	X			SSC	LC S
<i>Urosaurus graciosus</i>	Long-tailed brush lizard					LC
<i>Urosaurus nigricaudus</i>	Baja California brush lizard					LC
<i>Urosaurus ornatus</i>	Ornate tree lizard					LC
<i>Uta stansburiana elegans</i>	Western common side-blotched lizard					LC
<i>Uta stansburiana nevadensis</i>	Nevada common side-blotched lizard					LC
<i>Uta stansburiana stansburiana</i>	Northern common side-blotched lizard					LC
Scincidae						
<i>Plestiodon gilberti</i>	Gilbert's skink					LC
<i>Plestiodon skiltonianus skiltonianus</i>	Western skink					LC
<i>Plestiodon skiltonianus interparietalis</i>	Coronado skink	X				LC S
Teiidae						
<i>Aspidoscelis hyperythra</i>	Orange-throated whiptail	X				LC S

Taxon ¹	Common name	CDFG special animal	USFWS ²	CDFW ¹	IUCN ⁴	USFS ⁵	BLM
Squamata—Lizards							
<i>Aspidoscelis tigris munda</i>	California whiptail				LC		
<i>Aspidoscelis tigris stejnegeri</i>	Coastal whiptail	X		SSC	LC		
<i>Aspidoscelis tigris tigris</i>	Great Basin whiptail				LC		
Xantusiidae							
<i>Xantusia gracilis</i>	Sandstone night lizard	X		SSC	V		
<i>Xantusia henshawi</i>	Henshaw's night lizard				LC		
<i>Xantusia riversiana</i>	Island night lizard	X			LC		
<i>Xantusia vigilis sierrae</i> ¹³	Sierra night lizard	X		SSC	LC		
<i>Xantusia vigilis vigilis</i> ¹⁴	Desert night lizard				LC		
<i>Xantusia wigginsi</i>	Baja California night lizard				LC		
<i>Xantusia</i> sp. "Yucca Valley"	Yucca Valley night lizard				LC		
<i>Xantusia</i> sp. "San Jacinto"	San Jacinto night lizard				LC		
Squamata—Snakes							
Boidae							
<i>Charina bottae bottae</i>	Rubber boa				LC		
<i>Charina bottae umbratica</i>	Southern rubber boa	X		T	LC	S	
<i>Lichanura orcutti</i> ¹⁵	California rosy boa	X			LC	S	
Colubridae							
<i>Arizona elegans candida</i>	Mojave glossy snake				LC		
<i>Arizona elegans eburnata</i>	Desert glossy snake				LC		
<i>Arizona elegans occidentalis</i>	California glossy snake			SSC	LC		
<i>Bufo topiis rosaliae</i>	Baja California rat snake	X			LC		

<i>Chionactis occipitalis annulata</i>	Colorado shovel-nosed snake				LC
<i>Chionactis occipitalis occipitalis</i>	Mojave shovel-nosed snake				LC
<i>Chionactis occipitalis talpina</i>	Nevada shovel-nosed snake				LC
<i>Coluber constrictor mormon</i>	Western yellow-bellied racer				LC
<i>Contia longicauda</i>	Forest sharp-tailed snake				LC
<i>Contia tenuis</i>	Common sharp-tailed snake				LC
<i>Diadophis punctatus</i> "Coastal CA" ¹⁹	Ring-necked snake				LC
<i>Diadophis punctatus</i> "Eastern CA"	Ring-necked snake				LC
<i>Diadophis punctatus</i> "Southern CA"	Ring-necked snake	X			LC S
<i>Diadophis punctatus</i> "Great Basin" ²⁰	Ring-necked snake			SSC	LC
<i>Hypsiglena chlorophaea</i>	Northern desert night snake				LC
<i>Hypsiglena ochrorhyncha klauberi</i>	San Diego night snake				LC
<i>Hypsiglena ochrorhyncha nuchulata</i>	California night snake				LC
<i>Lampropeltis californiae</i>	Common kingsnake				LC
<i>Lampropeltis multifasciata</i> ²¹	California mountain kingsnake	X			LC S S
<i>Lampropeltis zonata</i>	California mountain kingsnake				LC S
<i>Masticophis flagellum piceus</i> ²²	Red coachwhip				LC
<i>Masticophis flagellum ruddocki</i>	San Joaquin coachwhip	X		SSC	LC
<i>Masticophis fuliginosus</i>	Baja California coachwhip			SSC	LC
<i>Masticophis lateralis euryxanthus</i>	Alameda striped racer	X	T	T	LC
<i>Masticophis lateralis lateralis</i>	California striped racer				LC
<i>Masticophis taeniatus</i>	Striped whipsnake				LC
<i>Phyllorhynchus decurtatus</i>	Spotted leaf-nosed snake				LC
<i>Pituophis catenifer affinis</i>	Sonoran gopher snake				LC
<i>Pituophis catenifer annectens</i>	San Diego gopher snake				LC

Taxon ¹	Common name	CDPG special animal	USFWS ²	CDFW ³	IUCN ⁴	USFS ⁵	BLM
Squamata—Snakes							
<i>Pituophis catenifer catenifer</i>	Pacific gopher snake				LC		
<i>Pituophis catenifer deserticola</i>	Great Basin gopher snake				LC		
<i>Pituophis catenifer pumilus</i>	Santa Cruz Island gopher snake	X			LC		
<i>Rhinocentillus lecontei</i>	Long-nosed snake				LC		
<i>Salvadora hexalepis hexalepis</i>	Desert patch-nosed snake				LC		
<i>Salvadora hexalepis mojavensis</i>	Mojave patch-nosed snake				LC		
<i>Salvadora hexalepis virgultea</i>	Coast patch-nosed snake	X		SSC	LC		
<i>Sonora semiannulata</i>	Western ground snake				LC		
<i>Tantilla hobartsmithi</i>	Southwestern black-headed snake				LC		
<i>Tantilla planiceps</i>	California black-headed snake				LC		
<i>Thamnophis atratus atratus</i>	Santa Cruz aquatic garter snake				LC		
<i>Thamnophis atratus hydrophilus</i>	Oregon aquatic garter snake				LC		
<i>Thamnophis atratus zaxanthus</i>	Diablo Range aquatic garter snake				LC		
<i>Thamnophis couchii</i>	Sierra (western aquatic) garter snake				LC		
<i>Thamnophis elegans elegans</i>	Mountain terrestrial garter snake				LC		
<i>Thamnophis elegans terrestris</i>	Coast terrestrial garter snake				LC		
<i>Thamnophis elegans vagrans</i>	Wandering terrestrial garter snake				LC		
<i>Thamnophis gigas</i>	Giant garter snake	X	T	T	V		
<i>Thamnophis hammondi</i>	Two-striped garter snake	X		SSC	LC	S	S
<i>Thamnophis marciatus</i>	Checkered garter snake						
<i>Thamnophis ordinoides</i>	Northwestern garter snake				LC		
<i>Thamnophis sirtalis fitchi</i>	Valley garter snake				LC		

<i>Thamnophis sirtalis infernalis</i> ²³	California red-sided garter snake	X		SSC ²⁴	LC	
<i>Thamnophis sirtalis tetralaenia</i>	San Francisco garter snake	X	E	E, FP	LC	
<i>Trimorphodon lambda</i>	Sonoran lyre snake					
<i>Trimorphodon lyrophanes</i>	Peninsular lyre snake					
Leptotyphlopidae						
<i>Rena humilis humilis</i> ²⁵	Southwestern blind snake				LC	
<i>Rena humilis cahuilae</i>	Desert blind snake				LC	
Viperidae						
<i>Crotalus atrox</i>	Western diamond-backed rattlesnake				LC	
<i>Crotalus cerastes cerastes</i>	Mojave Desert sidewinder				LC	
<i>Crotalus cerastes laterorepens</i>	Colorado Desert sidewinder				LC	
<i>Crotalus mitchellii</i>	Speckled rattlesnake				LC	
<i>Crotalus oreganus helleri</i> ²⁶	Southern Pacific rattlesnake				LC	
<i>Crotalus oreganus lutosus</i>	Great Basin rattlesnake				LC	
<i>Crotalus oreganus oreganus</i>	Northern Pacific rattlesnake				LC	
<i>Crotalus ruber</i>	Red diamond rattlesnake	X		SSC	LC	S
<i>Crotalus scutulatus</i>	Northern Mojave rattlesnake				LC	
<i>Crotalus stephensi</i>	Panamint rattlesnake				LC	

Testudines

Emydidae						
<i>Emys marmorata marmorata</i> ²⁷	Northern western pond turtle	X		SSC	V	S
<i>Emys marmorata pallida</i>	Southern western pond turtle	X		SSC	V	S S

(continued)

Taxon ¹	Common name	CDFG special animal	USFWS ²	CDFW ¹	IUCN ⁴	USFS ⁵	BLM
Testudines							
Kinosternidae							
<i>Kinosternon sonoriense</i>	Sonora mud turtle	X		SSC	V		
Testudinidae							
<i>Gopherus agassizii</i>	Mohave Desert tortoise	X	T	T	V		

1. Species, subspecies, or Distinct Population Segment (DPS)
 2. E: Endangered; T: Threatened.

3. E: Endangered; T: Threatened; FP: Fully Protect; SSC: Species of Special Concern.

4. E: Endangered; V: Vulnerable; NT: Near Threatened; LC: Least Concern; DD: Data Deficient

5. S: Sensitive.

6. Frost et al. (2006a) recommend placing all California bufonids except *Bufo alvarius* in the genus *Anaxyrus*. Frost et al. (2009b) recommend that *B. alvarius* be placed in the genus *Inhyla*.

7. Recuero et al. (2006a, 2006b) propose breaking *Pseudacris regilla* (sensu lato) into three distinct species. This proposal has not been widely accepted because the range boundaries of the three taxa are poorly characterized and significant haplotype sharing exists across these putative lineages that has not been studied.

8. This frog was widely introduced in California at one point, though presumed native populations were also present. The taxon may now be extirpated.

9. Frost et al. (2006a) recommend placing *Rana pipiens* and *R. muscosa* in the genus *Lithobates*.

10. It is likely that any populations on the eastern side of the Warner Mountains are actually *Rana boylei*. However, no specimens or data exist to clarify this issue. Until new data become available, *R. boylei* cannot be definitively included as a member of the California herpetofauna.

11. Following Rissler and Apodaca (2007).

12. An Owens Valley population was formerly presumed to be an undescribed taxon and has become widely recognized in the conservation community. Roviola (2010) refutes its status as a distinct lineage and we include the Owens Valley population with *Hydromantes platycephalus*.

13. Status applies only to Monterey County, CA, and south.

14. Papenfuss and Parham (2013) proposed splitting *Amniella pulchra* in California into five species.

15. Leach et al. (2009) revised the *Phrynosoma coronatum* complex, placing California populations of *P. coronatum* into *P. blainvillii*.

16. Schulte et al. (2006) propose that the *Sceloporus magister* subspecies be elevated to full species. This was refuted by Leach and Mulcahy (2007).

17. Leavitt et al. (2007) find a significant genetic structure within the *Xantusia vigilis* complex. Taxonomic revisions may occur in the near future within this clade.

18. Wood et al. (2008) divided the *rosy* bias into two species: *Lichanura orcutti* and *L. trivirgata*. Their mitochondrial data indicate that *L. trivirgata* is present in extreme southern California, though newer unpublished nuclear data suggest that the species break actually occurs farther south, in Baja California, Mexico (D. Wood, pers. comm.).

19. Feldman and Spicer (2006) and Fontanella et al. (2008) find evidence for lineages that are not concordant with

previously described subspecies boundaries. We follow the lineage designations from the latter study.

20. The Great Basin clade includes animals formerly assigned to *Diadophis punctatus regalis*. The SSC status refers only to populations occurring at isolated desert springs in Southern California.

21. Mountain kingsnake taxonomy is in flux. Rodriguez Robles et al. (1999b) refute the formerly recognized subspecies and find evidence for four distinct lineages. Myers et al. (2013) find evidence for two species (the arrangement that we follow here). *Lampropeltis multistriata* contains the former southern subspecies *Lampropeltis zonata parvifera* and *L. z. pulchra*. The conservation status applies to these two subspecies.

22. Nagy et al. (2004) propose combining *Masticophis* into the genus *Colebeles*.

23. Southern populations of this subspecies may represent a distinct taxon and are currently under study (C. Mahdi, pers. comm.; E. Ervin, pers. comm.).

24. SSC status applies to only the southern portion of the range.

25. Adalsteinsson et al. (2009) propose placing California *Leptotyphlops* in the genus *Rena*.

26. Some authors treat the subspecies of *Crotalus oregonus* as distinct species.

27. Some authors place the western pond turtles in the monotypic genus *Actinemys*. Spinks et al. (2014) recommend elevating both pond turtle subspecies to species status.

APPENDIX 2

Public Comment Announcement

We solicited public comment on this project by posting the announcement on the right on the websites of the following organizations: California Department of Fish and Wildlife, Center for North American Herpetology, Ecological Society of America (ECOLOG-L), Partners in Amphibian and Reptile Conservation, and The Wildlife Society. In addition, we circulated the announcement widely to colleagues via email. Following the public comment period, we also contacted experts on each taxon under consideration to request advice, data, and reviews of early drafts of this document.

California's list of Amphibian and Reptile Species of Special Concern (ARSSC) is a critical component of the management and protection of amphibians and reptiles in the state. The current California ARSSC list is undergoing a complete revision to better reflect those taxa that require some measure of conservation to stabilize populations and avoid future listing under the California Endangered Species Act. To date, the ARSSC revision team has developed a set of risk metrics, compiled a list of nominee taxa, and completed a preliminary risk assessment for each nominee based on literature reviews and locality information. Now, we need your help to make sure that we have the most accurate and complete list possible of SSC for potential inclusion in the final list. The best list will require input from as many knowledgeable biologists as possible. If you have data, well-documented field experience, or unpublished observations that are relevant to California's amphibian and reptile fauna, we invite you to share them with us.

Further details, risk assessments, and instructions for submitting feedback are available at <http://arssc.ucdavis.edu>. The public comment period closes August 31st, 2009.

*Bob Thomson
Amber Wright
Brad Shaffer*

*Center for Population Biology
University of California
Davis, CA 95616*

APPENDIX 3

Watch List

The watch list comprises taxa that were previously, but are no longer, considered Species of Special Concern. Here we include an explanation for each taxon's change in status and discuss future conservation concerns regarding Watch List taxa.

California tiger salamander (*Ambystoma californiense*)

Jennings and Hayes (1994a) identified this species as the highest-concern vernal pool-breeding amphibian in the state. In keeping with this assessment and recent research documenting its decline range-wide, *A. californiense* was listed under the California Endangered Species Act as a Threatened species in 2010, superseding Species of Special Concern status. See Bolster (2010) for the CDFW's recent status review. The species was also listed under the federal Endangered Species Act in 2000 (Santa Barbara; Endangered), 2003 (Sonoma; Endangered), and 2004 (Central; Threatened), as three separate Distinct Population Segments. Recent multi-locus phylogeographic work indicates that the Central Distinct Population Segment is composed of two separate lineages from the Inner Coast Range and Central Valley and that these may be best considered as separate units with different management needs (J. Johnson and B. Shaffer, unpublished data).

Orange-throated whiptail (*Aspidoscelis hyperythra*)

This taxon was included by Jennings and Hayes (1994a) primarily because of habitat loss within its

relatively narrow range. We place it on the Watch List because, thus far, it appears to tolerate habitat fragmentation better than many similarly distributed taxa, including the red diamond rattlesnake (*Crotalus ruber*), coast patch-nosed snake (*Salvadora hexalepis virgulata*), and California glossy snake (*Arizona elegans occidentalis*), all of which have experienced more severe declines; and it remains relatively common in many areas throughout its range. It is possible that further development and habitat fragmentation could cause more severe declines, so this taxon should be periodically reevaluated.

Baja California rat snake (*Bogertophis rosaliae*)

Jennings and Hayes (1994a) included the *B. rosaliae* primarily as a precaution. Virtually nothing was known about the species in California except that, if it ever naturally occurred in the state, it was probably rare and restricted in distribution (only a single specimen has ever been recorded). In the intervening time, no additional specimens have been reported, and no new information has become available for this species. If this species is found to be a native component of the California fauna, the conservation status should be reevaluated when more is known about the populations and habitat of the snake in California.

Yellow-blotched ensatina (*Ensatina eschscholtzii croceater*)

Jennings and Hayes (1994a) included this taxon primarily over concerns about land use changes within

its small range. We shared several of these concerns, although the severity of these threats appears to have decreased since 1994. As long as the planned preservation areas at Tejon Ranch remain in effect, a large amount of *E. e. croceater* habitat will remain protected, so designation as a Species of Special Concern may not be necessary. We include *E. e. croceater* on the Watch List to encourage reevaluation of habitat availability for this taxon in the future.

Large-blotched ensatina

(*Ensatina eschscholtzii klauberi*)

Jennings and Hayes (1994a) included this taxon primarily over concerns about ongoing development within its range. We agree that development has had, and is continuing to have, an impact on this species, although the severity of these impacts appears to be significantly less than those being experienced by other taxa with similar ranges. Further, the large-blotched *Ensatina* appears to be commonly found with stable populations throughout significant areas of its range, including protected parklands. If the extent of development increases within this salamander's range, it may become necessary to reconsider special concern status and more active management.

Mount Lyell web-toed salamander

(*Hydromantes platycephalus*)

This taxon was included by Jennings and Hayes (1994a) as a precaution, based on its patchy distribution and suspected susceptibility to local extirpations. We do not include *H. platycephalus* at this time because, although it is patchily distributed, the species appears to be stable throughout most of its range and is not experiencing appreciable risk from habitat disturbance (Wake and Papenfuss 2005). Additional populations have been found since the early 1990s, and the species appears to be relatively common at many sites. Although it is a California endemic, has a moderately small range, and is a narrow ecological specialist, this species does not appear to be currently at risk of immediate decline (Wake and Papenfuss 2005).

Owens Valley web-toed salamander

(*Hydromantes platycephalus*)

The Owens' Valley populations of *H. platycephalus* were included by Jennings and Hayes (1994a) as a precaution, both because little was known about the population biology of this elusive salamander and because it was strongly suspected that it was a distinct taxon. Research completed since 1994 suggests that these populations do not form a distinct lineage

but instead are part of the more broadly distributed *H. platycephalus* lineage (Rovito 2010). As with *H. platycephalus*, additional localities have been found and populations appear to be stable, leading us to conclude that Species of Special Concern designation is not required at the present time (Wake and Papenfuss 2005).

Southern California mountain kingsnakes

(*Lampropeltis zonata parvirubra* and *L. z. pulchra*)

The two southern California subspecies *L. z. parvirubra* and *L. z. pulchra* were considered Species of Special Concern by Jennings and Hayes (1994a) on the basis of suspected declines due to illegal collecting and habitat destruction from some collectors. We agree that this has occurred, although the current scale of exploitation does not appear to threaten this species' long-term survival. We placed the species on the Watch List in recognition that collection pressure and/or habitat destruction could cause the need to provide additional protections in the future.

Santa Cruz Island gopher snake

(*Pituophis catenifer pumilis*)

Jennings and Hayes (1994a) included this taxon primarily because of its small range (it is restricted to Santa Cruz and Santa Rosa islands) and threats from feral ungulates and pigs. We removed this species from special concern status because the invasive mammals causing the primary threats have been removed from the largest part of the range, Santa Cruz Island (USNPS 2010). This island is also well protected from future development because it is a national park.

Coronado skink

(*Plestiodon skiltonianus interparietalis*)

Jennings and Hayes (1994a) included *P. s. interparietalis* primarily because it has a relatively restricted range and has disappeared from some areas. As with *Aspidoscelis hyperythra*, we agree that some declines have occurred, although their severity appears to be modest. If these declines continue, further protections may be warranted in the future.

Del Norte salamander

(*Plethodon elongatus*)

Jennings and Hayes (1994a) included the Del Norte salamander because of concerns regarding habitat specialization by inland populations and the potential for timber harvest to destroy these habitats. Although these are valid concerns, as well as for two

close relatives of *P. elongatus*, the Scott Bar salamander (*Plethodon asupak*) and Siskiyou Mountains salamander (*P. stormi*), population status across most of the range of this taxon appears to be stable. Inland populations are patchy and likely more vulnerable to habitat degradation, which is why we place this taxon on our Watch List (H. Welsh, pers. comm.).

Mountain yellow-legged frogs

(*Rana muscosa* and *R. sierrae*)

Mountain yellow-legged frogs were designated as Species of Special Concern by Jennings and Hayes (1994a) under the name *R. muscosa*. Vredenburg et al. (2007) divided *R. muscosa* (sensu lato) into two species on the basis of morphometric measurements, differences in advertisement call, and mitochondrial DNA: the Sierra Madre yellow-legged frog (*R. muscosa*) in the south and the Sierra Nevada yellow-legged frog (*R. sierrae*) in the north. Both species were state listed in 2013, superseding Species of Special Concern status. See Bonham and Lockhart (2011) for the CDFW's recent status review of these taxa.

APPENDIX 4

Additional Taxa in Need of Research and Monitoring

We identified the following taxa that did not qualify for Species of Special Concern status but nonetheless would benefit from some level of additional research and/or monitoring. We provide a brief description of our concerns for each of these taxa below.

Orange-throated whiptail

(*Aspidoscelis hyperythra*)

Aspidoscelis hyperythra occurs in California in a relatively narrow region of southern California. Much of its available habitat has been destroyed or is threatened by ongoing urbanization and development. Further, many of the areas where habitat persists have become fragmented by development in intervening areas. The taxon remains locally common in several areas, although this should be reevaluated periodically. Further habitat modification could lead to more declines that warrant additional protections. Additional threats may arise from increasing intensity and/or frequency of wildfire in the region.

San Gabriel Mountains slender salamander

(*Batrachoseps gabrieli*)

Batrachoseps gabrieli occurs in a small area in Los Angeles and San Bernardino Counties (Stebbins 2003). Very few localities are known for this taxon, and its range is probably not fully characterized (Goodman et al. 1998, Hansen et al. 2005d). The salamander appears to be limited to talus slopes in the vicinity of oak, big cone spruce, and pine (Wake 1996, Goodman et al. 1998). It exhibits limited sur-

face activity and appears to specialize on an environment that is unlikely to be developed. This species' known range lies within the boundaries of the Angeles and San Bernardino National Forests and appears to be well protected at the present time. However, other narrowly distributed species of *Batrachoseps* have undergone large and unexplained declines, and it is possible that similar declines could occur for this species (Jennings and Hayes 1994a). For this reason, periodic monitoring and reevaluation of status of *B. gabrieli* is warranted.

Baja California rat snake

(*Bogertophis rosaliae*)

Bogertophis rosaliae is known only from a single road-killed specimen in California along Interstate 8 (specimen SDNHM 64416). It is unclear if this represents an escaped or discarded pet, a rare migrant from the known range farther south in Baja California, Mexico, or a regular, infrequently encountered component of the California reptile fauna. If a population does exist in California, ongoing development along the border in both the United States and Mexico is likely to isolate these populations from the main part of the range, which occurs farther south. If so, the California populations could be susceptible to stochastic effects associated with small populations, as well as habitat loss from development. In some areas this species appears to be associated with palm oases, which are uncommon habitat patches, so any degradation of this habitat may have severe impacts on the taxon.

If this species is native to California, it appears to be encountered exceedingly rarely and is never reported. Given this complete uncertainty concerning its status and validity as a native element of the California fauna, we place this taxon on the Watch List, primarily to highlight research needs. Surveys for this taxon should be encouraged, although in the absence of additional data, specimen collection should be strictly limited to only what is needed to learn more about its natural history and status within the state. However, we emphasize that tissue samples might help determine if any California specimens are native or introduced.

Yellow-blotched ensatina

(*Ensatina eschscholtzii croceater*)

Ensatina eschscholtzii croceater occurs in a relatively small area of Kern and Ventura Counties in southern California. Some localized populations may have undergone declines or extirpations due to development, although data on this are scarce. Workers have expressed concerns about land use practices and development in the Tehachapi Mountains, Bear Valley, Cummings Valley, and Tejon Ranch, particularly in areas of oak woodlands (pers. comm. in Jennings and Hayes 1994a). One of the main concerns for this taxon was that a large fraction of its range occurs on property owned by the Tejon Ranch Company, the largest contiguous private landholding in California, and that this land would be developed in a way that was incompatible with the salamander's survival. Since the previous evaluation, a large fraction of Tejon Ranch has been set aside for preservation—areas in which grazing, but not development, may continue (Tejon Ranch Conservancy 2008). In addition, many populations occur on National Forest and other public lands that are unlikely to experience intense habitat modification. The availability of suitable habitat should be monitored periodically, and habitat modification within its very restricted range should be avoided.

Southern California mountain kingsnake

(*Lampropeltis zonata* "Southern Clade" or *L. multifasciata*)

The southern clade of *L. zonata* includes the formerly recognized subspecies *L. z. pulchra* and *L. z. parvirubra* (Rodríguez-Robles et al. 1999h), and has more recently been recognized at the species level as *L. multifasciata* (Myers et al. 2013). This snake specializes on rocky outcrop habitats occurring primarily in a variety of woodland and chaparral habitats from sea level to nearly 3000 m (Stebbins 2003). It is

a popular species among herpetoculturists and collectors, and some have voiced concerns that habitat destruction has caused localized declines. Overzealous collection of this snake does tend to destroy the microhabitats within rocks, which can degrade the quality of sites for a long period of time, although the species exhibits a relatively narrow window of surface activity, and much of its habitat may be relatively inaccessible to collectors. Staub and Mulks (2009) surveyed the Mount Laguna region, San Diego County, from 2006 to 2008 and found that 75% of all rock piles surveyed had some degree of damage. They concluded that collecting is ongoing and is not restricted to the vicinity of roads, supporting the concerns that the intensity of ongoing collecting could harm this species. Managers should be wary of signs of habitat destruction, stemming from either collectors or other sources, particularly in areas that experience heavy human traffic such as Mount Laguna. If surveys demonstrate that these collecting activities are depleting populations, further management and enforcement of existing collecting prohibitions may be needed.

Del Norte salamander

(*Plethodon elongatus*)

Plethodon elongatus occurs from the California–Oregon border south into Humboldt and Trinity Counties. Optimal habitat for this taxon appears to be late-successional and mature forests, which may be increasingly impacted by timber harvest in the coming years (Welsh and Lind 1995; H. Welsh, pers. comm.). Prior to 2002, this species was managed under the Northwest Forest Plan (Welsh and Bury 2005, Survey and manage program 2010). These protections have now been removed, although much of the habitat that supports this taxon remains protected under the Plan (Northwest Forest Plan 1994). Monitoring efforts should focus on the impact of timber harvest on this species' ability to persist, particularly at inland sites.

Western black-headed snake

(*Tantilla planiceps*)

The natural history of *T. planiceps* is poorly understood in California. We have almost no information concerning this species' natural history, habitat requirements, or population densities. The snake seems to be patchily distributed and rarely seen, making the detection of population declines or extirpations difficult. In addition, much of its range occurs in areas that have experienced heavy development and habitat modification. Some workers have

suggested that changing wildfire regimes in southern California could be having a negative impact on this species; however, relevant data are very sparse. An important priority for this taxon is an increased research effort focused on distribution and habitat surveys so that its ecological requirements and population dynamics can be better characterized. As populations are discovered, tissue samples should be collected for molecular analyses of the degree of isolation and differentiation of these apparently disjunct populations.

Baja California night lizard

(*Xantusia wigginsi*)

Xantusia wigginsi was not known to be a part of the California lizard fauna until recent genetic studies established its presence in extreme southern California (Leavitt et. al. 2007). Virtually nothing is known about this taxon's range, life history, habitat requirements, or conservation status within California. Further research on this species is needed before assessments of its conservation status and management needs can be made.

This page intentionally left blank

GLOSSARY

- ADRESSED LIMBS** Position of the limbs such that the forelimbs are pressed backwards against the trunk of the animal, and the hind limbs are pressed forward against the trunk. The distance between addressed limbs, a character which measures the relative limb length with respect to the trunk length, is usually best measured in preserved specimens, since the limbs may be damaged in living animals.
- ALLOPATRIC** Occurring in separate areas; refers to species ranges that do not overlap.
- ALLOZYME** Alleles of an enzyme that vary in their speed of migration through an electrophoretic gel. A common way to quantify genetic variation before DNA sequencing became routine.
- AMPLEXUS** Mating behavior in many aquatic anurans and some salamanders in which the male grasps the female with the front legs.
- BD** *Batrachochytrium dendrobatidis*. A pathogenic fungus that causes the disease chytridiomycosis in many amphibians.
- CARAPACE** The dorsal half of a turtle shell.
- COSTAL GROOVES** Lateral indentations along the trunk of many salamanders.
- CRITICAL THERMAL MAXIMUM** The temperature above which a given species ceases to be able to maintain normal body function. Extended temperatures above this point generally lead to death.
- CRITICAL THERMAL MINIMUM** The temperature below which a given species ceases to be able to maintain normal body function. Extended temperatures below this point generally lead to death.
- CRYPTIC TAXA** Evolutionarily distinct lineages that are morphologically conserved and are difficult to distinguish from one another on the basis of morphology alone.
- DIAPAUSE** A delay in the life cycle of an organism, often occurring in response to adverse environmental conditions.
- DORSOLATERAL FOLDS** Ridges of the skin that run along either side of the back in many frogs.
- EXTANT** A taxon that is still in existence, opposite of extinct.
- HIBERNACULUM** A place used by one or more individuals to hibernate or undergo a period of dormancy. Frequently used to refer to areas that house many hibernating individuals of the same species, especially sites that are used repeatedly over many years. The plural is hibernacula.
- INTROGRESSION** Transfer of genetic molecules from one species to another. In our usage, this most commonly refers to the transfer of the mitochondrial genome among species due to hybridization.
- ISOLATION BY DISTANCE** The genetic signature that tends to arise from the tendency of individuals within a population to mate with nearby

- individuals, eventually leading to the gradual accumulation of genetic differentiation across the landscape.
- KEELED** A spine or ridge structure that runs along the central axis of a scale or scute.
- LATE-SERAL** Used to describe forests that are in a later stage of succession. Typified by the presence of large, old (>100 years) trees in the overstory.
- MICROSATELLITE** Short repetitive regions in the DNA that often exhibit a large amount of variation due to the very high rate of mutation in these regions of the genome. Frequently employed to measure population genetic variation within species, because their high mutation rate allows them to track changes in gene flow and population size quickly.
- MTDNA** An abbreviation for mitochondrial DNA, the separate chromosome found in the mitochondria of all plants and animals. Until recently, it has been the standard molecule of choice for most systematic, population genetic, and phylogeographic research.
- NASOLABIAL GROOVES** Characteristic grooves that run from each naris (external nostril) down to the upper lip in plethodontid salamanders.
- NUCHAL** Relating to or lying in the region of the nape.
- OCELLUS** An eye-like spot.
- OVIPAROUS** A mode of reproduction in which embryos develop inside of eggs.
- OVOVIVIPAROUS** A mode of reproduction in which embryos develop inside of eggs which are retained in the mother's body until hatching.
- PAEDOMORPHOSIS** The retention of larval traits into adulthood. In ambystomatid and dicamptodontid salamanders, it is also used to refer to reproduction in the larval condition.
- PARAPHYLETIC** A group of taxa, all descending from of a common ancestor, that does not contain all descendants of that ancestor. For examples, "reptiles" as traditionally defined are paraphyletic because they do not contain birds as a contained taxon.
- PARATOID GLANDS** External skin glands that lie along the back of the head or neck region and are prominent in most toads and several species of salamander.
- PCA** Principle component analysis. A multivariate ordination approach that reduced the variability among large sets of measured variables down to a (usually) smaller number of independent (orthogonal) variables.
- PIT TAG** Passive integrated transponder tag. A small injectable tag that emits a unique electronic signal that can be read using specialized instruments. A frequently used method for uniquely labeling individual organisms in a population.
- PLASTRON** The ventral part of a turtle shell.
- POLYTYPIC** Having several morphological forms. These may or may not correspond to evolutionary lineages.
- POND TYPE LARVAE** Salamander larvae that develop in ponds are characterized by having relatively large long fins associated with a relatively strong swimming ability.
- SCUTE** An enlarged scale, such as those on a turtle shell.
- SCL** Straight carapace length. The distance from the anterior to the posterior end of the carapace taken along the midline and measured as a straight distance (i.e., not measuring along the curvature of the shell). A standard way of measuring body length in turtles.
- SNP** Single nucleotide polymorphism. A homologous nucleotide position in a DNA sequence that is variable among conspecific individuals. SNPs are increasingly used instead of allozymes, microsatellites, and mtDNA for population genetic and species delimitation studies.
- STREAM TYPE LARVAE** Salamander larvae that develop in streams are typically smaller than pond type larvae and have smaller tail fins. Behaviorally, they tend not to swim in the open water and instead remain near the substrate.
- SVL** Snout to vent length. The distance from the tip of the snout to the anterior edge of the cloaca. A standard way of measuring length in many amphibians and reptiles.
- TL** Total length. The distance from the tip of the snout to the end of the tail.
- VIVIPAROUS** A mode of reproduction in which females give birth to live young that are not retained in shelled eggs (compare with ovoviviparous).

INDEX

Note: Page number followed by (f), (m), and (t) indicates figure, map, and table respectively.

- agricultural runoff, 116, 122
- agrochemicals, 75, 110
- Ambystoma californiense*, 38, 42, 368
- Ambystoma macrodactylum sigillatum*, 136–41, 136(f)
 - breeding activity and egg laying, 138
 - breeding season, 139
 - courtship, pattern of, 138
 - diet of, 139
 - disease and environmental contaminants, 140
 - distribution of, 137(m), 139–40
 - effects of climate change on, 140
 - habitat requirements for, 139
 - hibernation, 138
 - identification of, 136–38
 - in Klamath–Siskiyou bioregion, 139
 - life history of, 138–39
 - management of, 141
 - metapopulation dynamics, 141
 - migration to breeding habitat, 138
 - monitoring, research, and survey needs, 141
 - as Priority 2 Species of Special Concern, 136
 - ranavirus infections, 140
 - recolonization of, 140
 - reproduction, 138
 - risk factors, 136(t)
 - in spring-fed water bodies, 139
 - status of, 136, 141
 - susceptibility to local extirpations, 139
 - taxonomy of, 138
 - threats faced by, 140
 - trends in abundance, 140
 - upland habitat, 139
- amphibians and reptiles, management of, 39–43
 - Amphibian and Reptile Conservation Inventory and Monitoring guide, 43
 - in California, 355(t)–366(t)
 - mitigating effects of roads, 41–42
 - protection of
 - aquatic habitats, 39–40
 - terrestrial habitat patches, 40–41
 - translocation of animals, 42–43
- Aneides flavipunctatus niger*, 142–46, 142(f)
 - allozyme studies of, 144
 - breeding activity and egg laying, 144
 - distribution of, 143(m), 144–45
 - effects of climate change on, 145
 - habitat requirements, 144
 - identification of, 142–44
 - life history of, 144
 - management of, 145
 - mitochondrial DNA analysis, 144
 - monitoring, research, and survey needs, 145–46
 - as Priority 3 Species of Special Concern, 142
 - risk factors, 142(t)
 - status of, 142, 145
 - streamside microhabitats, 144
 - taxonomy of, 144
 - threats faced by, 145
 - trends in abundance, 145
- Anniella pulchra*, 186–91, 186(f)
 - breeding season, 188
 - climate change, impact of, 190
 - distribution of, 187(m), 189
 - feeding ecology of, 188

- genetic markers, 191
- gestation period, 188
- habitat requirements, 188–89
- habitat suitability for, 189
- human-disturbed habitats, 190
- identification of, 186–88
- life history of, 188
- management of, 190
- mating activity, 188
- microhabitat suitability, 189
- mitochondrial DNA analysis, 188
- monitoring, research, and survey needs, 190–91
- movement ecology, 188
- nuclear DNA analysis, 188
- as Priority 2 Species of Special Concern, 186
- risk factors, 186(t)
- sexual maturity, 188
- status of, 186, 190
- surface activity, 188
- taxonomy of, 188
- threats faced by, 189–90
- trends in abundance, 189
- Anza-Borrego Desert State Park, 248–49
- Apalone spinifer*, 302, 307
- aquatic habitats, in California, 31
 - acidification of, 73
- aquatic invasive predators, 31
- aquatic invertebrates, 307
- Argentine ant. See *Linepithema humile*
- Arizona elegans candida*, 257
- Arizona elegans eburnata*, 257
- Arizona elegans occidentalis*, 255–59, 255(f), 368
 - breeding activity and egg laying, 257
 - coloration of, 255
 - distinctiveness of, 257
 - distribution of, 256(m), 258
 - ecological specialization and endemism, 258
 - habitat requirements, 257–58
 - identification of, 255
 - life history of, 257
 - management of, 259
 - monitoring, research, and survey needs, 259
 - pitfall trapping of, 258
 - population sizes, 259
 - prey of, 257
 - as Priority 1 Species of Special Concern, 255
 - reproductive activity, 257
 - risk factors, 255(t)
 - status of, 255, 258–59
 - taxonomy of, 257
 - threats faced by, 258
 - trends in abundance, 258
- Arroyo toad. See *Bufo californicus*
- Ascaphus truei*, 51–58, 51(f)
 - breeding behavior, 53
 - climate change, impact of, 55
 - clutch size, 53
 - distribution of, 52(m), 55
 - habitat requirements, 53–55
 - identification, 51–53
 - larvae of, 51, 55
 - life history, 53
 - management recommendations, 57
 - marijuana cultivation and, 57
 - monitoring, research, and survey needs, 57–58
 - population decline, 55
 - risk factors, 51(t)
 - road construction, impact of, 56
 - sensitivity to warm temperatures, 54
 - sexual maturity of, 53
 - status determination, 57
 - status summary, 51
 - in streams, 54
 - taxonomic relationships, 53
 - thermal tolerances of, 54
 - threats faced by, 55–57
 - timber harvesting, impact of, 56
 - time to metamorphosis, 53
 - trends in abundance, 53
- Aspidoscelis hyperythra*, 368–69, 371
 - habitat destruction, impact of, 371
- Aspidoscelis tigris stejnegeri*, 192–96, 192(f)
 - distribution of, 193(m), 195
 - habitat loss and fragmentation, 195
 - habitat requirements for, 194–95
 - identification of, 192–94
 - life history of, 194
 - management of, 196
 - monitoring, research, and survey needs, 196
 - population size, 196
 - as Priority 2 Species of Special Concern, 192
 - reproductive activity, 194
 - risk factors, 192(t)
 - status of, 192, 196
 - taxonomy of, 194
 - threats faced by, 195
 - trends in abundance, 195
- assisted migration, importance of, 231
- Association of Fish and Wildlife Agencies, 44
- Baja California coachwhip. See *Masticophis fuliginosus*
- Baja California night lizard. See *Xantusia wigginsi*
- Baja California Peninsula, 300
- Baja California rat snake. See *Bogertophis rosaliae*
- Batrachochytrium dendrobatidis* (Bd), 45, 68, 73, 82, 91, 96–99, 104, 116–17, 122, 140–41, 184, 375
- Batrachoseps campi*, 147–50, 147(f)
 - distribution of, 148(m), 149
 - elevational range of, 149
 - habitat requirements for, 149

- identification of, 147–49
- life history of, 149
- management of, 150
- monitoring, research, and survey needs, 150
- as Priority 3 Species of Special Concern, 147
- risk factors, 147(t)
- status of, 147, 150
- surface activity of, 149
- taxonomy of, 149
- threats faced by, 149–50
- trends in abundance, 149
- Batrachoseps gabrieli*, 371
- Batrachoseps minor*, 151–55, 151(f)
 - distribution of, 152(m), 153
 - DNA sequences, 154
 - effect of invasion of exotic plants on, 154
 - habitat modification, 154
 - habitat requirements for, 153
 - identification of, 151–53
 - life history of, 153
 - management of, 154
 - mitochondrial DNA analysis, 153
 - monitoring, research, and survey needs, 154–55
 - as Priority 1 Species of Special Concern, 151
 - risk factors, 151(t)
 - sensitivity to habitat, 154
 - status of, 151, 154
 - taxonomy of, 153
 - threats faced by, 153–54
 - trends in abundance, 153
- Batrachoseps relictus*, 156–60, 156(f)
 - aquatic microhabitats of, 159
 - distinctiveness of, 158
 - distribution of, 157(m), 159
 - effects of climate change on, 159–60
 - habitat degradation from road construction, 159
 - habitat requirements for, 159
 - identification of, 156–58
 - life history of, 158–59
 - management of, 160
 - mitochondrial DNA analysis, 158
 - monitoring, research, and survey needs, 160
 - as Priority 1 Species of Special Concern, 156
 - risk factors, 156(t)
 - status of, 156, 160
 - surface activity of, 159
 - taxonomy of, 158
 - threats faced by, 159–60
 - trends in abundance, 159
- Bogertophis rosaliae*
 - decline in population of, 368
 - habitat of, 368
 - destruction of, 371
 - status of, 368, 372
- Breckenridge Mountain, 158–60
- Breeding Bird Surveys, 47
- Bufo alvarius*, 59–63, 59(f)
 - breeding behavior, 61, 63
 - distribution of, 60(m), 61–62
 - effect of
 - landscape modification, 62
 - pesticide applications, 62
 - habitat requirements, 61
 - identification, 59–61
 - life history, 61
 - management recommendations, 62–63
 - metapopulation dynamics, 63
 - monitoring, research, and survey needs, 63
 - osteological and genetic data, 61
 - poison of, 61
 - toxicity of, 61
 - population decline, 62
 - causes of, 62
 - as Priority 1 Species of Special Concern, 59
 - risk factors, 59(t)
 - status determination, 62
 - status of, 59, 62
 - taxonomy of, 61
 - threats faced by, 62
 - trends in abundance, 62
- Bufo californicus*, 64–68, 64(f)
 - Anaxyrus* genus, 66
 - breeding behavior, 66
 - distribution of, 65(m), 67
 - drought and El Niño cycles, effects of, 68
 - feeding habits, 66
 - in human-made habitats, 67
 - identification of, 64–66
 - larvae of, 66–67
 - life history of, 66–68
 - management recommendations, 68
 - monitoring, research, and survey, 68
 - mortality rate of, 67
 - nocturnal activity, patterns of, 66–67
 - nonnatural disturbances, 66
 - population size and connectivity, 68
 - predation pressure, 67
 - as Priority 1 Species of Special Concern, 64
 - radio tracking of, 67
 - reproductive season, 66
 - risk factors, 64(t)
 - seasonal activity period for, 66
 - sexual maturity, 66
 - status of, 64, 68
 - stream and riparian habitat, 67
 - survivorship of, 66–68
 - taxonomy of, 66
 - threats faced by, 68
 - trends in abundance, 67–68
 - USFWS recovery plan for, 68

- Bufo ranorus*, 69–76, 69(f)
 breeding activity and egg-laying, 71–72
 breeding choruses, 71
 human disturbance of, 73–75
 breeding season, 75
 disease outbreaks, 73, 75
 distribution of, 70 (m), 72–73
 drought, effects of, 74
 embryos and hatchlings, 73
 genetic differentiation, 71
 genetic distance between populations, 71
 habitat modification, impact of, 74
 habitat requirements, 72
 hibernation and breeding biology, 75
 identification of, 69–71
 insect food supply for, 74
 larvae of, 72
 life history of, 71–72
 livestock grazing, influence of, 74
 management program, 75
 metamorphosis of, 71
 metapopulation dynamics, 74
 monitoring, research, and survey needs, 75–76
 mortality rate, 73
 palatability of, 74
 population decline, 72–73
 causes of, 73, 75
 predators of, 74
 as Priority 1 Species of Special Concern, 75
 protection from predation, 72
 risk factors, 69(t)
 seasonal activity period, 72
 sexual maturity of, 75
 in snowmelt pools, 73
 status of, 69, 75
 stress and immunosuppression, 75
 taxonomy of, 71
 threats faced by, 73–75
 trends in abundance, 73
- Bureau of Land Management, 5
- California Aquatic Invasive Species Management Plan, 40
- California Climate Action Team, 44
 assessments of, 11
- California Coastal Act (1976), 83
- California Department of Fish and Wildlife (CDFW), 4, 6, 19
 Biogeographic Information and Observation System, 16
 Geographic Information System, 16
- California Endangered Species Act, 4, 6, 8, 14, 19, 33, 231, 368
- California Environmental Quality Act, 5, 36
- California Essential Habitat Connectivity Project, 41
- California Floristic Province, 4
- California giant salamander. See *Dicamptodon ensatus*
- California glossy snake. See *Arizona elegans occidentalis*
- California kangaroo rat. See *Dipodomys californicus*
- California legless lizard. See *Anniella pulchra*
- California Natural Community Conservation Planning, 41
- California Natural Diversity Database, 5, 13, 15, 33, 216
- California red-legged frog. See *Rana draytonii*
- California tiger salamander. See *Ambystoma californiense*
- California Wildlife Habitat Relationships, 16–17
- cannibalism, 86, 133, 163, 183
- carbon emissions, 11, 44
- Carson River, 300
- Cascades frog. See *Rana cascadae*
- Christmas Bird Counts, 47
- chytrid fungus. See *Batrachochytrium dendrobatidis*
- chytridiomycosis, 73, 91, 122
- citizen science, 47
- climate change
 conservation risk due to, 39
 impacts of, 44–45, 302
 vulnerability to, 11, 11(t)
- Coachella Valley, 229
 Multiple Species Habitat Conservation Plan, 232
- coastal tailed frog. See *Ascaphus truci*
- coastal whiptail. See *Aspidoscelis tigris stejnegeri*
- coast horned lizard. See *Phrynosoma blainvillii*
- coast patch-nosed snake. See *Salvadora hexalepis virgulata*
- coast range newt. See *Taricha torosa*
- Coleonyx variegatus abboti*, 197–201, 197(f)
 distinctiveness of, 199
 distribution of, 198(m), 199–200
 habitat degradation, 200
 habitat requirements, 199
 identification of, 197–99
 impact of
 artificial night lighting, 200
 climate change, 200
 development and agriculture, 200
 life history of, 199
 management of, 200
 monitoring, research, and survey needs, 200–201
 nuclear DNA markers, 199
 pitfall trapping, 199, 201
 as Priority 3 Species of Special Concern, 197
 reproductive activity, 199
 risk factors, 197(t)
 status of, 197, 200
 surface activity, 199
 taxonomy of, 199
 threats faced by, 200
 trends in abundance, 200
- Colorado Desert, 35
- Colorado Desert fringe-toed lizard. See *Uma notata*
- Colorado River, 307, 308

- common garter snake. See *Thamnophis sirtalis*
- common musk turtle. See *Sternotherus odoratus*
- Cope's leopard lizard. See *Gambelia copeii*
- Coronado skink. See *Plestiodon skiltonianus interparietalis*
- Couch's spadefoot. See *Scaphiopus couchii*
- counties and islands, of California, 50(m)
- Crotalus ruber*, 260–65, 260(f), 368
 - climate change, impact of, 264
 - coloration and behavior, 260
 - courtship and mating, 262
 - denning behavior, 262
 - distribution of, 261(m), 263–64
 - gestation period, 262
 - habitat requirements, 263
 - identification of, 260
 - life history of, 262–63
 - management of, 264
 - monitoring, research, and survey needs, 264–65
 - neonates of, 260
 - pitfall trapping, 265
 - prey of, 263
 - as Priority 3 Species of Special Concern, 260
 - radio tracking of, 262
 - reproductive activity, 262
 - risk factors, 260(t)
 - sexual maturity, 263
 - status of, 260, 264
 - taxonomy of, 262
 - threats faced by, 264
 - trends in abundance, 264
- Dale Dry Lake, San Bernardino County, 242
- Daphnia magna*, 45
- DDT pesticide, 116
- Death Valley, Inyo County, 269
- Del Norte salamander. See *Plethodon elongatus*
- desert glossy snake. See *Arizona elegans eburnata*
- desert mule deer. See *Odocoileus hemionus crooki*
- Diadophis punctatus regalis*, 266–70, 266(f)
 - breeding activities and egg laying, 268
 - climate changes, impact of, 269
 - denning activities, 268
 - distribution of, 267(m), 269
 - habitat requirements, 269
 - identification of, 266
 - life history of, 268–69
 - management of, 269–70
 - mitochondrial DNA, 268
 - monitoring, research, and survey needs, 270
 - prey of, 269
 - reproductive activity, 268
 - risk factors, 266(t)
 - risk of extirpation, 269
 - status of, 266, 269
 - taxonomy of, 268
 - threats faced by, 269
- trends in abundance, 269
- Dicamptodon ensatus*, 161–65, 161(f)
 - breeding activity and egg laying, 163
 - breeding and larval development, 163
 - breeding season, 163
 - cannibalism behavior, 163
 - distinctiveness of, 163
 - distribution of, 162(m), 163–64
 - effects of
 - climate change, 164
 - wildfire, 164
 - fragmentation of riparian habitat, 164
 - genetic markers, 163
 - habitat requirements for, 163
 - identification of, 161–63
 - larvae of, 163
 - habitat of, 163
 - in slow-moving water, 163
 - life history of, 163
 - management of, 165
 - monitoring, research, and survey needs, 165
 - mortality and habitat degradation, 164
 - paedomorphosis, prevalence of, 163
 - pitfall trapping, 165
 - as Priority 3 Species of Special Concern, 161, 164
 - risk factors, 161(t)
 - status of, 161, 164
 - taxonomy of, 163
 - threats faced by, 164
 - trends in abundance, 164
- Dipodomys californicus*, 221
- disease transmission, 42, 45
 - in *Bufo canorus*, 73, 75
- DNA sequencing, 110, 154, 252, 277
- East Bay Regional Park District, 103
- Elgaria panamintina*, 202–6, 202(f)
 - distribution of, 203(m), 205
 - habitat requirements, 205
 - identification of, 202–4
 - life history of, 204–5
 - management of, 205–6
 - monitoring, research, and survey needs, 206
 - predators of, 205
 - as Priority 3 Species of Special Concern, 202
 - reproductive activity, 204
 - risk factors, 202(t)
 - status of, 202, 205
 - taxonomy of, 204
 - thermal tolerances, 204
 - threats faced by, 205
 - trends in abundance, 205
- El Niño cycles, 68
- Emys marmorata*, 296–303, 296(f)
 - clutch sizes, 299
 - diet, 299

- distribution of, 297(m), 300–301
- growth rates of, 299
- habitat loss of, 301
- habitat requirements of, 299, 300, 303
- hatchlings of, 299
- human-modified habitats, impact of, 303
- identification of, 298
- introduced species, impact of, 308
- life history of, 299
- management recommendations, 302–3
- mating season, 299
- melanistic, 298
- monitoring, research, and survey needs, 303
- movement to overwintering sites, 299
- nest sites, 299, 303
- population sizes, 301
- predators of, 302, 303
- reproductive maturity, 299–300
- risk factors, 298(t)
- status, 298, 302
- taxonomy, 298–99
- threats faced by, 301–2
- trends in abundance, 301
- endemic and near endemic species, 33(t)
- Ensatina eschscholtzii croceator*, 368–69, 372
- environmental contamination, 73, 97–98, 110, 140
- environmental impact assessment, 36

- Fallen Leaf Lake, 109
- Federal Endangered Species Act, 8, 244
- fishbait industry, 109
- flat-tailed horned lizard. *See Phrynosoma mcallii*
- Flat-tailed Horned Lizard Interagency Coordinating Committee, 229, 231–32
- foothill yellow-legged frog. *See Rana boylei*
- forest ecosystems, 11

- Gambelia copeii*, 207–11, 207(f)
 - breeding season, 209
 - distribution of, 208(m), 210
 - gene flow and hybridization, 209
 - habitat loss due to development, 210
 - habitat requirements, 209–10
 - hibernation, 209
 - identification of, 207–9
 - life history of, 209
 - management of, 210
 - monitoring, research, and survey needs, 211
 - risk factors, 207(t)
 - sexual size dimorphism, 209
 - as Species of Special Concern, 207
 - status of, 207, 210
 - taxonomy of, 209
 - threats faced by, 210
 - trends in abundance, 210
- gene flow, isolation-by-distance model of, 95
- generalist predator, 44
- Gila monster. *See Heloderma suspectum*
- glyphosate, 110
- granite night lizard. *See Xantusia henshawi*
- Great Basin and Sonoran ecoregions, 35
- Greenhorn Mountains, 158–59

- Habitat Conservation Plans (HCP), 5, 31
- habitat corridors, 41–42, 58, 185, 196, 201, 211, 231, 245, 253, 264
- habitat fragmentation, 31, 35, 41, 195–96, 233, 243–44, 282–83, 368
- habitat loss and degradation, 34, 81, 171, 184, 189, 253
- habitat quality for species, determinants of, 231
- habitat restoration, 68, 92, 98–99, 108, 110, 135, 150, 190, 196, 231, 244, 290, 294–95, 303
- Havas National Wildlife Refuge, 308
- headstarting programs, 43, 303, 308
- Heloderma horridum*, 214
- Heloderma suspectum*, 212–17, 212(f)
 - daily activity pattern, 215
 - diet of, 215
 - distribution of, 213(m), 215–16
 - foraging behavior, 215
 - habitat requirements, 215
 - homing ability, 215
 - identification of, 214
 - life history of, 214–15
 - male–male combat, 215
 - management of, 216
 - monitoring, research, and survey needs, 216–17
 - nocturnal activity, 215
 - reproductive activity, 215
 - riparian or xeroriparian habitat, 215
 - risk factors, 214(t)
 - sexual maturity, 215
 - as Species of Special Concern, 214
 - status of, 214, 216
 - subspecies of, 214
 - taxonomy of, 214
 - threats faced by, 216
 - trends in abundance, 216
 - venom, use of, 215
- herpetofauna, 42, 259
- hibernation
 - of *Ambystoma macrodactylum sigillatum*, 138
 - of *Bufo canorus*, 75
 - of *Gambelia copeii*, 209
 - of *Kinosternon sonoriense*, 306
 - of *Phrynosoma mcallii*, 227
 - of *Rana pretiosa*, 114
- Humboldt Bay National Wildlife Refuge, 79
- Humboldt County General Plan, 81, 171
- Hydromantes platycephalus*, 39, 369

- Imperial Valley, 62
- introduced species, impact of, 4, 83, 104, 133, 183–84, 190, 232, 301–2, 308
- invasive species, 36, 40, 89, 134, 184–85, 303
- Inyo Mountains salamander. See *Batrachoseps campi*
- Kathleen, Hurricane, 122
- Kaweah River, 182
- Kern River Canyon, 158–59
- Kinosternon sonoriense*, 304–9, 304(f)
- clutch sizes, 306
 - decline of, 308
 - diet of, 306
 - distribution of, 305(m), 307
 - feeding habits of, 306
 - habitat loss of, 309
 - habitat requirements, 306–7
 - headstarting program, 308
 - hibernation of, 306
 - identification of, 304–6
 - life history of, 306
 - management recommendations, 308
 - monitoring, research, and survey needs, 308–9
 - population densities, 306
 - reproductive maturity, 306
 - risk factors, 304(t)
 - status of, 304, 308
 - taxonomy of, 306
 - terrestrial movements, 306
 - threats faced by, 308
 - trends in abundance, 307–8
- Klamath–Siskiyou bioregion, 139
- Klauher, L.M., 68
- Laguna Dam, 308
- Lampropeltis multifasciata*, 372
- Lampropeltis zonata parvirubra*, 369, 372
- Lampropeltis zonata pulchra*, 369, 372
- landscape genetics, 45, 146
- land use, conversion to agriculture, 302
- Lassen Volcanic National Park, 140
- lesser slender salamander. See *Batrachoseps minor*
- Linepithema humile*, 221–23
- livestock grazing, influence of, 74
- Lower Colorado River, 121–23, 216
- lowland leopard frog. See *Rana yavapaiensis*
- Lucas Creek, 159
- marijuana cultivation, 35, 55–58, 81, 83, 90–91, 170–72
- Masticophis flagellum ruddocki*, 271–74, 271(f)
- climate change, impact of, 274
 - denning activity, 273
 - diet of, 273
 - distribution of, 272(m), 273
 - habitat requirements for, 273
 - identification of, 271
 - life history of, 273
 - management of, 274
 - mating season, 273
 - mitochondrial DNA analysis, 273
 - monitoring, research, and survey needs, 274
 - mortality of, 274
 - prey of, 273
 - as Priority 2 Species of Special Concern, 271
 - risk factors, 271(t)
 - status of, 271, 274
 - surface activity, 273
 - taxonomy of, 273
 - threats faced by, 273–74
 - trends in abundance, 273
- Masticophis fuliginosus*, 275–78, 275(f)
- climate change, impact of, 278
 - diet of, 277
 - distinctiveness of, 275
 - distribution of, 276(m), 277
 - diurnal behavior, 277
 - foraging behavior, 277
 - habitat requirements for, 277
 - hatchlings of, 277
 - home range sizes, 277
 - identification of, 275–77
 - life history of, 277
 - management of, 278
 - mating activity, 277
 - mitochondrial DNA analysis, 277
 - monitoring, research, and survey needs, 278
 - prey of, 277
 - as Priority 3 Species of Special Concern, 275
 - radiotelemetry of, 277
 - risk factors, 275(t)
 - status of, 275, 278
 - taxonomy of, 277
 - threats faced by, 278
 - trends in abundance, 278
- Mattole Watershed, 54–55, 88, 169–70, 177
- McCloud River, 169
- Mendocino National Forest, 80
- Mesquite Mountains, 216
- metapopulation dynamics, 31, 41, 63, 74–75, 98, 134, 141, 185, 238
- Mexican beaded lizard. See *Heloderma horridum*
- migration of species
- assisted migration, 231
 - corridors for (See habitat corridors)
 - for ensuring gene flow, 231
 - migration tunnels, 42
- Mojave Desert, 3, 34–35, 149, 189, 222, 236, 269, 273
- Mojave fringe-toed lizard. See *Uma scoparia*
- Mojave glossy snake. See *Arizona elegans candida*

- Mojave River, 300
- Monterey peninsula, 186
- mortality, road-associated, 41–42
- mountain yellow-legged frogs. See *Rana muscosa*;
Rana sierrae
- Mount Laguna, 372
- Mount Lyell web-toed salamander. See *Hydromantes platycephalus*
- Natural Community Conservation Plan (NCCP), 5, 31, 196, 232
- nominee taxa, scoring of, 12
- nonnative plant species, 233
- northern leopard frog. See *Rana pipiens*
- northern red legged frog. See *Rana aurora*
- northern western pond turtle. See *Emys marmorata*
- North Fork Feather River, 90
- Northwest Forest Plan, 372
- nuclear DNA markers, 105, 110, 199, 277
- Ocotillo Wells State Vehicular Recreation Area, 227–28, 230
- Odocoileus hemionus crooki*, 63
- Old-growth Wildlife Project, 169
- orange-throated whiptail. See *Aspidoscelis hyperythra*
- Oregon spotted frog. See *Rana pretiosa*
- oviparous reptiles, 204, 281
- Owens River, 109
- Owens Valley, 39, 109
- web-toed salamander (See *Hydromantes platycephalus*)
- Pacific treefrog. See *Pseudacris regilla*
- Panamint alligator lizard. See *Elgaria panamintina*
- Peninsula Open Space Trust, 145
- peninsular leaf-toed gecko. See *Phyllodactylus nocticolus*
- Pepperwood Creek, 179
- Phrynosoma blainvillii*, 218–24, 218(f)
- agriculture and development, impact of, 222
- breeding season, 220
- cattle grazing and, 224
- climate change, impact of, 223
- deaths due to predation, 221
- diets of, 223
- distribution of, 219 (m), 221–22
- diurnal activity, 220
- effects of wildfire on, 223, 224
- habitat requirements, 221
- identification of, 218–19
- life history of, 220–21
- management of, 223
- microhabitat preferences, 221
- mitochondrial DNA analysis, 220
- monitoring, research, and survey needs, 223–24
- pitfall trapping, 221
- population decline, 222
- as Priority 2 Species of Special Concern, 218
- radio tracking of, 221
- reproductive activity, 220
- risk factors, 218(t)
- road mortality, 221
- sexual maturity, 220
- status of, 218, 223
- surface activity, 221
- survival rates, 221
- taxonomy of, 220
- thermoregulation, 221
- threats faced by, 222–23
- trends in abundance, 222
- Phrynosoma mcallii*, 225–33, 225(f)
- breeding activity and egg laying, 227
- breeding season, 227
- clutch size, 227
- daily activity patterns, 227
- diet of, 228, 231
- distribution of, 226 (m), 229
- habitat loss and fragmentation, 230
- habitat requirements, 228–29
- hibernation, 227
- identification of, 225–27
- life history of, 227–28
- management of, 231–32
- mark-recapture studies of, 230, 232
- monitoring, research, and survey needs, 232–33
- mortality, causes of, 230
- phylogeographic study of, 227
- populations of, 229
- predators of, 228
- as Priority 2 Species of Special Concern, 225, 231
- radiotelemetry studies of, 227–28
- renewable energy projects, impact of, 231
- risk factors, 225(l)
- scat counts, 232
- sexual maturity, 228
- shade-seeking behavior, 227
- status of, 225, 231
- surface activity of, 227
- survivorship of, 228
- taxonomy of, 227
- threats faced by, 230–31
- trends in abundance, 229–30
- Phyllodactylus nocticolus*, 248
- Phylogeography, 45, 166, 194, 241, 309
- pitfall trapping, 165, 173, 199, 201, 221, 258, 265
- Pituophis catenifer pumilis*, 369
- Plestiodon skiltonianus interparietalis*, 369
- Plethodon asupak*, 370
- Plethodon elongatus*, 369, 372
- habitat, 372
- timber harvest, impact of, 372

- Plethodon stormi*, 370
 Prairie Creek Redwoods State Park, 170
Pseudacris regilla, 75
- radiotelemetry, 67, 87, 92, 105, 228, 259, 262, 264, 277, 278, 288
- Rana aurora*, 77–83, 77(f)
 artificial habitats, 80
 breeding activity and egg-laying, 79, 81
 breeding season, 79
 in coastal streams, 80
 distribution of, 78(m), 80
 embryonic development, 79
 habitat requirements for, 79–80
 identification of, 77–79
 impact of
 agricultural and residential development, 81
 climate changes, 82
 forest conversion, 81
 habitat loss and degradation, 81, 83
 marijuana cultivation, 81, 83
 pesticides and herbicides, 81
 sea-level-rise, 83
 life history of, 79
 management of, 83
 mesocosm experiment, 82
 metamorphosis of, 79
 monitoring, research, and survey needs, 83
 population decline, 80–81
 predators of, 81, 83
 as Priority 2 Species of Special Concern, 77, 82
 radio tracking of, 80, 81
 recolonization of, 81
 risk factors, 77(t)
 saltwater tolerance, 83
 status of, 77, 82
 taxonomy of, 79
 terrestrial *versus* aquatic life stages, 82
 threats faced by, 81–82
 trends in abundance, 80–81
- Rana boylei*, 53, 84–92, 84(f)
 breeding activity and egg-laying, 86
 breeding and oviposition, 87
 cannibalistic behavior, 87
 distribution of, 85(m), 90
 extirpations, 89
 food resources, 87
 habitat requirements for, 87–88
 hydrological effects, 92
 identification of, 84–86
 larvae of, 87
 life history of, 86–87
 management of, 91–92
 metamorphosis of, 87
 monitoring, research, and survey needs, 92
 movements to or from breeding sites, 87
 onset and duration of breeding, 86
 pesticides and herbicides, impact of, 90
 population dynamics, 92
 as Priority 1 Species of Special Concern, 84
 radiotelemetry studies, 87, 92
 risk factors, 84(t)
 skeletochronology of, 87
 in snowmelt-fed streams, 91
 status of, 84, 91
 survivorship of, 92
 taxonomy of, 86
 threats faced by, 89–91
 trends in abundance, 89
- Rana cascadae*, 93–99, 93(f)
 breeding activity and egg-laying, 95
 diet of, 95
 distribution of, 94(m), 96
 embryo development, 95
 habitat loss and modification, 97
 habitat requirements, 95–96
 identification of, 93–95
 isolation-by-distance model of gene flow, 95
 life history of, 95
 management of, 98–99
 monitoring, research, and survey needs, 99
 mortality of, 97
 pesticide use, impact of, 97
 population dynamics of, 96, 98
 predators of, 96, 97
 as Priority 2 Species of Special Concern, 93
 recolonization of, 98
 risk factors, 93(t)
 status of, 93, 98
 taxonomy of, 95
 threats faced by, 96–98
 trends in abundance, 96
 UV-B radiation, impact of, 97, 98
- Rana draytonii*, 100–105, 100(f)
 breeding activity and egg-laying, 102
 breeding season, 103
 diet of, 102
 distribution of, 101(m), 103–4
 effect of introduced species on, 104
 elevational range of, 104
 foraging behavior, 102
 habitat requirements, 102–3
 impact of
 cattle-grazing, 105
 climate change, 104–5
 pesticide use, 105
 urbanization and agriculture, 104
 larvae of, 102
 life history of, 102
 management of, 105

- mitochondrial DNA variation, 102, 105
- monitoring, research, and survey needs, 105
- population trends of, 103–4
- prey types, 102
- as Priority 1 Species of Special Concern, 100, 105
- radiotelemetry studies, 105
- risk factors, 100(t)
- status of, 100, 105
- taxonomy of, 102
- threats faced by, 104
- trends in abundance, 104
- USFWS recovery plan for, 105
- Rana muscosa*, 370
- Rana pipiens*, 106–111, 106(f)
 - distribution of, 107(m), 108–9
 - habitat requirements, 108
 - identification of, 106–8
 - impact of
 - climate changes, 109
 - habitat modification, 109
 - livestock grazing, 109
 - pesticide use, 109–10
 - life history of, 108
 - management of, 110
 - mitochondrial DNA variation, 110
 - monitoring, research, and survey needs, 110–11
 - morphology and vocalizations, 108
 - mortality of, 110
 - as Priority 1 Species of Special Concern, 106
 - risk factors, 106(t)
 - status of, 106, 110
 - taxonomy of, 108
 - threats faced by, 109–10
 - trends in abundance, 109
- Rana pretiosa*, 112–18, 112(f)
 - breeding activity and egg laying, 114
 - distribution of, 113(m), 115–16
 - genetic differentiation and population structure, 117–18
 - habitat requirements, 114–15
 - hibernation, 114
 - identification of, 112–14, 117
 - life history of, 114
 - management of, 117
 - mitochondrial DNA variation, 114
 - monitoring, research, and survey needs, 117–18
 - prey of, 114
 - as Priority 1 Species of Special Concern, 112, 117
 - risk factors, 112(t)
 - status of, 112, 117
 - survivorship of, 114
 - taxonomy of, 114
 - threats faced by, 116–17
 - trends in abundance, 116
- Rana sierrae*, 370
- ranavirus infections, 45, 140–41
- Rana yavapaiensis*, 119–24, 119(f)
 - coloration of, 121
 - diseases, 122
 - distribution of, 120(m), 122
 - habitat requirements for, 121–22
 - hybridization of, 122–23
 - identification of, 119–21
 - larvae of, 123
 - life history of, 121
 - management of, 123
 - molecular markers, 123
 - molecular phylogenetic analyses, 123
 - monitoring, research, and survey needs, 123–24
 - natural vulnerability, 123
 - population decline, 122–23
 - causes of, 124
 - predators of, 122–23
 - as Priority 1 Species of Special Concern, 119
 - reproductive biology of, 121
 - risk factors, 119(t)
 - salt tolerance in, 122
 - status of, 119, 123
 - taxonomy of, 121
 - threats faced by, 122–23
 - trends in abundance, 122
- range maps for each species, development of, 17(m)
- red-bellied newt. See *Taricha rivularis*
- Red Creek, Tehama County, 88
- red diamond rattlesnake. See *Crotalus ruber*
- red-eared slider. See *Trachemys scripta elegans*
- regal ring-necked snake. See *Diadophis punctatus regalis*
- relictual slender salamander. See *Batrachoseps relictus*
- relocation programs, genetic consequences of, 42
- Rhyacotriton variegatus*, 166–73, 166(f)
 - breeding activities and egg laying, 168
 - breeding season, 168
 - clutch size, 168
 - coastal populations, 171
 - diet of, 169
 - distribution of, 167(m), 169–70
 - geographic ranges and genetic distinctiveness, 168
 - habitat loss and degradation, 171
 - habitat requirements for, 169
 - identification of, 166–68
 - impacts of
 - climate change, 170
 - herbicides and pesticides, 171
 - marijuana cultivation, 170–72
 - roadbuilding, 170
 - timber harvesting, 170
 - landscape genetic studies of, 173
 - larvae of, 168
 - life history of, 168–69
 - management of, 172
 - mitochondrial DNA analysis, 168

- monitoring, research, and survey needs, 172–73
- pitfall trapping, 173
- as Priority 1 Species of Special Concern, 166
- risk factors, 166(t)
- sexual maturity, 168
- status of, 166, 171–72
- in stream channel, 169
- taxonomy of, 168
- thermal tolerances of, 169
- threats faced by, 170–71
- trends in abundance, 170
- river management, 91–92
- road construction, impact of, 41–42, 55–56, 159–60, 164
- Roundup®, 110
- Sacramento Valley, 300
- Salinas River Valley, 86, 88
- Salvadora hexalepis virgulata*, 279–84, 279(f), 368
 - climate change, impact of, 283
 - diet of, 281
 - distinctiveness of, 281
 - distribution of, 280(m), 282
 - genetic analysis of, 281
 - habitat requirements, 282
 - identification of, 279
 - life history of, 281–82
 - management of, 283
 - monitoring, research, and survey needs, 283–84
 - movement ecology of, 281
 - population decline, 282
 - prey of, 281
 - as Priority 2 Species of Special Concern, 279, 283
 - risk factors, 279(t)
 - status of, 279, 283
 - taxonomy of, 281
 - threats faced by, 281–82
- San Bernardino Mountains, 183
- San Diego banded gecko. See *Coleonyx variegatus abbotii*
- San Diego Multiple Species Habitat Conservation Plan, 31
- sandstone night lizard. See *Xantusia gracilis*
- San Felipe Creek drainage, 121–23
- San Gabriel Mountains slender salamander. See *Batrachoseps gabrieli*
- San Gabriel River, 301
- San Joaquin coachwhip. See *Masticophis flagellum ruddocki*
- San Joaquin Valley, 273, 300
- San Luis Obispo County, 153
- San Sebastian Marsh, 122
- San Simeon State Park, 287–88
- Santa Catalina Mountains, 306
- Santa Cruz black salamander. See *Aneides flavipunctatus niger*
- Santa Cruz Island, 369
- Santa Cruz Island gopher snake. See *Pituophis catenifer pumilis*
- Santa Cruz Mountains, 144–45, 163–64
- Santa Lucia Mountains, 145, 164
- Santa Ynez Mountains, 184
- Saproglenia*, 82
- Scaphiopus couchii*, 125–29, 125(f)
 - breeding activity and egg laying, 127
 - breeding season, 127
 - development rate of, 127
 - distribution of, 126(m), 127–28
 - emergence behavior, 127
 - foraging behavior, 127
 - habitat requirements, 127
 - identification of, 125
 - life history of, 127
 - management of, 128–29
 - metamorphosis of, 127
 - monitoring, research, and survey needs, 129
 - movement ecology of, 129
 - natural breeding sites, 129
 - physiological limits of, 128, 129
 - population dynamics of, 128
 - as Priority 3 Species of Special Concern, 125
 - risk factors, 125(t)
 - status of, 125, 128
 - taxonomy of, 127
 - terrestrial habitat, 129
 - threats faced by, 128
 - trends in abundance, 128
- Scott Bar salamander. See *Plethodon asupak*
- septicemia, 73
- Sierra Nevada, 11, 253, 269
- Sierra night lizard. See *Xantusia vigilis sierrae*
- Siskiyou Mountains salamander. See *Plethodon stormi*
- Six Rivers National Forest, 168, 170
- Smith River, 168
- snake trapping, 259
- Sonora mud turtle. See *Kinosternon sonoriense*
- Sonoran Desert toad. See *Bufo alvarius*
- southern California mountain kingsnakes. See *Lampropeltis multifasciata*; *Lampropeltis zonata parvirubra*
- southern long-toed salamander. See *Ambystoma macrodactylum sigillatum*
- southern torrent salamander. See *Rhyacotriton variegatus*
- southern western pond turtles. See *Emys marmorata*
- Southwest Climate Science Center, 44
- Spea hammondi*, 130–35, 130(f)
 - breeding activity and egg laying, 132
 - breeding aggregations, 132
 - breeding season, 132
 - development of, 133
 - diet of, 133

- distribution of, 131(m), 133
- growth rate of, 132
- habitat requirements, 133
- identification of, 130
- intraspecific variation in, 132
- larvae of, 132–33
- life history of, 132–33
- management of, 134–35
- metamorphosis of, 133
- migration patterns and distances, 135
- mitochondrial DNA, 130
- mitochondrial introgression, 132
- monitoring, research, and survey needs, 135
- movement and surface activity, 133
- natural vernal pool phenology, 135
- predators of, 133
- as Priority 1 Species of Special Concern, 130
- risk factors, 130(t)
- sexual characters, 132
- status of, 130, 134
- survivorship of, 133
- taxonomy of, 130–32
- threats faced by, 134
- trends in abundance, 133–34
- Species of Special Concern
 - aquatic or terrestrial habitat of, 27(f)
 - committed to extinction, 45
 - comparison of, 37(t)–38(t)
 - conservation-related research, 44
 - criteria for listing, 5
 - definition of, 8, 9
 - distribution of, 34(m)
 - down-grade status, 36
 - ecological patterns in metric scores, 30
 - ecoregion of, 26(m)
 - endemic and near endemic species, 33(t)
 - geographic patterns in, 32–36
 - identification of, 4
 - list of amphibian and reptiles, 20(t)–21(t)
 - museum collections and other data sources, 16(t)
 - nominee list, development of, 8
 - number of taxa in each status category among, 22(f)
 - occurrence of, 36(m)
 - patterns in the metric scores for, 22–25
 - principal components analysis of, 22
 - public comment, 12–13, 25, 367
 - ranking and determination of, 13
 - ranking scores among, 29
 - reptile and amphibian taxa under, 30(f)
 - research, survey, and monitoring needs, 43–48
 - risk metrics of, 8–12
 - distribution trend, 9–10, 10(t)
 - ecological tolerance, 10–11, 11(t)
 - endemism, 10, 10(t)
 - population concentration and migration, 10, 10(t)
 - population trend, 11, 11(t)
 - projected impacts, 12, 12(t)
 - range size, 9, 9(t)
 - vulnerability to climate change, 11, 11(t)
 - scoring of nominee taxa, 12
 - Spearman's rank correlations, 23(t)
 - total score/total possible for taxa evaluated for, 21(f)
 - watch list of, 13, 368–70
- spiny softshell turtle. See *Apalone spinifer*
- spring snowmelt, 11, 56, 90, 171
- State Wildlife Action Plans (SWAP), 5
- Sternotherus odoratus*, 306
- Stevens Creek watershed, 177
- stinkpot turtle. See *Sternotherus odoratus*
- stream-breeding amphibians, 164, 178
- stream hydrology, 90, 178
- Susan River, 300
- Tantilla planiceps*, 372–73
 - ecological requirements, 373
 - habitat requirements, 372
 - natural history of, 372
 - population densities, 372
- Taricha rivularis*, 174–79, 174(f)
 - aquatic breeding habitats, 177
 - breeding activities and egg laying, 176
 - breeding season, 176
 - buffer protections, ecological effects of, 178
 - diet of, 177
 - distinctiveness of, 176
 - distribution of, 175(m), 177
 - effects of climate change on, 177
 - habitat loss and fragmentation, 178
 - habitat requirements for, 177
 - identification of, 174–76
 - life history of, 176–77
 - management of, 178
 - mitochondrial DNA analysis, 176
 - monitoring, research, and survey needs, 179
 - mortality and habitat degradation, 178
 - as Priority 2 Species of Special Concern, 174, 178
 - risk factors, 174(t)
 - sexual maturity, 176
 - status of, 174, 178
 - taxonomy of, 176
 - temperature tolerances of, 177
 - threats faced by, 177–78
 - trends in abundance, 177
- Taricha torosa*, 180–85, 180(f)
 - aquatic breeding habitats, 182
 - breeding activities and egg laying, 182
 - distribution of, 181(m), 183
 - habitat corridor, 185
 - habitat requirements for, 183
 - identification of, 180–82
 - larvae of, 182–83
 - life history of, 182–83

- management of, 185
- metapopulation dynamics, 185
- monitoring, research, and survey needs, 185
- predators of, 184
- as Priority 2 Species of Special Concern, 180, 185
- risk factors, 180(t)
- road mortality, 185
- status of, 180, 185
- stream- and pond-breeding sites, 185
- survivorship of, 184
- taxonomy of, 182
- threats faced by, 184–85
- trends in abundance, 183–84
- Thamnophis hammondi*, 285–90, 285(f)
 - climate change, impact of, 289
 - distinctiveness of, 287
 - distribution of, 286(m), 288–89
 - foraging and basking activities, 287
 - habitat and movement ecology, 288
 - habitat requirements, 288
 - identification of, 285–87
 - life history of, 287–88
 - management of, 290
 - mating activities, 287
 - monitoring, research, and survey needs, 290
 - prey of, 288
 - as Priority 2 Species of Special Concern, 285
 - radiotelemetry study of, 288
 - risk factors, 285(t)
 - status of, 285, 290
 - surface activity, 289
 - taxonomy of, 287
 - threats faced by, 289
 - trends in abundance, 289
- Thamnophis sirtalis*, 291–95, 291(f)
 - captive breeding, 295
 - diet of, 294
 - distribution of, 292(m), 294
 - extirpations and population declines, 294
 - habitat requirements, 294
 - identification of, 291–93
 - impact of
 - climate change, 295
 - floods, 294
 - nonnative species, 295
 - life history of, 293–94
 - management of, 295
 - monitoring, research, and survey needs, 295
 - as Priority 1 Species of Special Concern, 291
 - risk factors, 291(t)
 - sexual maturity of, 293
 - status of, 291, 295
 - taxonomy of, 293
 - threats faced by, 294–95
 - trends in abundance, 294
- Thousand Palms Oasis Preserve, 230
- timber harvest, impact of, 35, 41, 55–57, 82–83, 159–60, 164, 169–70, 172, 178, 372
- Tioga Pass meadow, 72, 73
- Torrey Pines State Reserve, Point Loma, 258
- Trachemys scripta elegans*, 298, 302, 307
- translocation of animals, 42–43
- Truckee River, 300
- Tule River drainage, 158
- turtle trapping surveys, 307
- two-striped garter snake. *See* *Thamnophis hammondi*
- Uma notata*, 234–38, 234(f)
 - climate change, impact of, 237
 - distinctiveness of, 236
 - distribution of, 235(m), 237
 - habitat loss and fragmentation, 237
 - habitat requirements for, 237
 - identification of, 234–36
 - life history of, 236–37
 - management of, 237–38
 - mitochondrial DNA analysis, 236, 238
 - monitoring, research, and survey needs, 238
 - mortality, causes of, 237
 - as Priority 2 Species of Special Concern, 234
 - recolonization of, 238
 - risk factors, 234(t)
 - status of, 234, 237
 - taxonomy of, 236
 - threats faced by, 237
 - trends in abundance, 237
- Uma scoparia*, 239–45, 239(f)
 - breeding activity and egg laying, 242
 - breeding season, 241
 - burrowing and sand-swimming behavior, 242
 - clutch size, 242
 - diet of, 242
 - distinctiveness of, 241
 - distribution of, 240(m), 242–43
 - effects of climate change on, 243–44
 - efficiency at capturing prey, 243
 - extirpations of, 243
 - habitat loss and fragmentation, 243
 - habitat patches, for long-term persistence, 242
 - habitat requirements, 242
 - human-assisted translocation, 244, 245
 - identification of, 239–41
 - life history of, 241–42
 - management of, 244
 - mitochondrial DNA analysis, 245
 - monitoring, research and survey needs, 244–45
 - noise pollution effects, 244
 - phylogenetic analyses of, 241
 - population size and density, 242, 244–45
 - as Priority 3 Species of Special Concern, 239, 244
 - push-up behavior, 241
 - quality of available food, 242

- risk factors, 239(t)
- status of, 239, 244
- taxonomy of, 241
- threats faced by, 243–44
- trends in abundance, 243
- United States Department of Agriculture (USDA)
 - ecoregion subsections used in developing range maps, 18(m)
- urban development, impact of, 302
- US Fish and Wildlife Service (USFWS), 45, 105
- US Forest Service, 5
 - Old-growth Wildlife Project, 169
- US Geological Survey (USGS), 258
- venomous reptiles
 - Bufo alvarius*, 59–64
 - Crotalus ruber*, 260–65
 - Heloderma suspectum*, 212–17
 - predator avoidance mechanism, 215
- Warner Mountains, 115–17
- watch list, of Species of Special Concern
 - Baja California rat snake, 368
 - California tiger salamander, 368
 - Coronado skink, 369
 - Del Norte salamander, 369–70
 - large-blotched ensatina, 369
 - mountain yellow-legged frogs, 370
 - Mount Lyell web-toed salamander, 369
 - orange-throated whiptail, 368
 - Owens Valley web-toed salamander, 369
 - Santa Cruz Island gopher snake, 369
 - southern California mountain kingsnakes, 369
 - yellow-blotched ensatina, 368–69
- western black-headed snake. See *Tantilla planiceps*
- western pond turtle. See *Emys marmorata*
- western spadefoot. See *Spea hammondi*
- wildfires, 164, 184, 195, 210, 223, 258, 283
- Wildlife Action Plan (California), 5
- wildlife corridors. See habitat corridors
- Xantusia gracilis*, 246–49, 246(f)
 - behavioral differences in captivity, 248
 - distinctiveness of, 248
 - distribution of, 247(m), 248
 - habitat requirements, 248
 - identification of, 246–48
 - life history of, 248
 - management of, 249
 - monitoring, research, and survey needs, 249
 - as Priority 3 Species of Special Concern, 246
 - risk factors, 246(t)
 - status of, 246, 249
 - taxonomy of, 248
 - threats faced by, 249
 - trends in abundance, 248–49
- Xantusia henshawi*, 199
- Xantusia vigilis sierrae*, 250–54, 250(f)
 - daily activity cycles, 252
 - diet of, 252
 - distribution of, 251(m), 253
 - elevational range, 253
 - habitat requirements for, 252–53
 - identification of, 250–52
 - life history of, 252
 - management of, 253
 - migration corridors, 254
 - mitochondrial and nuclear DNA sequence, 252
 - monitoring, research, and survey needs, 253–54
 - phylogeographic/species boundary, 253
 - as Priority 3 Species of Special Concern, 250
 - risk factors, 250(t)
 - sexual maturity of, 252
 - status of, 250, 253
 - taxonomy of, 252
 - threats faced by, 253
 - trends in abundance, 253
- Xantusia wigginsi*, 373
- yellow-blotched ensatina. See *Ensatina eschscholtzii croceater*
- yellow-legged frogs. See *Rana boylei*
- Yosemite National Park, 71, 74–76
- Yosemite toad. See *Bufo canorus*
- Yuma Desert, 232

U.S. Fish and Wildlife Service

California - Great Basin Region

Migratory Birds Program

Recommended Buffer Zones for Ground-based Human Activities around Nesting Sites of Golden Eagles in California and Nevada

May 2021

For most ground-based human activities, we recommend a one-mile no-disturbance buffer surrounding golden eagle nesting sites in California and Nevada; see table below for specifics on activity and buffer recommendations. Recommended buffers may increase or decrease, depending on specific site or activity circumstances. Buffers may be reduced in consultation with the U.S. Fish and Wildlife Service (Service) when the nest is not in use or activities are not in line-of-sight of the nest^a. In parts of California, eagles maintain year-round territories that may require additional protection. We recommend consultation with the Service for determining buffer zones for high intensity or long duration activities, unique circumstances, activities not listed in the table below, or when historic levels of human activity are a consideration.

Activity	Recommended No-Disturbance Buffer
Use of Motorized Vehicles off-road and on water: Including, but not limited to, passenger vehicles, all-terrain vehicles, dirt bikes, and snowmobiles. Any passenger vehicle driving on dirt or gravel roads that are not part of a routinely used transportation corridor. Also includes motorized boating activities.	1 mile
Pedestrian and Non-Motorized Activity^b: Including, but not limited to, walking, running, hiking, biking, camping, rock climbing, bird watching, fishing, hunting, horseback riding, canoeing, kayaking, and biological surveys.	1 mile
Developed Sites: Including, but not limited to, facilities, developed campground sites, and designated snowmobile and off-road vehicle courses.	1 mile
Industrial, Municipal, and Construction Activity: Including, but not limited to, urbanization; mining; oil and gas development; solar development; logging; power line construction; road construction & maintenance; facilities construction; and agricultural operations.	1 mile
Blasting and other loud non-regular noise: Including, but not limited to, detonation devices, fireworks classified by the Federal Department of Transportation as Class B explosives, recreational shooting, and outdoor concerts.	2 miles

^a An *in-use nest* is defined as a "golden eagle nest characterized by the presence of one or more eggs, dependent young, or adult eagles on the nest in the past 10 days during the breeding season" (50 CFR 22.3) and "(b)reeding begins... with the start of courtship..." (*Programmatic Environmental Impact Statement for the Eagle Rule Revision*, United States Department of the Interior, Fish and Wildlife Service, December 2016).

^b Many existing nest sites experience some level of intermittent and on-going low levels of disturbance from these types of human activities, and the resident pair of eagles may have acclimated to these existing levels of disturbance. However, increases in human activity may not be tolerated by nesting eagles.

^c Qualified biologists conducting ground-based eagle monitoring may follow distance recommendations in Pagel et al (2010).

Joshua tree (*Yucca brevifolia*) seeds are dispersed by seed-caching rodents¹

Stephen B. VANDER WALL², Department of Biology and the Program in Ecology, Evolution and Conservation Biology, University of Nevada, Reno, Nevada 89557, USA, e-mail: sv@unr.edu

Todd ESQUE, United States Geological Survey, Henderson, Nevada, USA.

Dustin HAINES, Department of Ecology, Evolution, and Behavior, University of Minnesota, Mankato, Minnesota 55455, USA.

Megan GARNETT, United States Geological Survey, Henderson, Nevada, USA.

Ben A. WAITMAN, Department of Biology and the Program in Ecology, Evolution and Conservation Biology, University of Nevada, Reno, Nevada 89557, USA, and United States Geological Survey, Henderson, Nevada, USA.

Abstract: Joshua tree (*Yucca brevifolia*) is a distinctive and charismatic plant of the Mojave Desert. Although floral biology and seed production of Joshua tree and other yuccas are well understood, the fate of Joshua tree seeds has never been studied. We tested the hypothesis that Joshua tree seeds are dispersed by seed-caching rodents. We radioactively labelled Joshua tree seeds and followed their fates at five source plants in Potosi Wash, Clark County, Nevada, USA. Rodents made a mean of 30.6 caches, usually within 30 m of the base of source plants. Caches contained a mean of 5.2 seeds buried 3–30 mm deep. A variety of rodent species appears to have prepared the caches. Three of the 836 Joshua tree seeds (0.4%) cached germinated the following spring. Seed germination using rodent enclosures was nearly 15%. More than 82% of seeds in open plots were removed by granivores, and neither microsite nor supplemental water significantly affected germination. Joshua tree produces seeds in indehiscent pods or capsules, which rodents dismantle to harvest seeds. Because there is no other known means of seed dispersal, it is possible that the Joshua tree-rodent seed dispersal interaction is an obligate mutualism for the plant.

Keywords: Joshua tree, scatter hoarding, seed caching, seed dispersal, *Yucca brevifolia*

Résumé : L'arbre de Josué (*Yucca brevifolia*) est une plante distinctive et charismatique du désert de Mojave. Quoique la biologie de la floraison et la production de graines de cet arbre et des autres yuccas sont bien comprises, le sort des graines de l'arbre de Josué n'a jamais été étudié. Nous avons testé l'hypothèse que les graines de cet arbre sont dispersées par des rongeurs cacheurs de graines. Nous avons marqué radioactivement des graines d'arbres de Josué et suivi leur sort à partir de cinq plants sources à Potosi Wash, Clark County, Nevada, É-U. Les rongeurs ont fait en moyenne 30,6 caches habituellement à 30 m de la base des plants sources. Les caches contenaient en moyenne 5,2 graines enterrées à des profondeurs de 3 à 30 mm. Une variété d'espèces de rongeurs semblaient avoir fait ces caches. Trois des 836 graines d'arbres de Josué cachées (0,4%) ont germé le printemps suivant. La germination de graines dans des enclos à rongeurs atteignait presque 15%. Plus de 82% des graines dans les sites ouverts ont été enlevées par des granivores et ni le microsite ni l'ajout d'eau n'ont affecté significativement la germination. L'arbre de Josué produit des graines dans des capsules indéhiscentes que les rongeurs brisent pour récolter les graines. Dans le cas de l'arbre de Josué, puisque aucun autre moyen de dispersion des graines n'est connu, il est possible que la dispersion par les rongeurs soit une interaction mutualiste obligée pour cette plante.

Mots-clés : arbre de Josué, cache de graines, dispersion de graines, éparpiller en divers caches, *Yucca brevifolia*

Nomenclature: Hickman, 1993; Peilmyr, 2003

Introduction

Yuccas (*Yucca* and *Hesperoyucca*) are prominent and charismatic plants of warm temperate and arid regions of the southern United States and Mexico. Yuccas have received a great deal of scientific attention, especially in the area of floral biology. The yucca moth–yucca flower pollination interaction, often cited in textbooks and popular works, is one of the most convincing examples of a coevolved, obligate mutualism (Peilmyr *et al.*, 1996). The

interaction, which may have arisen more than 40 million years ago (Peilmyr & Leebens-Mack, 1999), was first described by George Engelmann and Charles Riley in the 1870s and is still an object of intense scientific investigation (Peilmyr, 2003). Although much is known about their genesis, virtually nothing is known about the fate of yucca seeds (e.g., seed dispersal, post-dispersal seed predation, seedling establishment).

Some yuccas (e.g. Joshua tree, *Yucca brevifolia*) produce relatively large seeds. These large seeds are likely to be attractive food items for granivorous rodents, birds, and ants. Further, a number of seed-caching rodents live in the

¹Rec. 2006-04-18, acc. 2006-06-20.
Associate Editor: Johannes Kollmann
²Author for correspondence.

same habitat as these yuccas, and it is possible that a fraction of the yucca seeds are scatter hoarded by rodents and that rodents might serve as dispersal agents for these yuccas. Some alternative means of seed dispersal have been suggested. In 1893, William Trelease wrote of Joshua tree fruits that "their rounded form and very light specific gravity render them well developed 'tumble fruits' and point to their dissemination across the desert by the aid of strong winds that prevail there." A number of authors state that yuccas (i.e., *Y. elata*, *Y. filamentosa*, *Y. whipplei*, and *Y. glauca*) are wind/gravity dispersed (Powell & Mackie, 1966; Wallen & Ludwig, 1978; Aker, 1982; Dodd & Linhart, 1994; Massey & Hamrick, 1998; Huth & Pellmyr, 2000), but the effectiveness of this mechanism has never been assessed. Wallen and Ludwig (1978) suggested that *Y. baccata* is dispersed by herbivores, but did not present conclusive evidence. Yeaton *et al.* (1985) also noted that *Y. schottigera*, in addition to *Y. baccata*, is dispersed by vertebrate means, but neither produced evidence or specified which vertebrates. Janzen (1986) suggested that the seeds of *Y. decipiens* in the southern Chihuahuan Desert, Mexico, might have been dispersed by extinct megafauna.

A novel and important reproductive feature of some *Yucca* species is the production of indehiscent seed pods. This is a derived condition, nested within the Agavaceae, while all other members have dry, dehiscent capsules. There are two types of indehiscence: one found in the fleshy-fruited yuccas of the section *Sarcocarpa* and the other in the spongy-fruited *Yucca brevifolia*, the sole member of the section *Chilocarpa*. Developmentally, they differ in that the mesocarp layer thickens in *Sarcocarpa*, whereas the exocarp layer thickens in *Y. brevifolia*, creating a less dense fruit in the latter. Many yucca species produce dehiscent seed pods, and these may well be dispersed by wind or some other abiotic vector, but it is unclear how seeds from the species with indehiscent pods are dispersed. The yuccas with indehiscent pods produce relatively large seeds that would appear to be sought after by granivores, however. The conjunction of these two traits, indehiscent pods and large seeds, led us to hypothesize that these yuccas, which include Joshua tree, are dispersed by seed-caching rodents that remove the seeds directly from pods. The goal of this study was to assess whether Joshua tree, one of the most familiar and important yuccas in the southwestern United States, is dispersed by seed-caching rodents.

Methods

We studied the dispersal of Joshua tree seeds at Potosi Wash on the southwest slope of the Spring Mountains, about 50 km southwest of Las Vegas, Clark County, Nevada (35° 52' N, 115° 33' W, elev. 1660–1700 m). The site is Mojave Desert scrub dominated by blackbrush (*Coleogyne ramosissima*), Nevada ephedra (*Ephedra nevadensis*), desert almond (*Prunus fasciculata*), and creosote bush (*Larrea tridentata*), with moderately dense populations of Joshua tree and banana yucca (*Y. baccata*) just below pinon-juniper woodland. The site is federal land administered by the US Department of Agriculture, Forest Service and US Department of the Interior, Bureau of Land Management. We studied seedling recruitment of Joshua tree along Walking

Box Ranch Road off Nevada route 164 about 12 km west of Searchlight, Nevada.

On 15 August 2002, we placed 200 Joshua tree seeds around each of two source plants (≈20 seeds at each of 10 scrapes in the soil within 1 m of the tree trunk). On 26 September 2002, we placed three more batches of 200 seeds under three other source plants in a similar fashion. Seeds were numbered individually and labelled with radioactive scandium, a gamma-emitting radionuclide with a half-life of 83.8 d. Scandium is biologically inert (does not pass through the food chain), decays to non-toxic, non-radioactive titanium, and appears harmless to rodents and seeds. The way we presented seeds was unnatural, but it is not feasible to present radioactively labelled seeds within pods and, in this study, we were interested in the dispersal process, not how seeds might be harvested from pods. We returned the day after deploying seed to check seed fates. We collected unharvested seeds and then searched for missing seeds using Geiger counters, following the procedure of Vander Wall (2002). We excavated caches to determine number of seeds and burial depth. We recorded the numbers on the seeds and returned them to the cache sites. We recorded caching microhabitat (under shrub [> 10 cm from canopy edge], edge of shrub [< 10 cm of canopy edge], or space between shrubs [> 10 cm from canopy edge]) and substrate (mineral soil or plant litter) and measured the coordinates of cache sites using the source plant as the origin of a graph space and cardinal directions as axes. We returned to source plants later in the fall of 2002 and spring of 2003 and resurveyed the area to determine which caches were present and to search for new (secondary) caches. After the last survey, all remaining labelled seeds were gathered and disposed of.

During fall 2003, after production of relatively large seed crops, we monitored fruits of Joshua tree, banana yucca, and Mojave yucca at our Potosi Wash study site and at Wallace Canyon in the Spring Mountains of southern Nevada to determine how seeds might be removed. Like Joshua tree, banana and Mojave yuccas produce indehiscent seed pods with relatively large seeds.

To determine rodent species composition and relative abundances at the Potosi Wash site, we trapped rodents on the evenings of 30 April and 1 May 2003. We used 98 Sherman live traps in a 7 × 7 grid with 12 m spacing and 2 traps at each point. Two trapping grids were laid out, one located in burned shrublands and the other in unburned vegetation. Bait consisted of mixed bird seed. We baited and set traps just before dusk, and we checked and closed traps at dawn. Trapping sessions lasted two nights.

We conducted seed germination trials using Joshua tree seeds gathered from Carpenter Canyon (Spring Mountains, southern Nevada) in October 2003. We removed seeds from pods and placed them in water; seeds that floated were considered inviable and were discarded. We also excluded any seeds with visible damage from moth larvae. Viable seeds were air dried and stored in paper bags.

To prevent pilferage by rodents, we conducted seed germination experiments within protective enclosures constructed with 6 mm hardware cloth 50 × 50 × 20 cm high, buried 10 cm deep in the soil. In each cage, we planted seeds ≈3 cm deep in a 5 × 10 grid. Wire mesh lids were placed

over the cages. We repeated this procedure in 24 exclosures (1200 seeds), spacing them ≈ 30 m apart along a transect. We placed half (12) of the exclosures under shrubs and the other half in the open, alternating site conditions along the transect. Six exclosures in each microsite treatment were watered each week for five weeks with one litre of water, again alternating along the transect. The other six received no supplemental water. We monitored exclosures weekly for seedling emergence. We initiated this study in March of 2004 and gathered the final data on 29 January 2005.

Nearby, we established 12 unfenced plots to measure the rate of rodent seed pilferage. Fifty Joshua tree seeds were buried ≈ 3 cm deep in a 5×10 pattern within a 50×50 -cm plot. As with the exclosures, half of these plots were watered each week. We checked plots weekly for seed removal and seedling emergence; however, the number of seeds removed was difficult to determine without disturbing the plots. At the end of two months, the top 5 cm of soil in the plots was removed and sifted for remaining seeds.

We analyzed germination and seed removal data using a two-factor ANOVA for microsite and supplemental watering. Data were square root transformed to meet assumptions of equal variance where necessary (Box & Cox, 1964). Data are reported as mean \pm 1 SD unless otherwise noted.

Results

Rodents removed labelled seeds from four of the source plants within 1 d, and seeds at the fifth source plant were removed during the second day. Only 5 of 1000 seeds (0.5%) were not harvested by rodents. None of the seeds had been eaten under the source plants. When we searched for seeds with Geiger counters, we found $84.1 \pm 13.1\%$ of removed seeds (range 67.7 to 97.5%) in caches. We found a total of 153 primary caches (30.6 ± 7.4 caches-plant⁻¹) located 0.5–56.6 m away from the base of source plants (Table 1). Primary caches contained a mean of 5.2 ± 3.9 seeds (range = 1–16 seeds) and were widely scattered around source plants (Figure 1). Mean maximum dispersal distance was 30.0 ± 16.8 m (range = 12.5–56.6 m). Five banana yucca seeds were found among three of the caches, indicating that rodents were foraging for and caching indigenous yucca seeds. We also found one lizard containing 44 Joshua tree seeds in a dead, hollow, banana yucca stem.

We resurveyed the areas around source plants later in the fall and found that 70 caches were still intact (45.8%), four caches had most seeds removed but one or more seeds still present (2.6%), 79 caches were missing (51.6%), and

50 new caches had been made (Table 1). Rodents made these new (secondary) caches from seeds they had taken from the original or primary caches. Secondary caches were smaller, containing an average of 3.3 ± 2.5 seeds (1–14 seeds; Table 1), and usually were widely scattered (Figure 1). The transport distance between primary and secondary caches averaged 6–13 m (full range = 0.2–32.2 m) (Table II).

Combining both primary and secondary caches, rodents placed nearly equal numbers of caches under shrubs (32.0%), near the edge of shrubs (36.5%), and in the open (31.5%). Caches were usually close to or under black-brush, ephedra, lycium (*Lycium andersonii*), or creosote bush, which are the most abundant shrubs at the study site. Rodents made caches in sandy or gravelly soil (68.2%) or under thin deposits of plant litter (31.8%).

During the course of the fall and winter, rodents ate many of the seeds in primary and secondary caches. On 23 March 2003, we found only 3 seedlings emerging from caches (the total number was probably greater because most seed had been removed, and rodents may have cached many seeds elsewhere).

From 392 trap-nights, we caught only 21 rodents (13 individuals captured). Trapping yielded two longtail pocket mice (*Chaetodipus formosus*), one canyon mouse (*Peromyscus crinitus*), and 10 Merriam's kangaroo rats (*Dipodomys merriami*). All but one of the Merriam's kangaroo rats were found on the burned site, and all other rodents were caught on the unburned site. The diurnal whitetail antelope ground squirrels (*Ammospermophilus leucurus*) were seen on the site but not captured.

The fruits of Joshua tree, banana yucca, and Mojave yucca that we monitored in fall 2003 had been gnawed on by rodents, both still attached to the plants and on the ground nearby. Rodents had dragged some pods away from productive plants, opened the pods, and appeared to have removed seeds. Based on teeth marks on the pods and foot prints in the soil, this activity appeared to be caused by antelope ground squirrels and perhaps woodrats (*Neotoma* sp.), but other species of rodents (e.g., mice or kangaroo rats) may have removed seeds from the partially opened pods. Over a period of 2–3 months, rodents dismantled most of the pods, and most of the seeds in these pods disappeared. We did not observe any other means of seed removal.

A total of 178 Joshua tree seedlings (14.8%) emerged in the germination exclosure study. More seedlings emerged under shrubs ($22.8 \pm 19.4\%$) than in the open ($8.8 \pm 8.6\%$), but the difference is only marginally significant ($F_{1,20} = 3.658$, $P < 0.070$). More seedlings emerged in

TABLE 1. Results of radioactive seed studies around five source plants. Data are means \pm 1 SD. Primary caches are from seeds taken from beneath source plants; secondary caches are from seeds taken from primary caches. Distance is the shortest distance between source plant and cache. Cache depth was measured to the top of seeds. Between 67.7 and 97.5% of the 200 seeds placed at each source plant were found.

Tree	Primary caches				Secondary caches			
	n	Seeds-cache ⁻¹	Distance (m)	Depth (mm)	n	Seeds-cache ⁻¹	Distance (m)	Depth (mm)
1	26	6.4 \pm 3.7	12.7 \pm 7.1	17.0 \pm 11.5	5	2.6 \pm 1.5	15.9 \pm 2.5	5.0 \pm 0.7
2	18	3.6 \pm 3.3	18.1 \pm 13.7	19.5 \pm 21.6	6	2.2 \pm 0.8	11.1 \pm 8.0	3.7 \pm 1.6
3	27	7.2 \pm 4.4	10.6 \pm 6.6	31.8 \pm 9.3	16	2.9 \pm 1.7	9.0 \pm 4.4	7.1 \pm 7.9
4	39	2.3 \pm 0.9	15.3 \pm 9.5	3.1 \pm 5.0	0	-	-	-
5	23	8.3 \pm 3.8	8.6 \pm 2.8	7.0 \pm 7.5	23	3.9 \pm 3.3	13.7 \pm 7.5	8.5 \pm 13.9




FIGURE 1. The dispersion of caches around Joshua tree 5 (large circle). Closed circles represent primary caches; open circles represent empty primary caches; triangles represent secondary caches made from the emptied primary caches. Units are in metres.

TABLE II. The distance that rodents moved seeds between primary and secondary caches at four source plants. There are more 1°-to-2° cache distances than secondary caches (Table I) because rodents often buried seeds from two primary caches in the same secondary cache.

Source plant	n	Mean \pm SD	Range
1	7	13.2 \pm 8.4	0.9–33.0
2	7	11.0 \pm 12.1	0.2–32.2
3	16	6.0 \pm 2.8	1.6–12.3
4	0		
5	27	12.0 \pm 6.9	3.6–24.9

unwatered enclosures (16.6 \pm 18.2%) than in watered enclosures (13.2 \pm 13.8%), but the difference was not significant ($F_{1,20} = 0.303$, $P = 0.585$). The shrub-water interaction also was not significant ($F_{1,20} = 0.017$, $P = 0.897$). Over a period of two months, 82% of the 600 seeds in unfenced plots were removed by animals. There was no significant effect of watering ($F_{1,6} = 0.17$, $P = 0.6888$) on the number of seeds removed, but animals took significantly more seeds under shrub canopies (16.2 \pm 15.9) than in the open (1.5 \pm 1.8) ($F_{1,6} = 5.49$, $P = 0.0472$).

Discussion

Our source plant studies demonstrated that rodents will gather and cache Joshua tree seeds. Since we saw essentially the same pattern at all five source plants, scatter hoarding of these yucca seeds appears to be a consistent behavioural response. However, we are not certain which rodent species were responsible for the caches. At source plant 2, several primary caches were pilfered during the day while we were excavating and mapping primary caches, suggesting that white-tailed antelope ground squirrels, the only diurnal rodent at the site, were probably involved. At source

plant 4, caches were uniformly small (1–4 seeds) and shallow (≈ 3 mm deep), suggesting that a small rodent such as a *Peromyscus* might have cached the seeds. Deer mice and piñon mice make caches similar to the caches at source plant 4 (Vander Wall *et al.*, 2001; Hollander & Vander Wall, 2004). Caches made at source plants 1, 3, and 5 were larger and deeper and might have been made by a larger rodent such as *Dipodomys merriami*.

Only three of the Joshua tree seeds remained in caches until they germinated in the spring and established seedlings. This is only a tiny percentage (0.36%) of the 836 seeds found in primary caches and 0.66% of the 456 seeds still in place at our last autumn survey (mid October to mid November). However, this study took place in the middle of an extensive drought period in which conditions for germination were far from optimal. Labeled seeds were removed after the last cache survey, and thus it was not possible to quantify germination in subsequent years, which may have been closer to optimal conditions.

The germination study inside enclosures, which prevented pilferage by rodents, indicates that seeds in shallow surface caches similar to those made by rodents can germinate and establish seedlings. In the laboratory, germination rates of moistened Joshua tree seeds are typically 90–95% over ≈ 5 d (Garrett & Esque, unpubl. data). The much lower percentage in the field (14.8%) indicates that many of the viable seeds did not germinate successfully during trials. More seeds successfully germinated and produced seedlings under shrubs, but the variation was so great that this trend was not statistically significant. Because we controlled for the loss of seeds by rodents, we assume that abiotic factors, soil invertebrates, or microbes reduced germination.

The pilfering of seeds from caches and the moving of pilfered seeds to secondary caches suggest that the dynamics of scatter-hoarded seeds is similar to that observed with some pine seeds (Vander Wall & Joyner, 1998; Vander Wall, 2002). Most yucca seeds reside at several different sites before finally being consumed or germinating, and removal of seeds from cache sites is not an accurate measure of seed predation. The re-caching of seeds has important consequences for seed dispersal, as seeds are dispersed from original caches they may be moved further from the original source plant, split into additional caches, and moved to a greater variety of habitats (Hoshizaki, Suzuki & Sasaki, 1997; Bohning-Gaese, Gaese & Rabemanantsoa, 1999; Roth & Vander Wall, 2005). Because many seeds were moved from their original cache site and probably re-cached, our estimate of establishment from natural caches is likely to be an underestimate of true seedling establishment. We were unable to conduct extensive, repeated surveys of the areas around source plants, so many secondary (and tertiary) caches were probably missed, and the fate of these seeds is unknown. Also, some of the re-cached seeds were probably carried farther from the source plant, suggesting that maximum seed dispersal distances were probably greater than the mean maximum distance of 30 m recorded for primary caches.

Rodents dismantled mature pods of Joshua tree and other yucca species and removed seeds. Most of the seeds appeared to have been carried away intact, because there was little evidence that rodents were eating seeds near the



**State Policy for Water Quality Control: State Wetland
Definition and Procedures for Discharges of
Dredged or Fill Material to Waters of the State**

**[Also for Inclusion in the Water Quality Control Plan for
Ocean Waters of California, and the Water Quality Control
Plan for Inland Surface Waters and Enclosed Bays and
Estuaries for Waters of the United States]**

STATE WATER RESOURCES CONTROL BOARD

**Adopted April 2, 2019
and Revised April 6, 2021**

I.	Introduction	1
II.	Wetland Definition	1
III.	Wetland Delineation	3
IV.	Procedures for Regulation of Discharges of Dredged or Fill Material to Waters of the State....	4
A.	Project Application Submittal for Individual Orders	5
B.	Permitting Authority Review and Approval of Applications for Individual Orders	9
C.	General Orders.....	12
D.	Activities and Areas Excluded from the Application Procedures for Regulation of Discharges of Dredged or Fill Material to Waters of the State	13
V.	Definitions	15
	Appendix A: State Supplemental Dredge or Fill Guidelines.....	19
	Subpart A – General.....	19
	Subpart B – Compliance with Guidelines.....	20
	Subpart E –Special Aquatic Sites	21
	Subpart H – Actions to Minimize Adverse Effects.....	22
	Subpart J – Compensatory Mitigation for Losses of Aquatic Resources	25

Procedures for Discharges of Dredged or Fill Material to Waters of the State

I. Introduction¹

The mission of the State Water Resources Control Board and the Regional Water Quality Control Boards (Water Boards) includes the preservation, enhancement, and restoration of the quality of California's water resources for the protection of the environment and all beneficial uses for the benefit of present and future generations. In accordance with the Porter-Cologne Water Quality Control Act (Water Code, § 13000 et seq.), the Water Boards are authorized to regulate discharges of waste, which includes discharges of dredged or fill material, that may affect the quality of waters of the state. As described below, waters of the state include some, but not all, features that are defined as wetlands, as well as other features, including the ocean, lakes, and rivers. These wetlands provide environmental and economic benefits to the people of this state, including flood and stormwater control, surface and ground water supply, fish and wildlife habitat, erosion control, pollution treatment, nutrient cycling, and public enjoyment. Wetlands ameliorate the effects of global climate change by providing floodwater storage, sequestering carbon, and maintaining vulnerable plant and animal communities. Many of these invaluable areas statewide have been lost to fill and development. Presently, wetlands are threatened by impacts from increasing population growth, land development, sea level rise, and climate change. These Procedures for the Discharges of Dredged or Fill Material to Waters of the State (Procedures) conform to Executive Order W-59-93, commonly referred to as California's "no net loss" policy for wetlands. In accordance with Executive Order W-59-93, the Procedures ensure that the Water Boards' regulation of dredge or fill activities will be conducted in a manner "to ensure no overall net loss and long-term net gain in the quantity, quality, and permanence of wetlands acreage and values..." The Water Boards are committed to increasing the quantity, quality, and diversity of wetlands that qualify as waters of the state.

These Procedures contain a wetland definition in section II and wetland delineation procedures in section III, both of which apply to all Water Board programs. The wetland definition encompasses the full range of wetland types commonly recognized in California, including some features not protected under federal law, and reflects current scientific understanding of the formation and functioning of wetlands. These Procedures also include procedures for the submission, review and approval of applications for activities that could result in the discharge of dredged or fill material to any waters of the state in section IV. The Procedures include elements of the Clean Water Act Section 404(b)(1) Guidelines, thereby bringing uniformity to Water Boards' regulation of discharges of dredged or fill material to all waters of the state. The effective date of these Procedures shall be May 28, 2020.

II. Wetland Definition

The Water Boards define an area as wetland as follows:

An area is wetland if, under normal circumstances, (1) the area has continuous or recurrent saturation of the upper substrate caused by groundwater, or shallow surface water, or both; (2) the duration of such saturation is sufficient to cause anaerobic conditions in the upper substrate; and (3) the area's vegetation is dominated by hydrophytes or the area lacks vegetation.

¹ [NOTE: These Procedures will be incorporated into the Water Quality Control Plans for (1) Inland Surface Waters Enclosed Bays and Estuaries for waters for which water quality standards are required by the Federal Water Pollution Control Act and acts amendatory thereof or supplementary thereto and (2) Ocean Waters of California. Future incorporation of the Procedures, as adopted, into the water quality control plans will be considered non-substantive amendments. At that time, formatting and other organizational edits necessary for incorporation into the water quality control plans will be addressed.]

Procedures for Discharges of Dredged or Fill Material to Waters of the State

39 The Water Code defines "waters of the state" broadly to include "any surface water or groundwater,
40 including saline waters, within the boundaries of the state." "Waters of the state" includes all "waters of
41 the U.S."² The following wetlands are waters of the state:

- 42 1. Natural wetlands,
- 43 2. Wetlands created by modification of a surface water of the state,³ and
- 44 3. Artificial wetlands⁴ that meet any of the following criteria:
 - 45 a. Approved by an agency as compensatory mitigation for impacts to other waters of the state,
46 except where the approving agency explicitly identifies the mitigation as being of limited
47 duration;
 - 48 b. Specifically identified in a water quality control plan as a wetland or other water of the state;
 - 49 c. Resulted from historic human activity, is not subject to ongoing operation and maintenance,
50 and has become a relatively permanent part of the natural landscape; or
 - 51 d. Greater than or equal to one acre in size, unless the artificial wetland was constructed, and
52 is currently used and maintained, primarily for one or more of the following purposes (i.e.,
53 the following artificial wetlands are not waters of the state unless they also satisfy the criteria
54 set forth in 2, 3a, or 3b):
 - 55 i. Industrial or municipal wastewater treatment or disposal,
 - 56 ii. Settling of sediment,
 - 57 iii. Detention, retention, infiltration, or treatment of stormwater runoff and other
58 pollutants or runoff subject to regulation under a municipal, construction, or industrial
59 stormwater permitting program,
 - 60 iv. Treatment of surface waters,
 - 61 v. Agricultural crop irrigation or stock watering,
 - 62 vi. Fire suppression,
 - 63 vii. Industrial processing or cooling,
 - 64 viii. Active surface mining – even if the site is managed for interim wetlands functions
65 and values,

² Therefore, wetlands that meet the current definition, or any historic definition, of waters of the U.S. are waters of the state. In 2000, the State Water Resources Control Board determined that all waters of the U.S. are also waters of the state by regulation, prior to any regulatory or judicial limitations on the federal definition of waters of the U.S. (California Code of Regulations title 23, section 3831(w).) This regulation has remained in effect despite subsequent changes to the federal definition. Therefore, waters of the state includes features that have been determined by the U.S. Environmental Protection Agency (U.S. EPA) or the U.S. Army Corps of Engineers (Corps) to be "waters of the U.S." In an approved jurisdictional determination, "waters of the U.S." identified in an aquatic resource report verified by the Corps upon which a permitting decision was based; and features that are consistent with any current or historic final judicial interpretation of "waters of the U.S." or any current or historic federal regulation defining "waters of the U.S." under the federal Clean Water Act.

³ "Created by modification of a surface water of the state" means that the wetland that is being evaluated was created by modifying an area that was a surface water of the state at the time of such modification. It does not include a wetland that is created in a location where a water of the state had existed historically, but had already been completely eliminated at some time prior to the creation of the wetland. The wetland being evaluated does not become a water of the state due solely to a diversion of water from a different water of the state.

⁴ Artificial wetlands are wetlands that result from human activity.

Procedures for Discharges of Dredged or Fill Material to Waters of the State

- 66 ix. Log storage,
67 x. Treatment, storage, or distribution of recycled water, or
68 xi. Maximizing groundwater recharge (this does not include wetlands that have
69 incidental groundwater recharge benefits); or
70 xii. Fields flooded for rice growing.⁵

71 All artificial wetlands that are less than an acre in size and do not satisfy the criteria set forth in 2, 3.a,
72 3.b, or 3.c are not waters of the state. If an aquatic feature meets the wetland definition, the burden is
73 on the applicant to demonstrate that the wetland is not a water of the state.

74 III. Wetland Delineation

75 The permitting authority shall rely on any wetland area delineation from a final aquatic resource report
76 verified by the U.S. Army Corps of Engineers (Corps) for the purposes of determining the extent of
77 wetland waters of the U.S. A delineation of any wetland areas potentially impacted by the project that
78 are not delineated in a final aquatic resource report verified by the Corps shall be performed using the
79 methods described in the three federal documents listed below (collectively referred to as "1987 Manual
80 and Supplements") to determine whether the area meets the state definition of a wetland as defined
81 above. As described in the 1987 Manual and Supplements, an area "lacks vegetation" if it has less
82 than 5 percent areal coverage of plants at the peak of the growing season. The methods shall be
83 modified only to allow for the fact that the lack of vegetation does not preclude the determination of
84 such an area that meets the definition of wetland. Terms as defined in these Procedures shall be used
85 if there is conflict with terms in the 1987 Manual and Supplements.

- 86 • Environmental Laboratory. 1987. U.S. Army Corps of Engineers Wetlands Delineation
87 Manual. Technical Report Y-87-1. U.S. Army Engineer Waterways Experiment Station,
88 Vicksburg, MS.
- 89 • U.S. Army Corps of Engineers. 2008. Regional Supplement to the Corps of Engineers
90 Wetland Delineation Manual: Arid West Region (Version 2.0). ed. J. S. Wakeley, R. W.
91 Lichvar, and C. V. Noble. ERDC/EL TR-08-28. Vicksburg, MS: U.S. Army Engineer Research
92 and Development Center.
- 93 • U.S. Army Corps of Engineers. 2010. Regional Supplement to the Corps of Engineers
94 Wetland Delineation Manual: Western Mountains, Valleys, and Coast Region (Version 2.0).
95 ed. J. S. Wakeley, R. W. Lichvar, and C. V. Noble. ERDC/EL TR-10-3. Vicksburg, MS: U.S.
96 Army Engineer Research and Development Center.

⁵ Fields used for the cultivation of rice (including wild rice) that have not been abandoned due to five consecutive years of non-use for the cultivation of rice (including wild rice) that are determined to be a water of the state in accordance with these Procedures shall not have beneficial use designations applied to them through the Water Quality Control Plan for the Sacramento and San Joaquin River Basins, except as otherwise required by federal law for fields that are considered to be waters of the United States. Further, agricultural inputs legally applied to fields used for the cultivation of rice (including wild rice) shall not constitute a discharge of waste to a water of the state. Agricultural inputs that migrate to a surface water or groundwater may be considered a discharge of waste and are subject to waste discharge requirements or waivers of such requirements pursuant to the Water Board's authority to issue or waive waste discharge requirements or take other actions as applicable.

Procedures for Discharges of Dredged or Fill Material to Waters of the State

97 **IV. Procedures for Regulation of Discharges of Dredged or Fill Material to** 98 **Waters of the State**

99 The purpose of this section is to establish application procedures for discharges of dredged or fill
100 material to waters of the state, which includes both waters of the U.S. and non-federal waters of the
101 state. This section supplements existing state requirements for discharges of dredged or fill material to
102 waters of the U.S.⁶ These Procedures include Appendix A, which contains relevant portions of the
103 U.S. EPA's Section 404(b)(1) "Guidelines for Specification of Disposal Sites for Dredge or Fill
104 Material"⁷ (Guidelines), with minor modifications to make them applicable to the state dredged or fill
105 program (hereafter State Supplemental Dredge or Fill Guidelines).⁸ This section applies to all
106 applications for discharges of dredged or fill material to waters of the state submitted after [insert date
107 that is nine months after approval by the Office of Administrative Law].⁹ The Procedures do not apply
108 to applications that are submitted prior to [insert date that is nine months after approval by the Office of
109 Administrative Law].

110 Unless excluded by section IV.D, applicants must file an application with the Water Boards for any
111 activity that could result in the discharge of dredged or fill material to waters of the state in accordance
112 with California Code of Regulations, title 23, section 3855.¹⁰ This application requirement applies to
113 new discharges, proposed material changes in the character, location, or volume of existing
114 discharges, and upon renewal of existing Orders for existing discharges. The permitting authority may
115 amend an existing Order solely for the purpose of extending the expiration date without requiring a new
116 application.

117 The applicant may consult with the Water Boards to determine whether a project could result in a
118 discharge of dredged or fill material to waters of the state and/or discuss submittals that would meet the
119 application requirements listed below. Discharges of dredged or fill material or other waste material to

⁶ California Code of Regulations, title 23, sections 3630-3669 (state's Clean Water Act (CWA) section 401 (33 USC § 1341) water quality certification program)

⁷ 40 C.F.R. § 230.

⁸ The State Supplemental Dredge or Fill Guidelines are included as Appendix A. Because the State Supplemental Dredge or Fill Guidelines are derived directly from the U.S. EPA's 404(b)(1) Guidelines, it uses slightly different terms than terms used in sections I through V of these Procedures. The State Supplemental Dredge or Fill Guidelines will be applied in a manner consistent with sections I through V of these Procedures.

⁹ In cases where the applicant is a state agency and is acting as the CEQA lead agency for one or more projects otherwise subject to this section, and that state agency is a party to an existing written agreement (e.g., memorandum of understanding) with the State Water Board that sets out alternative procedures and requirements regarding the submission, review or approval of project applications, the permitting authority shall apply the terms and conditions of the agreement in lieu of the terms and conditions of this section. After adoption of these Procedures, the State Water Board may also enter into such written agreements after consideration at a public meeting; such an agreement may include, for example, early consultation regarding potential project applications, early identification and analysis of project alternatives and mitigation measures, and dispute resolution. Any written agreements, whether existing or entered into after the adoption of these Procedures, may be amended in writing at any time by joint agreement of the parties, and such amended agreements shall govern in lieu of the terms and conditions of this section. All other applicable laws, including requirements for public notice and comment, apply to the permitting authorities' approval of projects under such an agreement.

¹⁰ Note that California Code of Regulations, title 23, section 3855 applies only to individual water quality certifications, but these Procedures extend the application of section 3855 to individual waste discharge requirements for discharges of dredged or fill material to waters of the state and waivers thereof.

Procedures for Discharges of Dredged or Fill Material to Waters of the State

120 areas that are not waters of the state, but that could affect the quality of waters of the state, may be
121 addressed under other Water Board regulatory programs.

122 **A. Project Application Submittal for Individual Orders**

123 The requirements set forth in sections IV.A and IV.B apply only to individual orders. Applicants must
124 submit the items listed in subsection 1 to the permitting authority. In addition, applicants shall consult
125 with the permitting authority about the items listed in subsection 2. Within 30 days of receiving the
126 items listed in subsection 1, the permitting authority may require the applicant to submit one or more of
127 the items in subsection 2 for a complete application. Applicants are encouraged to consult with the
128 permitting authority to determine the appropriate level of detail for the items in subsections 1 (and 2, if
129 applicable). Within 30 days of receiving all of the required items, the permitting authority shall determine
130 whether the application is complete and notify the applicant accordingly. If the applicant's federal
131 license or permit application includes any of the information required in subsections 1 or 2 below, the
132 applicant may submit the federal application materials to satisfy the corresponding state application
133 information. If federal application materials are submitted as part of the state application, the applicant
134 shall indicate where the corresponding state application information can be found in the federal
135 application materials.

136 **1. Items Required for a Complete Application**

- 137 a. All items listed in California Code of Regulations, title 23, section 3856 "Contents of a Complete
138 Application."¹¹
- 139 b. If the Corps requires an aquatic resource delineation report, a copy of the report verified by the
140 Corps.
- 141 c. A delineation of any waters that are not delineated in an aquatic resource delineation report
142 verified by the Corps. If such waters include wetlands, the wetlands must be delineated as
143 described in section III.
- 144 d. The dates upon which the overall project activity will begin and end, and, if known, the date(s)
145 upon which the discharge(s) will take place.
- 146 e. Map(s) with a scale of at least 1:24000 (1" = 2000') and of sufficient detail to accurately show
147 (1) the boundaries of the lands owned or to be utilized by the applicant in carrying out the
148 proposed activity, including the grading limits, proposed land uses, and the location, dimensions
149 and type of any structures erected (if known) or to be erected and (2) all aquatic resources that
150 may qualify as waters of the state, within the boundaries of the project, and all aquatic
151 resources that may qualify as waters of the state outside of the boundary of the project that
152 could be impacted by the project. A map verified by the Corps may satisfy this requirement if it
153 includes all potential waters of the state. The permitting authority may require that the map(s)
154 be submitted in electronic format (e.g., GIS shapefiles).
- 155 f. A description of the waters proposed to be impacted by the dredge or fill activity. The
156 description should include the beneficial uses as listed in the applicable water quality control
157 plan; a description of the activity at each individual discharge or dredge location; quantity of
158 impacts to waters proposed to receive a discharge of dredged or fill material at each location
159 rounded to at least the nearest one-hundredth (0.01) of an acre, nearest linear foot, and quantity

¹¹ Note that California Code of Regulations, title 23, section 3856 applies only to individual water quality certifications, but these Procedures extend the application of section 3856 to individual waste discharge requirements for discharges of dredged or fill material to waters of the state and waivers thereof.

Procedures for Discharges of Dredged or Fill Material to Waters of the State

- 160 of impacts to waters proposed to be dredged at each dredging location to the nearest cubic yard
161 (as applicable); assessment of potential direct and indirect impacts resulting from the discharge
162 or dredging activity and potential mitigation measures for those potential impacts, identification
163 of existing water quality impairment(s); the source of water quality impairment(s), if known; and
164 the presence of rare, threatened or endangered species¹² habitat.
- 165 g. An alternatives analysis,¹³ unless any of the following exemptions apply. The exemption from
166 the alternatives analysis requirement does not preclude a permitting authority from requiring the
167 applicant to demonstrate in its application that the project complies with section IV.B.1.a.
- 168 i. The project includes discharges to waters of the state outside of federal jurisdiction, but the
169 entire project would meet the terms and conditions of one or more Water Board-certified
170 Corps' General Permits, including any Corps District's regional terms and conditions, if all
171 discharges were to waters of the U.S. The permitting authority will verify that the entire
172 project would meet the terms and conditions of the Corps' General Permit(s) if all
173 discharges, including discharges to waters of the state outside of federal jurisdiction, were
174 to waters of the U.S. based on information supplied by the applicant.
- 175 ii. The project includes only discharges to waters of the U.S. and meets the terms and
176 conditions for coverage under an uncertified Corps' General Permit, including any Corps
177 District's regional terms and conditions. This exemption does not apply if the discharge of
178 dredged or fill material will directly impact:
- 179 a) more than two-tenths (0.2) of an acre or 300 linear feet of waters of the state;
180 b) rare, threatened, or endangered species habitat in waters of the state;
181 c) wetlands or eel grass beds; or
182 d) Outstanding National Resource Waters or Areas of Special Biological
183 Significance.
- 184 iii. The project would be conducted in accordance with a watershed plan that has been
185 approved for use by the permitting authority and analyzed in an environmental document
186 that includes a sufficient alternatives analysis, monitoring provisions, and guidance on
187 compensatory mitigation opportunities.
- 188 iv. The project is an Ecological Restoration and Enhancement Project.
- 189 v. The project has no permanent impacts to aquatic resources and no impacts to rare,
190 threatened or endangered species habitat in waters of the state, wetlands or eel grass
191 beds, Outstanding National Resource Waters or Areas of Special Biological Significance,

¹² "Rare, threatened, or endangered species" as used in the Procedures refers to plant and animal species listed as rare, threatened, or endangered pursuant to the California Endangered Species Act of 1964 (Fish & Game Code, § 2050 et seq.), the Native Plant Protection Act of 1977 (Fish & Game Code, § 1900 et seq.), or the Federal Endangered Species Act of 1973 (16 U.S.C. § 1531 et seq.).

¹³ "Alternatives analysis" as used in these Procedures refer to the analysis required by section IV.A.1.h and is a means to comply with the State Supplemental Dredge or Fill Guidelines, section 230.10(a). An alternatives analysis also may be required in order to comply with other statutory or regulatory requirements, such as CEQA or a Regional Board water quality control plan discharge prohibition. The exemptions and the tiers set forth below do not affect any alternatives analysis conducted pursuant to another statutory or regulatory requirement. To the extent that the permitting authority is acting as the lead agency under CEQA, it may be necessary for the permitting authority to conduct further analysis to comply with CEQA.

Procedures for Discharges of Dredged or Fill Material to Waters of the State

- 192 and all implementation actions in the restoration plan can reasonably be concluded within
193 one year.
- 194 h. If none of the above exemptions apply, the applicant must submit an alternatives analysis
195 consistent with the requirements of section 230.10 of the State Supplemental Dredge or Fill
196 Guidelines that allows the permitting authority to determine whether the proposed project is the
197 Least Environmentally Damaging Practicable Alternative (LEDPA). If the applicant submitted
198 information to the Corps to support an alternatives analysis, the applicant shall provide that
199 information to the permitting authority. Such information may satisfy some or all of the following
200 requirements in accordance with section IV.B.3. Alternatives analyses shall be completed in
201 accordance with the following tiers. The level of effort required for an alternatives analysis
202 within each of the three tiers shall be commensurate with the significance of the impacts
203 resulting from the discharge.¹⁴
- 204 i. Tier 3 projects include any discharge of dredged or fill material that directly impacts more
205 than two-tenths (0.2) of an acre or 300 linear feet of waters of the state, rare, threatened or
206 endangered species habitat in waters of the state, wetlands or eel grass beds, or
207 Outstanding National Resource Waters or Areas of Special Biological Significance, and is
208 not a project that inherently cannot be located at an alternate location. Tier 3 projects shall
209 provide an analysis of off-site and on-site alternatives.
- 210 ii. Tier 2 projects include any discharge of dredged or fill material that directly impacts more
211 than one tenth (0.1) and less than or equal to two tenths (0.2) of an acre or more than 100
212 and less than or equal to 300 linear feet of waters of the state unless it meets the criteria
213 for a Tier 3 project, or any project that inherently cannot be located at an alternate location
214 (unless it meets the size requirements set forth in Tier 1). Tier 2 projects shall provide an
215 analysis of only on-site alternatives. For routine operation and maintenance of existing
216 facilities, analysis of on-site alternatives is limited to operation and maintenance
217 alternatives for the facility.
- 218 iii. Tier 1 projects include any discharge of dredged or fill material that directly impacts less
219 than or equal to one tenth (0.1) of an acre or less than or equal to 100 linear feet of waters
220 of the state, unless it meets the criteria for a Tier 3 project. Tier 1 projects shall provide a
221 description of any steps that have been or will be taken to avoid and minimize loss of, or
222 significant adverse impacts to, beneficial uses of waters of the state.
- 223 2. Additional Information Required for a Complete Application
- 224 a. If required by the permitting authority on a case-by-case basis, supplemental field data from the
225 wet season to substantiate dry season delineations, as is consistent with the 1987 Manual and
226 Supplements.
- 227 b. If compensatory mitigation is required by the permitting authority, on a case-by-case basis, a
228 draft compensatory mitigation plan developed using a watershed approach containing the items
229 listed below. Compensatory mitigation plans are not required for Ecological Restoration and
230 Enhancement Projects. For permittees who intend to fulfill their compensatory mitigation

¹⁴ As used below, "impacts" include both permanent and temporary impacts.

Procedures for Discharges of Dredged or Fill Material to Waters of the State

- 231 obligations by securing credits from approved mitigation banks or in-lieu fee programs, their
232 mitigation plans need include only items i, ii, and iii, as described below, as well as information
233 required in the State Supplemental Dredge or Fill Guidelines, section 230.94 (c)(5) and (c)(6),
234 and the name of the specific mitigation bank or in-lieu fee program proposed to be used.
- 235 Draft compensatory mitigation plans shall comport with the State Supplemental Dredge or Fill
236 Guidelines, Subpart J, and include the items listed below.
- 237 i. A watershed profile for the project evaluation area for both the proposed dredged or fill
238 project and the proposed compensatory mitigation project.
 - 239 ii. An assessment of the overall condition of aquatic resources proposed to be impacted by
240 the project and their likely stressors, using an assessment method approved by the
241 permitting authority.
 - 242 iii. A description of how the project impacts and compensatory mitigation would not cause a
243 net loss of the overall abundance, diversity, and condition of aquatic resources, based on
244 the watershed profile. If the compensatory mitigation is located in the same watershed as
245 the project, no net loss will be determined on a watershed basis. If the compensatory
246 mitigation and project impacts are located in multiple watersheds, no net loss will be
247 determined considering all affected watersheds collectively. The level of detail in the plan
248 shall be sufficient to accurately evaluate whether compensatory mitigation offsets the
249 adverse impacts attributed to a project.
 - 250 iv. Preliminary information about ecological performance standards, monitoring, and long-term
251 protection and management, as described in the State Supplemental Dredge or Fill
252 Guidelines.
 - 253 v. A timetable for implementing the compensatory mitigation plan.
 - 254 vi. If the compensatory mitigation plan includes buffers, design criteria and monitoring
255 requirements for those buffers.
 - 256 vii. If the compensatory mitigation involves restoration or establishment as the form of
257 mitigation, applicants shall notify, as applicable, state and federal land management
258 agencies, airport land use commission, fire control districts, flood control districts, local
259 mosquito-vector control district(s), and any other interested local entities prior to initial site
260 selection. These entities should be notified as early as possible during the initial
261 compensatory mitigation project design stage.
 - 262 viii. If required by the permitting authority, an assessment of reasonably foreseeable impacts to
263 the compensatory mitigation associated with climate change, and any measures to avoid
264 or minimize those potential impacts.
- 265 c. If required by the permitting authority on a case-by-case basis, if project activities include in-
266 water work or water diversions, a proposed water quality monitoring plan to monitor compliance
267 with water quality objectives of the applicable water quality control plan. At a minimum, the plan
268 should include type and frequency of sampling for each applicable parameter.
 - 269 d. In all cases where temporary impacts are proposed, a draft restoration plan that outlines design,
270 implementation, assessment, and maintenance for restoring areas of temporary impact to pre-
271 project conditions. The design components shall include the objectives of the restoration plan;
272 grading plan of disturbed areas to pre-project contours; a planting palette with plant species

Procedures for Discharges of Dredged or Fill Material to Waters of the State

273 native to the area; seed collection locations; and an invasive species management plan. The
274 implementation component shall include all proposed actions to implement the plan (e.g., re-
275 contouring, initial planting, site stabilization, removal of temporary structures) and a schedule for
276 completing those actions. The maintenance and assessment components shall include a
277 description of performance standards used to evaluate attainment of objectives; the timeframe
278 for determining attainment of performance standards; and maintenance requirements (e.g.,
279 watering, weeding, replanting and invasive species control). If temporary impacts are proposed
280 to be restored through passive restoration, the draft restoration plan shall include an explanation
281 of how passive restoration will restore the area to pre-project conditions, assessment
282 components, and an estimated date for expected restoration. The level of detail in the
283 restoration plan shall be sufficient to accurately evaluate whether the restoration addresses the
284 adverse temporary impacts attributed to a project. The applicant shall submit a final restoration
285 plan that describes the restoration of all temporarily disturbed areas to pre-project conditions,
286 consistent with section IV.B.4.

287 For Ecological Restoration and Enhancement Projects, a restoration plan for temporary impacts
288 provided as part of the binding stream or wetland enhancement or restoration agreement or
289 wetland establishment agreement may satisfy this requirement.

290 e. For all Ecological Restoration and Enhancement Projects, a draft assessment plan including the
291 following: project objectives; description of performance standards used to evaluate attainment
292 of objectives; protocols for condition assessment; the timeframe and responsible party for
293 performing condition assessment; and assessment schedule. A draft assessment plan shall
294 provide for at least one assessment of the overall condition of aquatic resources and their likely
295 stressors, using an appropriate assessment method approved by the permitting authority, prior
296 to restoration and/or enhancement and two years following restoration and/or enhancement to
297 determine success of the restoration and/or enhancement. An assessment plan approved by a
298 federal or state resource agency, or a local agency with the primary function of managing land
299 or water for wetland habitat purposes in accordance with a binding stream or wetland
300 enhancement agreement, restoration agreement, or establishment agreement, will satisfy these
301 requirements. An assessment plan approved by a non-governmental conservation organization
302 or a state or federal agency that is statutorily tasked with natural resource management may
303 satisfy some or all of these requirements.

304 **B. Permitting Authority Review and Approval of Applications for Individual Orders**

- 305 1. The permitting authority will evaluate the potential impacts on the aquatic environment from the
306 proposed project and determine whether the proposed project complies with these Procedures.
307 The permitting authority has the discretion to approve a project only if the applicant has
308 demonstrated the following:
- 309 a. A sequence of actions has been taken to first avoid, then to minimize, and lastly compensate for
310 adverse impacts that cannot be practicably avoided or minimized to waters of the state;
 - 311 b. The potential impacts will not contribute to a net loss of the overall abundance, diversity, and
312 condition of aquatic resources in a watershed (or multiple watersheds when compensatory
313 mitigation is permitted in another watershed as set forth in section IV.B.5(d));
 - 314 c. The discharge of dredged or fill material will not violate water quality standards and will be
315 consistent with all applicable water quality control plans and policies for water quality control;
316 and
 - 317 d. The discharge of dredged or fill material will not cause or contribute to significant degradation of

Procedures for Discharges of Dredged or Fill Material to Waters of the State

318 the waters of the state.

319 2. The permitting authority shall rely on any final aquatic resource report verified by the Corps to
320 determine boundaries of waters of the U.S. For all other wetland area delineations, the permitting
321 authority shall review and approve delineations that are performed using the methods described in
322 section III.

323 3. Alternatives Analysis Review Requirements:

324 a. The purpose of the alternatives analysis is to identify the LEDPA. The permitting authority will
325 be responsible for determining the sufficiency of an alternatives analysis except as described in
326 3(b) below. In all cases, the alternatives analysis must establish that the proposed project
327 alternative is the LEDPA in light of all potential direct, secondary (indirect), and cumulative
328 impacts on the physical, chemical, and biological elements of the aquatic ecosystem.

329 b. Discharges to waters of the U.S.

330 In reviewing and approving the alternatives analysis for discharges of dredged or fill material
331 that impact waters of the U.S., the permitting authority shall defer to the Corps' determinations
332 on the adequacy of the alternatives analysis, or rely on a draft alternatives analysis if no final
333 determination has been made, unless the Executive Officer or Executive Director determines
334 that (1) the permitting authority was not provided an adequate opportunity to collaborate in the
335 development of the alternatives analysis, (2) the alternatives analysis does not adequately
336 address aquatic resource issues identified in writing by the Executive Officer or Executive
337 Director to the Corps during the development of the alternatives analysis, or (3) the proposed
338 project and all of the identified alternatives would not comply with water quality standards.

339 If the project also includes discharges to waters of the state outside of federal jurisdiction, the
340 permitting authority shall require the applicant to supplement the alternatives analysis to include
341 waters of the state outside of federal jurisdiction unless the applicant has consulted with the
342 permitting authority and the alternatives analysis addresses all issues identified by the
343 permitting authority during the consultation process. If an alternatives analysis is not required
344 by the Corps for discharges of dredged or fill material to waters of the U.S., the permitting
345 authority shall require an alternatives analysis for the entire project in accordance with the State
346 Supplemental Dredge or Fill Guidelines, unless the project is exempt under section IV.A.1(g)
347 above.

348 The permitting authority shall not apply the presumption set forth in the State Supplemental
349 Dredge or Fill Guidelines, section 230.10(a)(3) to any non-vegetated waters of the U.S. that the
350 Corps does not classify as a special aquatic site (as defined in subpart E of U.S. EPA's section
351 404(b)(1) Guidelines).

352 4. Prior to or concurrent with issuance of the Order, the permitting authority will approve the final
353 restoration plan for temporary impacts. Generally, the permitting authority will approve the final
354 restoration plan when it issues the Order. The permitting authority may approve the final restoration
355 plan after it issues the Order. In such cases the permitting authority shall include as a condition of
356 the Order that the applicant receive approval of the final restoration plan prior to initiating the
357 temporary impacts and shall specify a process for approving the final restoration plan.

358 5. Compensatory Mitigation

359 a. Compensatory mitigation, in accordance with the State Supplemental Dredge or Fill Guidelines,
360 Subpart J, may be required to ensure that an activity complies with these Procedures.
361 Consistent with section 230.93(a)(2) of the State Supplemental Guidelines, subject to the

Procedures for Discharges of Dredged or Fill Material to Waters of the State

362 permitting authority's approval, compensatory mitigation may be performed using methods of
363 restoration, enhancement, establishment, and in certain circumstances preservation.
364 Restoration should generally be the first option considered because the likelihood of success is
365 greater and the impacts to potentially ecologically important uplands are reduced compared to
366 establishment, and the potential gains in terms of aquatic resource functions are greater,
367 compared to enhancement and preservation.

368 b. Where feasible, the permitting authority will consult and coordinate with any other public
369 agencies that have concurrent mitigation requirements in order to achieve multiple
370 environmental benefits with a single mitigation project, thereby reducing the cost of compliance
371 to the applicant.

372 c. Amount: The amount of compensatory mitigation will be determined on a project-by-project
373 basis in accordance with State Supplemental Dredge or Fill Guidelines, section 230.93(f). The
374 permitting authority may take into account recent anthropogenic degradation to the aquatic
375 resource and the potential and existing functions and conditions of the aquatic resource. The
376 permitting authority may reduce the amount of compensatory mitigation if buffer areas adjacent
377 to the compensatory mitigation are also required to be maintained as part of the compensatory
378 mitigation management plan. The amount of compensatory mitigation required by the
379 permitting authority will vary depending on which of the following strategies the applicant uses to
380 locate the mitigation site within a watershed.

381 Strategy 1: Applicant locates compensatory mitigation using a watershed approach based on a
382 watershed profile developed from a watershed plan that: (1) has been approved for use by the
383 permitting authority and analyzed in an environmental document, (2) includes monitoring
384 provisions, and (3) includes guidance on compensatory mitigation opportunities.

385 Strategy 2: Applicant locates compensatory mitigation using a watershed approach based on a
386 watershed profile developed for a project evaluation area, and demonstrates that the mitigation
387 project will contribute to the sustainability of watershed functions and the overall health of the
388 watershed area's aquatic resources.

389 Generally, the amount of compensatory mitigation required under Strategy 1 will be less than
390 the amount of compensatory mitigation required under Strategy 2 since the level of certainty
391 that a compensatory mitigation project will meet its performance standards increases if the
392 compensatory mitigation project complies with a watershed plan as described above. Certainty
393 increases when there is a corresponding increase in understanding of watershed conditions,
394 which is increased when using a watershed plan as described above to determine
395 compensatory mitigation requirements.

396 A minimum of one-to-one mitigation ratio,¹⁵ measured as area or length, is required to
397 compensate for wetland or stream losses when compensatory mitigation is required. Subject to
398 the permitting authority's approval, the ratio may be satisfied using any of the methods identified
399 in section IV.B.5(a). A higher overall mitigation ratio shall be used where necessary to ensure
400 replacement of lost aquatic resource functions, as described in the State Supplemental Dredge
401 or Fill Guidelines, section 230.93(f). Where temporary impacts will be restored to pre-project
402 conditions, the permitting authority may require compensatory mitigation for temporal loss from
403 the temporary impacts.

¹⁵ For temporary impacts, the minimum one-to-one mitigation ratio for wetland or stream losses is not applicable for temporal losses for impacts that are fully restored to pre-project conditions.

Procedures for Discharges of Dredged or Fill Material to Waters of the State

404 d. Type and Location: The permitting authority will evaluate the applicant's proposed mitigation
405 type and location based on the applicant's use of a watershed approach based on a watershed
406 profile. The permitting authority will determine the appropriate type and location of
407 compensatory mitigation based on watershed conditions, impact size, location and spacing,
408 aquatic resource values, relevant watershed plans, and other considerations.

409 In general, the required compensatory mitigation should be located within the same watershed
410 as the impact site, but the permitting authority may approve compensatory mitigation in a
411 different watershed. For example, if a proposed project may affect more than one watershed,
412 then the permitting authority may determine that locating all required project mitigation in one
413 area is ecologically preferable to requiring mitigation within each watershed.

414 e. Final Compensatory Mitigation Plan: The permitting authority will review and approve the final
415 compensatory mitigation plan submitted by the applicant to ensure mitigation comports with the
416 State Supplemental Dredge or Fill Guidelines, Water Code requirements, applicable water
417 quality standards, and other appropriate requirements of state law. The level of detail in the
418 final plan shall be sufficient to accurately evaluate whether compensatory mitigation offsets the
419 adverse impacts attributed to a project considering the overall size and scope of impact. The
420 compensatory mitigation plan shall be sufficient to provide the permitting authority with a
421 reasonable assurance that replacement of the full range of lost aquatic resource(s) and/or
422 functions will be provided in perpetuity.

423 Generally, the permitting authority will approve the final compensatory mitigation plan when it
424 issues the Order. Where compliant with CEQA, the permitting authority may approve the final
425 compensatory mitigation plan after it issues the Order. In such cases the permitting authority
426 shall include as a condition of the Order that the applicant receive approval of the final mitigation
427 plan prior to discharging dredged or fill material to waters of the state and shall specify a
428 process for approving the final mitigation plan.

429 f. Financial Security: Where deemed necessary by the permitting authority, provision of a
430 financial security (e.g., letter of credit or performance bond) shall be a condition of the Order. In
431 this case, the permitting authority will approve the financial security to ensure compliance with
432 compensatory mitigation plan requirements. The financial security shall be in a form consistent
433 with the California Constitution and state law.

434 g. Term of Mitigation Obligation: The permitting authority may specify in the Order the conditions
435 that must be met in order for the permitting authority to release the permittee from the mitigation
436 obligation, including compensatory mitigation performance standards and long-term
437 management funding obligations.

438 6. The permitting authority shall provide public notice in accordance with Water Code section 13167.5
439 for waste discharge requirements. The permitting authority shall provide public notice of an
440 application for water quality certification in accordance with California Code of Regulations, title 23,
441 section 3858. If the permitting authority receives comments on the application or there is
442 substantial public interest in the project, the permitting authority shall also provide public notice of
443 the draft Order, or draft amendment of the Order, unless circumstances warrant otherwise.

444 7. The permitting authority will review and approve the final monitoring and reporting requirements for
445 all projects. Monitoring and reporting may be required to demonstrate compliance with the terms of
446 the Order.

447 C. General Orders

Procedures for Discharges of Dredged or Fill Material to Waters of the State

448 Discharges of dredged or fill material to waters of the state that are regulated under a general order are
449 not subject to the requirements set forth in sections IV.A and IV.B. Applicants applying to enroll under a
450 general order shall follow the instructions specified in the general order for obtaining coverage.

451 The permitting authority may issue general orders for specific classes of dredged or fill discharge
452 activities that are similar; involve the same or similar types of discharges and possible adverse impacts
453 requiring the same or similar conditions or limitations in order to alleviate potential adverse impacts to
454 water quality; and are determined by the permitting authority to more appropriately be regulated under
455 a general order rather than under an individual Order.

456 General orders shall be reviewed, noticed, and issued in accordance with the applicable requirements
457 of division 7 of the Water Code and the California Code of Regulations, division 3 of title 23.

458 **D. Activities and Areas Excluded from the Application Procedures for Regulation of** 459 **Discharges of Dredged or Fill Material to Waters of the State**

460 The application procedures specified in sections IV.A and IV.B do not apply to proposed discharges of
461 dredged or fill material to waters of the state from the following activities¹⁶ or to the following areas.
462 These exclusions do not, however, affect the Water Board's authority to issue or waive waste discharge
463 requirements (WDRs) or take other actions for the following activities or areas to the extent authorized
464 by the Water Code.

465 1. Activities excluded from application procedures in sections IV.A and IV.B:

466 a. Activities that are exempt under CWA section 404(f) (33 USC § 1344(f)).¹⁷ The permitting
467 authority shall use 33 CFR 323.4 (1986) and 40 CFR 232.3 (1988) to determine whether certain
468 activities are exempt under CWA section 404(f). These regulations are hereby incorporated by
469 reference and shall apply to all waters of the state. Consistent with CWA section 404(f)(2) and
470 40 CFR section 232.3, any discharge of dredged or fill material to a water of the state incidental
471 to any of these activities is not exempt under CWA section 404(f) and shall be subject to the
472 application procedures set forth in sections IV.A and IV.B, if (1) the purpose of the activity is
473 bringing a water of the state into a use to which it was not previously subject, where the flow or
474 circulation of water of the state may be impaired or the reach of such waters be reduced, or (2)
475 the discharge contains any toxic pollutant listed in CWA section 307.

476 b. Suction dredge mining activities for mineral recovery regulated under CWA section 402.

477 c. Routine and emergency operation and maintenance activities conducted by public agencies,
478 water utilities, or special districts that result in discharge of dredged or fill material to artificial,
479 existing waters of the state:

480 i. currently used and maintained primarily for one or more of the purposes listed in section
481 II.3.d. (ii), (iii), (iv), (x), or (xi); or

482 ii. for the purpose of preserving the line, grade, volumetric or flow capacity within the
483 existing footprint of a flood control or stormwater conveyance facility.

484 This exclusion does not relieve public agencies, water utilities or special districts of their

¹⁶ Note that not all activities identified in this section necessarily result in discharges of dredged or fill material to waters of the state.

¹⁷ Unless otherwise specified, all federal statutes and regulations that are incorporated by reference into these Procedures are the versions of those federal statutes and regulations that are in effect as of April 2, 2019.

Procedures for Discharges of Dredged or Fill Material to Waters of the State

485 obligation to submit an application for a water quality certification consistent with California
486 Code of Regulations, title 23, section 3856 or waste discharge requirements consistent with
487 Water Code section 13260, whichever is applicable, to the permitting authority for these
488 activities; or their responsibility to avoid and minimize adverse impacts to aquatic resources and
489 beneficial uses from these activities. The permitting authority has full discretion to determine
490 whether an activity described above qualifies for this exclusion based on the application
491 submitted and other relevant information. If the permitting authority determines that an activity
492 qualifies for this exclusion, the permitting authority retains full authority and discretion under the
493 Porter-Cologne Water Quality Control Act to determine how to regulate the discharge of
494 dredged or fill material. Where a permitting authority has already determined it appropriate to
495 regulate these types of activities in specific instances, this exclusion in no way disturbs or limits
496 the permitting authority's current regulation of these types of activities. This exclusion does not
497 apply to the discharge of dredged or fill material to a water of the state approved by an agency
498 as compensatory mitigation.

499 d. Routine operation and maintenance activities that result in discharge of dredged or fill material
500 to artificially-created waters currently used and maintained primarily for one or more of the
501 purposes listed in section II.3.d. (i), (ii), (iii), (vi), (vii), (x), or (xi). This exclusion does not apply
502 to the discharge of dredged or fill material to (a) a water of the U.S., (b) a water specifically
503 identified in a water quality control plan, (c) a water created by modification of a water of the
504 state, or (d) a water approved by an agency as compensatory mitigation.

505 2. Areas excluded from application procedures in sections IV.A and IV.B:

506 a. Wetland areas that qualify as prior converted cropland (PCC) within the meaning of 33 CFR
507 section 328.3(b)(2). The applicant may establish that the area is PCC by providing relevant
508 documentary evidence that the area qualifies as PCC and has not been abandoned due to five
509 consecutive years of non-use for agricultural purposes, or by providing a current PCC
510 certification by the Natural Resources Conservation Service, the Corps, or the U.S. EPA to the
511 permitting authority.

512 b. Wetlands that are, or have been, in rice cultivation (including wild rice) within the last five years
513 as of April 2, 2019 and have not been abandoned due to five consecutive years of non-use in
514 rice production.

515 c. The following features used for agricultural purposes:

516 i. Ditches with ephemeral flow that are not a relocated water of the state or excavated in a
517 water of the state;

518 ii. Ditches with intermittent flow that are not a relocated water of the state or excavated in a
519 water of the state, or that do not drain wetlands other than any wetlands described in
520 sections (iv) or (v);

521 iii. Ditches that do not flow, either directly or through another water, into another water of
522 the state;

523 iv. Artificially irrigated areas that would revert to dry land should application of waters to that
524 area cease; or

Procedures for Discharges of Dredged or Fill Material to Waters of the State

525 v. Artificial, constructed lakes and ponds created in dry land such as farm and stock
526 watering ponds, irrigation ponds, and settling basins.

527 The exclusions in section IV.D.2 do not apply to discharges of dredged or fill material that convert
528 wetland areas to a non-agricultural use.

529 For requests for approvals from the Division of Water Rights for activities associated with (1) an
530 appropriation of water subject to Part 2 (commencing with section 1200) of Division 2 of the Water
531 Code, (2) a hydroelectric facility where the proposed activity requires a Federal Energy Regulatory
532 Commission (FERC) license or amendment to a FERC license, or (3) any other diversion of water for
533 beneficial use where approval by the Division of Water Rights is required, the Division of Water Rights
534 will inform the applicant whether the application procedures in sections IV.A and IV.B will apply to the
535 application.

536 V. Definitions

537 The following definitions apply to these Procedures, including the State Supplemental Dredge or Fill
538 Guidelines. Unless otherwise indicated, any term that is not defined in these Procedures shall have the
539 same meaning as defined in Water Code section 13050, and title 23, section 3831 of the California
540 Code of Regulations.

541 **Abundance** means an estimate of the amount of aquatic resources by type in a watershed area, and
542 what types of aquatic resources are most and least prevalent.

543 **Active Surface Mining** means operations that, in accordance with Division 2, Chapter 9 of the Surface
544 Mining and Reclamation Act of 1975, have an approved reclamation plan, and for which reclamation
545 has not been certified as complete by the local lead agency with the concurrence of the Department of
546 Conservation.

547 **Alternatives Analysis** is the process of analyzing project alternatives, including the proposed project,
548 to determine the alternative that is the least environmentally damaging practicable alternative (LEDPA).

549 **Application** means a written request, including a report of waste discharge or request for water quality
550 certification, for authorization of any activity that may result in the discharge of dredged or fill material
551 and is subject to these Procedures.

552 **Discharge of Dredged or Fill Material** shall have the same meanings as they are used in the federal
553 Clean Water Act and 40 CFR section 232.2, but (1) shall include discharges to waters of the state that
554 are not waters of the U.S. and (2) any demonstrations described in 40 CFR section 232.2(3)(i) shall be
555 made to the permitting authority instead of the Corps or U.S. EPA. Placement of dredged or fill material
556 in a manner that could not affect the quality of waters of the state is not considered a discharge of
557 dredged or fill material.

558 **Diversity** means the relative proportion of aquatic resource types, classification, connectivity, and
559 spatial distribution in a watershed area.

560 **Ecological Restoration and Enhancement Project** means the project is voluntarily undertaken for the
561 purpose of assisting or controlling the recovery of an aquatic ecosystem that has been degraded,
562 damaged or destroyed to restore some measure of its natural condition and to enhance the beneficial
563 uses, including potential beneficial uses of water.

564 Such projects are undertaken:

Procedures for Discharges of Dredged or Fill Material to Waters of the State

- 565 1) in accordance with the terms and conditions of a binding stream or wetland enhancement or
566 restoration agreement, or a wetland establishment agreement, between the real property
567 interest owner or the entity conducting the habitat restoration or enhancement work and:
- 568 a. a federal or state resource agency, including, but not limited to, the U.S. Fish and
569 Wildlife Service, Natural Resources Conservation Service, Farm Service Agency,
570 National Marine Fisheries Service, National Oceanic and Atmospheric Administration,
571 U.S. Forest Service, U.S. Bureau of Land Management, California Department of Fish
572 and Wildlife, California Wildlife Conservation Board, California Coastal Conservancy or
573 the Delta Conservancy;
- 574 b. a local agency with the primary function of managing land or water for wetland habitat
575 purposes; or
- 576 c. a non-governmental conservation organization; or
- 577 2) by a state or federal agency that is statutorily tasked with natural resource management.

578 These projects do not include the conversion of a stream or natural wetland to uplands or stream
579 channelization. It is recognized that Ecological Restoration and Enhancement Projects may require
580 ongoing maintenance or management to maximize fish, wildlife, habitat, or other ecological benefits, or
581 filling gullied stream channels and similar rehabilitative activities to re-establish stream and meadow
582 hydrology. Changes in wetland plant communities that occur when wetland hydrology is more fully
583 restored during rehabilitation activities are not considered a conversion to another aquatic habitat type.
584 These projects also do not include actions required under a Water Board Order for mitigation, actions to
585 service required mitigation, or actions undertaken for the primary purpose of land development.

586 **Environmental Document** means a document prepared for compliance with the California
587 Environmental Quality Act (CEQA) or the National Environmental Policy Act (NEPA).

588 **Hydrophyte** means any macrophyte that grows in water or on a substrate that is at least periodically
589 deficient in oxygen as a result of excessive water content; plants typically found in wet habitats.

590 **LEDPA** means the least environmentally damaging practicable alternative. The determination of
591 practicable alternatives shall be consistent with the State Supplemental Dredge or Fill Guidelines,
592 section 230.10(a).

593 **Normal Circumstances** is the soil and hydrologic conditions that are normally present, without regard
594 to whether the vegetation has been removed. The determination of whether normal circumstances
595 exist in a disturbed area involves an evaluation of the extent and relative permanence of the physical
596 alteration of wetland hydrology and hydrophytic vegetation, and consideration of the purpose and cause
597 of the physical alterations to hydrology and vegetation.

598 **Order** means waste discharge requirements, waivers of waste discharge requirements, or water quality
599 certification.

600 **Permitting Authority** means the entity or person issuing the Order (i.e., the applicable Water Board,
601 Executive Director or Executive Officer, or his or her designee).

602 **Project** means the whole of an action that includes a discharge of dredged or fill material to waters of
603 the state.

604 **Project Evaluation Area** means an area that includes the project impact site, and/or the compensatory
605 mitigation site, and is sufficiently large to evaluate the effects of the project and/or the compensatory

Procedures for Discharges of Dredged or Fill Material to Waters of the State

606 mitigation on the abundance, diversity, and condition of aquatic resources in an ecologically meaningful
607 unit of the watershed. The size and location of the ecologically meaningful unit shall be based on a
608 reasonable rationale.

609 **Water Boards** mean any of the nine Regional Water Quality Control Boards, the State Water
610 Resources Control Board, or all of them collectively.

611 **Watershed** means a land area that drains to a common waterway, such as a stream, lake, estuary,
612 wetland, or ultimately the ocean.

613 **Watershed Approach** means an analytical process for evaluating the environmental effects of a
614 proposed project and making decisions that support the sustainability or improvement of aquatic
615 resources in a watershed. The watershed approach recognizes that the abundance, diversity, and
616 condition of aquatic resources in a watershed support beneficial uses. Diversity of aquatic resources
617 includes both the types of aquatic resources and the locations of those aquatic resources in a
618 watershed. Consideration is also given to understanding historic and potential aquatic resource
619 conditions, past and projected aquatic resource impacts in the watershed, and terrestrial connections
620 between aquatic resources. The watershed approach can be used to evaluate avoidance and
621 minimization of direct, secondary (indirect), and cumulative project impacts. It also can be used in
622 determining compensatory mitigation requirements.

623 **Watershed Plan** means a document, or a set of documents, developed in consultation with relevant
624 stakeholders, a specific goal of which is aquatic resource restoration, establishment, enhancement, and
625 preservation within a watershed. A watershed plan addresses aquatic resource conditions in the
626 watershed, multiple stakeholder interests, and land uses. Watershed plans should include information
627 about implementing the watershed plan. Watershed plans may also identify priority sites for aquatic
628 resource restoration and protection. Examples of watershed plans include special area management
629 plans, advance identification programs, and wetland management plans. The permitting authority may
630 approve the use of other plans, including for example, Habitat Conservation Plans (HCPs), Natural
631 Community Conservation Plans (NCCPs), or municipal stormwater permit watershed management
632 programs as watershed plans, if they substantially meet the stated above. Any NCCP approved by the
633 California Department of Fish and Wildlife before December 31, 2020, and any regional HCP approved
634 by the United States Fish and Wildlife Service before December 31, 2020, which includes biological
635 goals for aquatic resources, shall be used by the permitting authority as a watershed plan for such
636 aquatic resources, unless the permitting authority determines in writing that the HCP or NCCP does not
637 substantially meet the definition of a watershed plan for such aquatic resources.

638 **Watershed Profile** means a compilation of data or information on the abundance, diversity, and
639 condition of aquatic resources in a project evaluation area. The watershed profile shall include a map
640 and a report characterizing the location, abundance and diversity of aquatic resources in the project
641 evaluation area, assessing the condition of aquatic resources in the project evaluation area, and
642 describing the environmental stress factors affecting that condition.

643 The watershed profile shall include information sufficient to evaluate direct, secondary (indirect), and
644 cumulative impacts of project and factors that may favor or hinder the success of compensatory
645 mitigation projects and help define watershed goals. It may include such things as current trends in
646 habitat loss or conservation, cumulative impacts of past development activities, current development
647 trends, the presence and need of sensitive species, and chronic environmental problems or site
648 conditions such as flooding or poor water quality.

649 The scope and detail of the watershed profile shall be commensurate with the magnitude of impact
650 associated with the proposed project. Information sources include online searches, maps, watershed

Procedures for Discharges of Dredged or Fill Material to Waters of the State

651 plans, and possibly some fieldwork if necessary. In some cases, field data may need to be collected in
652 the project evaluation area to confirm the reported condition. Some or all of the information may be
653 obtained from a watershed plan. Watershed profiles for subsequent projects in a watershed can be
654 used to track the cumulative effectiveness of the permitting authority's decisions.

655 **Wetland Delineation** means the application of a technical and procedural method to identify the
656 boundary of a wetland area within a specified study site by identifying the presence or absence of
657 wetland indicators at multiple points at the site and by establishing boundaries that group together sets
658 of points that share the same status as wetland versus non-wetland.

Procedures for Discharges of Dredged or Fill Material to Waters of the State

659 **Appendix A: State Supplemental Dredge or Fill Guidelines**

660 It is the intent of the Water Boards to be consistent with the U.S. EPA's 404(b)(1) Guidelines where
661 feasible. Due to jurisdictional and procedural differences, some modifications to the U.S. EPA's
662 404(b)(1) Guidelines were necessary. Generally, these changes or deletions were made to reduce
663 redundancy (especially where sufficiently described elsewhere in these Procedures) and to account for
664 other state requirements. Note that the numbering scheme of the U.S. EPA's 404(b)(1) Guidelines has
665 been retained in these State Supplemental Dredge or Fill Guidelines for the benefit of practitioners who
666 are familiar with the U.S. EPA's 404(b)(1) Guidelines. The State Supplemental Dredge or Fill
667 Guidelines describe how the Water Boards will implement the U.S. EPA's 404(b)(1) Guidelines under
668 these Procedures. The definitions contained herein apply to these Procedures, including the State
669 Supplemental Dredge or Fill Guidelines.

670 **Subpart A – General**

671 § 230.3 Definitions.

672 For purposes of these Procedures, the following terms shall have the meanings indicated:

673 (c) The terms aquatic environment and aquatic ecosystem mean waters of the state, including
674 wetlands, that serve as habitat for interrelated and interacting communities and populations of
675 plants and animals.

676 (h) The term discharge point means the point within the disposal site at which the dredged or fill
677 material is released.

678 (i) The term disposal site means that portion of the "waters of the state" where the discharge of
679 dredged or fill material is permitted and involves a bottom surface area and any overlying volume of
680 water. In the case of wetlands or ephemeral streams on which surface water is not present, the
681 disposal site consists of the wetland or ephemeral stream surface area.

682 (k) The term extraction site means the place from which the dredged or fill material proposed for
683 discharge is to be removed.

684 (n) The term permitting authority means as defined above in the main text of these Procedures.

685 (q) The term practicable means available and capable of being done after taking into consideration
686 cost, existing technology, and logistics in light of overall project purposes.

687 (q1) Special aquatic sites means those sites identified in subpart E. Special aquatic sites are
688 geographic areas, large or small, possessing special ecological characteristics of productivity,
689 habitat, wildlife protection, or other important and easily disrupted ecological values. These areas
690 are generally recognized as significantly influencing or positively contributing to the general overall
691 environmental health or vitality of the entire ecosystem of a region. (See § 230.10 (a)(3))

692 § 230.6 Adaptability

693 (a) The manner in which these Guidelines are used depends on the physical, biological, and
694 chemical nature of the proposed extraction site, the material to be discharged, and the candidate
695 disposal site, including any other important components of the ecosystem being evaluated.
696 Documentation to demonstrate knowledge about the extraction site, materials to be extracted, and
697 the candidate disposal site is an essential component of guideline application. These Guidelines
698 allow evaluation and documentation for a variety of activities, ranging from those with large,
699 complex impacts on the aquatic environment to those for which the impact is likely to be innocuous.

Procedures for Discharges of Dredged or Fill Material to Waters of the State

700 It is unlikely that the Guidelines will apply in their entirety to any one activity, no matter how
701 complex. It is anticipated that substantial numbers of applications will be for minor, routine activities
702 that have little, if any, potential for significant degradation of the aquatic environment. It generally is
703 not intended or expected that extensive testing, evaluation or analysis will be needed to make
704 findings of compliance in such routine cases.

705 (b) The Guidelines user, including the agency or agencies responsible for implementing the
706 Guidelines, must recognize the different levels of effort that should be associated with varying
707 degrees of impact and require or prepare commensurate documentation. The level of
708 documentation should reflect the significance and complexity of the discharge activity.

709 (c) An essential part of the evaluation process involves making determinations as to the relevance
710 of any portion(s) of the Guidelines and conducting further evaluation only as needed. However,
711 where portions of the Guidelines review procedure are "short form" evaluations, there still must be
712 sufficient information (including consideration of both individual and cumulative impacts) to support
713 the decision of whether to specify the site for disposal of dredged or fill material and to support the
714 decision to curtail or abbreviate the evaluation process. The presumption against the discharge in
715 [§ 230.10](#) applies to this decision-making.

716 **Subpart B – Compliance with Guidelines**

717 § 230.10 Restrictions on Discharge

718 (a) No discharge of dredged or fill material shall be permitted if there is a practicable alternative to
719 the proposed discharge which would have less adverse impact on the aquatic ecosystem, so long
720 as the alternative does not have other significant adverse environmental consequences.

721 (1) For the purpose of this requirement, practicable alternatives include, but are not limited to:

722 (i) Activities which do not involve a discharge of dredged or fill material to waters of the state
723 or ocean waters;

724 (ii) Discharges of dredged or fill material at other locations in waters of the state or ocean
725 waters;

726 (2) An alternative is practicable if it is available and capable of being done after taking into
727 consideration cost, existing technology, and logistics in light of overall project purposes. If it is
728 otherwise a practicable alternative, an area not presently owned by the applicant which could
729 reasonably be obtained, utilized, expanded or managed in order to fulfill the basic purpose of
730 the proposed activity may be considered.

731 (3) Where activity associated with a discharge which is proposed for a special aquatic site (as
732 defined in subpart E) does not require access or proximity to or siting within the special aquatic
733 site in question to fulfill its basic purpose (i.e., is not "water dependent"), practicable alternatives
734 that do not involve special aquatic sites are presumed to be available, unless clearly
735 demonstrated otherwise. In addition, where a discharge is proposed for a special aquatic site,
736 all practicable alternatives to the proposed discharge which do not involve a discharge into a
737 special aquatic site are presumed to have less adverse impact on the aquatic ecosystem,
738 unless clearly demonstrated otherwise.

739 (b) No discharge of dredged or fill material shall be permitted if it:

740 (1) Causes or contributes, after consideration of disposal site dilution and dispersion, to
741 violations of any applicable State water quality standard;

Procedures for Discharges of Dredged or Fill Material to Waters of the State

742 (2) Violates any applicable toxic effluent standard or prohibition under section 307 of the Clean
743 Water Act;

744 (c) No discharge of dredged or fill material shall be permitted which will cause or contribute to
745 significant degradation of the waters of the state. Under these Guidelines, effects contributing to
746 significant degradation considered individually or collectively, include:

747 (1) Significantly adverse effects of the discharge of pollutants on human health or welfare,
748 including but not limited to effects on municipal water supplies, plankton, fish, shellfish, wildlife,
749 and special aquatic sites;

750 (2) Significantly adverse effects of the discharge of pollutants on life stages of aquatic life and
751 other wildlife dependent on aquatic ecosystems, including the transfer, concentration, and
752 spread of pollutants or their byproducts outside of the disposal site through biological, physical,
753 and chemical processes.

754 (3) Significantly adverse effects of the discharge of pollutants on aquatic ecosystem diversity,
755 productivity, and stability. Such effects may include, but are not limited to, loss of fish and
756 wildlife habitat or loss of the capacity of a wetland to assimilate nutrients, purify water, or reduce
757 wave energy; or

758 (4) Significantly adverse effects of the discharge of pollutants on recreational, aesthetic, and
759 economic values.

760 (d) No discharge of dredged or fill material shall be permitted unless appropriate and practicable
761 steps have been taken which will minimize potential adverse impacts of the discharge on the
762 aquatic ecosystem. Subpart H identifies such possible steps.

763 **Subpart E –Special Aquatic Sites**

764 § 230.40 Sanctuaries and refuges

765 (a) Sanctuaries and refuges consist of areas designated under State and Federal laws or local
766 ordinances to be managed principally for the preservation and use of fish and wildlife resources.

767 § 230.41 Wetlands.

768 (a)(1) Wetlands are as defined above in the main text of these Procedures.

769 § 230.42 Mud Flats.

770 (a) Mud flats are broad flat areas along the sea coast and in coastal rivers to the head of tidal influence
771 and inland lakes, ponds, and riverine systems. When mud flats are inundated, wind and wave action
772 may resuspend bottom sediments. Coastal mud flats are exposed at extremely low tides and
773 inundated at high tides with the water table at or near the surface of the substrate. The substrate of
774 mud flats contains organic material and particles smaller in size than sand. They are either
775 unvegetated or vegetated only by algal mats.

776 § 230.43 Vegetated shallows.

777 (a) Vegetated shallows are permanently inundated areas that under normal circumstances support
778 communities of rooted aquatic vegetation, such as turtle grass and eel grass in estuarine or marine
779 systems as well as a number of freshwater species in rivers and lakes.

780 § 230.45 Riffle and Pool Complexes.

781 (a) Steep gradient sections of streams are sometimes characterized by riffle and pool complexes. Such
782 stream sections are recognizable by their hydraulic characteristics. The rapid movement of water over

Procedures for Discharges of Dredged or Fill Material to Waters of the State

783 a coarse substrate in riffles results in a rough flow, a turbulent surface, and high dissolved oxygen
784 levels in the water. Pools are deeper areas associated with riffles. Pools are characterized by a slower
785 stream velocity, a streaming flow, a smooth surface, and a finer substrate. Riffle and pool complexes
786 are particularly valuable habitat for fish and wildlife.

787 **Subpart H – Actions to Minimize Adverse Effects**

788 Note: There are many actions which can be undertaken in response to 230.10(d) to minimize the
789 adverse effects of discharges of dredged or fill material. Some of these, grouped by type of activity,
790 are listed in this subpart. Additional criteria for compensation measures are provided in subpart J of
791 these Procedures.

792 § 230.70 Actions concerning the location of the discharge.

793 The effects of the discharge can be minimized by the choice of the disposal site. Some of the ways
794 to accomplish this are by:

- 795 (a) Locating and confining the discharge to minimize smothering of organisms;
- 796 (b) Designing the discharge to avoid a disruption of periodic water inundation patterns;
- 797 (c) Selecting a disposal site that has been used previously for dredged material discharge;
- 798 (d) Selecting a disposal site at which the substrate is composed of material similar to that being
799 discharged, such as discharging sand on sand or mud on mud;
- 800 (e) Selecting a disposal site, the discharge point, and the method of discharge to minimize the
801 extent of any plume;
- 802 (f) Designing the discharge of dredged or fill material to minimize or prevent the creation of standing
803 bodies of water in areas of normally fluctuating water levels, and minimize or prevent the drainage
804 of areas subject to such fluctuations.

805 § 230.71 Actions concerning the material to be discharged

806 The effects of a discharge can be minimized by treatment of, or limitations on the material itself, such
807 as:

- 808 (a) Disposal of dredged material in such a manner that physiochemical conditions are maintained,
809 and the potency and availability of pollutants are reduced.
- 810 (b) Limiting the solid, liquid, and gaseous components of material to be discharged at a particular
811 site;
- 812 (c) Adding treatment substances to the discharge material;
- 813 (d) Utilizing chemical flocculants to enhance the deposition of suspended particulates in diked
814 disposal areas.

815 § 230.72 Actions controlling the material after discharge.

816 The effects of the dredged or fill material after discharge may be controlled by:

- 817 (a) Selecting discharge methods and disposal sites where the potential for erosion, slumping or
818 leaching of materials into the surrounding aquatic ecosystem will be reduced. These sites or
819 methods include, but are not limited to:
 - 820 (1) Using containment levees, sediment basins, and cover crops to reduce erosions;

Procedures for Discharges of Dredged or Fill Material to Waters of the State

821 (2) Using lined containment areas to reduce leaching where leaching of chemical constituents
822 from the discharged material is expected to be a problem;

823 (b) Capping in-place contaminated material with clean material or selectively discharging the most
824 contaminated material first to be capped with the remaining material;

825 (c) Maintaining and containing discharged material properly to prevent point and nonpoint sources
826 of pollution;

827 (d) Timing the discharge to minimize impact, for instance during periods of unusual high-water
828 flows, wind, wave, and tidal actions.

829 § 230.73 Actions affecting the method of dispersion.

830 The effects of a discharge can be minimized by the manner in which it is dispersed, such as:

831 (a) Where environmentally desirable, distributing the dredged material widely in a thin layer at the
832 disposal site maintain natural substrate contours and elevation;

833 (b) Orienting a dredged or fill material mound to minimize undesirable obstruction to the water
834 current or circulation pattern, and utilizing natural bottom contours to minimize the size of the
835 mound;

836 (c) Using silt screens or other appropriate methods to confine suspended particulate/turbidity to a
837 small area where settling or removal can occur;

838 (d) Making use of currents and circulation patterns to mix, disperse and dilute the discharge;

839 (e) Minimizing water column turbidity by using a submerged diffuser system. A similar effect can be
840 accomplished by submerging pipeline discharges or otherwise releasing materials near the bottom;

841 (f) Selecting sites or managing discharges to confine and minimize the release of suspended
842 particulates to give decreased turbidity levels and to maintain light penetration for organisms;

843 (g) Setting limitations on the amount of material to be discharged per unit of time or volume of
844 receiving water.

845 § 230.74 Actions related to technology.

846 Discharge technology should be adapted to the needs of each site. In determining whether the
847 discharge operation sufficiently minimizes adverse environmental impacts, the applicant should
848 consider:

849 (a) Using appropriate equipment or machinery, including protective devices, and the use of such
850 equipment or machinery in activities related to the discharge of dredged or fill material;

851 (b) Employing appropriate maintenance and operation on equipment or machinery, including
852 adequate training, staffing, and working procedures;

853 (c) Using machinery and techniques that are especially designed to reduce damage to wetlands.
854 This may include machines equipped with devices that scatter rather than mound excavated
855 materials, machines with specially designed wheels or tracks, and the use of mats under heavy
856 machines to reduce wetland surface compaction and rutting;

857 (d) Designing access roads and channels spanning structures using culverts, open channels, and
858 diversions that will pass both low and high-water flows, accommodate fluctuating water levels, and
859 maintain circulation and faunal movement;

860 (e) Employing appropriate machinery and methods of transport of the material for discharge.

Procedures for Discharges of Dredged or Fill Material to Waters of the State

861 § 230.75 Actions affecting plant and animal populations.

862 Minimization of adverse effects on populations of plant and animals can be achieved by:

863 (a) Avoiding changes in water current and circulation patterns which would interfere with the
864 movement of animals;

865 (b) Selecting sites or managing discharges to prevent or avoid creating habitat conducive to the
866 development of undesirable predators or species which have a competitive edge ecologically over
867 indigenous plants or animals;

868 (c) Avoiding sites having unique habitat or other value, including habitat of threatened or
869 endangered species;

870 (d) Using planning and construction practices to institute habitat development and restoration to
871 produce a new or modified environmental state of higher ecological value by displacement of some
872 or all of the existing environmental characteristics. Habitat development and restoration techniques
873 can be used to minimize adverse impacts and to compensate for destroyed habitat. Additional
874 criteria for compensation measures are provided in subpart J of this part. Use techniques that have
875 been demonstrated to be effective in circumstances similar to those under consideration wherever
876 possible. Where proposed development and restoration techniques have not yet advanced to the
877 pilot demonstration stage, initiate their use on a small scale to allow corrective action if
878 unanticipated adverse impacts occur;

879 (e) Timing discharge to avoid spawning or migration seasons and other biologically critical time
880 periods;

881 (f) Avoiding the destruction of remnant natural sites within areas already affected by development.

882 § 230.76 Actions affecting human use.

883 Minimization of adverse effects on human use potential may be achieved by:

884 (a) Selecting discharge sites and following discharge procedures to prevent or minimize any
885 potential damage to the aesthetically pleasing features of the aquatic site (e.g. viewsapes),
886 particularly with respect to water quality;

887 (b) Selecting disposal sites which are not valuable as natural aquatic areas;

888 (c) Timing the discharge to avoid the seasons or periods when human recreational activity
889 associated with the aquatic site is most important;

890 (d) Following discharge procedures which avoid or minimize the disturbance of aesthetic features
891 on an aquatic site or ecosystem;

892 (e) Selecting sites that will not be detrimental or increase incompatible human activity, or require the
893 need for frequent dredge or fill maintenance activity in remote fish and wildlife areas;

894 (f) Locating the disposal site outside of the vicinity of a public water supply intake.

895 § 230.77 Other actions.

896 (a) In the case of fills, controlling runoff and other discharges from activities to be conducted on the
897 fill;

898 (b) In the case of dams, designing water releases to accommodate the needs of fish and wildlife;

899 (c) In dredging projects funded by Federal agencies other than the Corps of Engineers, maintain
900 desired water quality of the return discharge through agreement with the Federal funding authority

Procedures for Discharges of Dredged or Fill Material to Waters of the State

901 on scientifically defensible pollutant concentration levels in addition to any applicable water quality
902 standards;

903 (d) When a significant ecological change in the aquatic environment is proposed by the discharge of
904 dredged or fill material, the permitting authority should consider the ecosystem that will be lost as
905 well as the environmental benefits of the new system.

906 **Subpart J – Compensatory Mitigation for Losses of Aquatic Resources**

907 § 230.91 Purpose and general considerations.

908 (a) Purpose.

909 (1) The purpose of this subpart is to establish standards and criteria for the use of all types of
910 compensatory mitigation, including on-site and off-site permittee-responsible mitigation,
911 mitigation banks, and in-lieu fee mitigation to offset unavoidable impacts to waters of the state
912 authorized through the issuance of Orders.

913 (d) Accounting for regional variations. Where appropriate, the permitting authority shall account for
914 regional characteristics of aquatic resource types, functions and services when determining
915 performance standards and monitoring requirements for compensatory mitigation projects.

916 § 230.92 Definitions.

917 For the purposes of this subpart, the following terms are defined:

918 Adaptive management means the development of a management strategy that anticipates likely
919 challenges associated with compensatory mitigation projects and provides for the implementation of
920 actions to address those challenges, as well as unforeseen changes to those projects. It requires
921 consideration of the risk, uncertainty, and dynamic nature of compensatory mitigation projects and
922 guides modification of those projects to optimize performance. It includes the selection of appropriate
923 measures that will ensure that the aquatic resource functions are provided and involves analysis of
924 monitoring results to identify potential problems of a compensatory mitigation project and the
925 identification and implementation of measures to rectify those problems.

926 Buffer means an upland, wetland, and/or riparian area that protects and/or enhances aquatic
927 resource functions associated with waters of the state from disturbances associated with adjacent
928 land uses.

929 Compensatory mitigation means the restoration (re-establishment or rehabilitation), establishment
930 (creation), enhancement, and/or in certain circumstances preservation of aquatic resources for the
931 purposes of offsetting unavoidable adverse impacts which remain after all appropriate and practicable
932 avoidance and minimization has been achieved.

933 Compensatory mitigation project means compensatory mitigation implemented by the permittee as a
934 requirement of an Order (i.e., permittee-responsible mitigation), or by a mitigation bank or an in-lieu
935 fee program.

936 Condition means the relative ability of an aquatic resource to support and maintain a community of
937 organisms having a species composition, diversity, and functional organization comparable to
938 reference aquatic resources in the region.

939 Credit means a unit of measure (e.g., a functional or areal measure or other suitable metric)
940 representing the accrual or attainment of aquatic functions at a compensatory mitigation site. The
941 measure of aquatic functions is based on the resources restored, established, enhanced, or
942 preserved.

Procedures for Discharges of Dredged or Fill Material to Waters of the State

- 943 Days means calendar days.
- 944 Debit means a unit of measure (e.g., a functional or areal measure or other suitable metric)
945 representing the loss of aquatic functions at an impact or project site. The measure of aquatic
946 functions is based on the resources impacted by the authorized activity.
- 947 Enhancement means the manipulation of the physical, chemical, or biological characteristics of an
948 aquatic resource to heighten, intensify, or improve a specific aquatic resource function(s).
949 Enhancement results in the gain of selected aquatic resource function(s), but may also lead to a
950 decline in other aquatic resource function(s). Enhancement does not result in a gain in aquatic
951 resource area.
- 952 Establishment (creation) means the manipulation of the physical, chemical, or biological
953 characteristics present to develop an aquatic resource that did not previously exist at an upland site.
954 Establishment results in a gain in aquatic resource area and functions.
- 955 Functional capacity means the degree to which an area of aquatic resource performs a specific
956 function.
- 957 Functions means the physical, chemical, and biological processes that occur in ecosystems.
- 958 Impact means adverse effect.
- 959 In-kind means a resource of a similar structural and functional type to the impacted resource.
- 960 In-lieu fee program means a program involving the restoration, establishment, enhancement, and/or
961 preservation of aquatic resources through funds paid to a governmental or non-profit natural
962 resources management entity to satisfy compensatory mitigation requirements for Orders. Similar to
963 a mitigation bank, an in-lieu fee program sells compensatory mitigation credits to permittees whose
964 obligation to provide compensatory mitigation is then transferred to the in-lieu program sponsor.
965 However, the rules governing the operation and use of in-lieu fee programs are somewhat different
966 from the rules governing operation and use of mitigation banks. The operation and use of an in-lieu
967 fee program are governed by an in-lieu fee program instrument.
- 968 In-lieu fee program instrument means the legal document for the establishment, operation, and use of
969 an in-lieu fee program.
- 970 Instrument means mitigation banking instrument or in-lieu fee program instrument.
- 971 Mitigation bank means a site, or suite of sites, where resources (e.g., wetlands, streams, riparian
972 areas) are restored, established, enhanced, and/or preserved for the purpose of providing
973 compensatory mitigation for impacts authorized by Orders. In general, a mitigation bank sells
974 compensatory mitigation credits to permittees whose obligation to provide compensatory mitigation is
975 then transferred to the mitigation bank sponsor. The operation and use of a mitigation bank are
976 governed by a mitigation banking instrument.
- 977 Mitigation banking instrument means the legal document for the establishment, operation, and use of
978 an in-lieu fee program.
- 979 Off-site means an area that is neither located on the same parcel of land as the impact site, nor on a
980 parcel of land contiguous to the parcel containing the impact site.
- 981 On-site means an area located on the same parcel of land as the impact site, or on a parcel of land
982 contiguous to the impact site.
- 983 Out-of-kind means a resource of a different structural and functional type from the impacted resource.

Procedures for Discharges of Dredged or Fill Material to Waters of the State

- 984 Performance standards are observable or measurable physical (including hydrological), chemical
985 and/or biological attributes that are used to determine if a compensatory mitigation project meets its
986 objectives.
- 987 Permittee-responsible mitigation means an aquatic resource restoration, establishment,
988 enhancement, and/or preservation activity undertaken by the permittee (or an authorized agent or
989 contractor) to provide compensatory mitigation for which the permittee retains full responsibility.
- 990 Preservation means the removal of a threat to, or preventing the decline of, aquatic resources by an
991 action in or near those aquatic resources. This term includes activities commonly associated with the
992 protection and maintenance of aquatic resources through the implementation of appropriate legal and
993 physical mechanisms. Preservation does not result in a gain of aquatic resource area or functions.
- 994 Re-establishment means the manipulation of the physical, chemical, or biological characteristics of a
995 site with the goal of returning natural/historic functions to a former aquatic resource. Re-
996 establishment results in rebuilding a former aquatic resource and results in a gain in aquatic resource
997 area and functions.
- 998 Reference aquatic resources are a set of aquatic resources that represent the full range of variability
999 exhibited by a regional class of aquatic resources as a result of natural processes and anthropogenic
1000 disturbances.
- 1001 Rehabilitation means the manipulation of the physical, chemical, or biological characteristics of a site
1002 with the goal of repairing natural/historic functions to a degraded aquatic resource. Rehabilitation
1003 results in a gain in aquatic resource function, but does not result in a gain in aquatic resource area.
- 1004 Restoration means the manipulation of the physical, chemical, or biological characteristics of a site
1005 with the goal of returning natural/historic functions to a former or degraded aquatic resource. For the
1006 purpose of tracking net gains in aquatic resource area, restoration is divided into two categories: re-
1007 establishment and rehabilitation.
- 1008 Service area means the geographic area within which impacts can be mitigated at a specific
1009 mitigation bank or an in-lieu fee program, as designated in its instrument.
- 1010 Services mean the benefits that human populations receive from functions that occur in ecosystems.
- 1011 Sponsor means any public or private entity responsible for establishing, and in most circumstances,
1012 operating a mitigation bank or in-lieu fee program.
- 1013 Temporal loss is the time lag between the loss of aquatic resource functions caused by the permitted
1014 impacts and the replacement of aquatic resource functions at the compensatory mitigation site.
1015 Higher compensation ratios may be required to compensate for temporal loss. When the
1016 compensatory mitigation project is initiated prior to, or concurrent with, the permitted impacts, the
1017 permitting authority may determine that compensation for temporal loss is not necessary, unless the
1018 resource has a long development time.
- 1019 Watershed means a land area that drains to a common waterway, such as a stream, lake, estuary,
1020 wetland, or ultimately the ocean.
- 1021 Watershed approach is defined above in the main text of these Procedures.
- 1022 Watershed plan is defined above in the main text of these Procedures.
- 1023 § 230.93 General compensatory mitigation requirements.
- 1024 (a) General Considerations.

Procedures for Discharges of Dredged or Fill Material to Waters of the State

1025 (1) The fundamental objective of compensatory mitigation is to offset environmental losses
1026 resulting from unavoidable impacts to waters of the state authorized by Orders. The permitting
1027 authority must determine the compensatory mitigation to be required in an Order, based on
1028 what would be environmentally preferable. In making this determination, the permitting authority
1029 must assess the likelihood for ecological success and sustainability, and the location of the
1030 compensation site relative to the impact site and their significance within the watershed, and the
1031 costs of the compensatory mitigation project. In many cases, the environmentally preferable
1032 compensatory mitigation may be provided through mitigation banks or in-lieu fee programs
1033 because they usually involve consolidating compensatory mitigation projects where ecologically
1034 appropriate, consolidating resources, providing financial planning and scientific expertise (which
1035 often is not practical for permittee-responsible compensatory mitigation projects), reducing
1036 temporal losses of functions, and reducing uncertainty over project success. Compensatory
1037 mitigation requirements must be commensurate with the amount and type of impact that is
1038 associated with a particular Order. Applicants are responsible for proposing an appropriate
1039 compensatory mitigation option to offset unavoidable impacts.

1040 (2) Compensatory mitigation may be performed using methods of restoration, enhancement,
1041 establishment, and in certain circumstances preservation. Restoration should generally be the
1042 first option considered because the likelihood of success is greater and the impacts to
1043 potentially ecologically important uplands are reduced compared to establishment, and the
1044 potential gains in terms of aquatic resource functions are greater, compared to enhancement
1045 and preservation.

1046 (3) Compensatory mitigation projects may be sited on public or private lands. Credits for
1047 compensatory mitigation projects on public land must be based solely on aquatic resource
1048 functions provided by the compensatory mitigation project, over and above those provided by
1049 public programs already planned or in place. All compensatory mitigation projects must comply
1050 with the standards in section IV of these Procedures, if they are to be used to provide
1051 compensatory mitigation for activities authorized by Orders, regardless of whether they are sited
1052 on public or private lands and whether the sponsor is a governmental or private entity.

1053 (b) Type and location of compensatory mitigation.

1054 (1) In general, the required compensatory mitigation should be located within the same
1055 watershed as the impact site, and should be located where it is most likely to successfully
1056 replace lost functions and services, taking into account such watershed scale features as
1057 aquatic habitat diversity, habitat connectivity, relationships to hydrologic sources (including the
1058 availability of water rights), trends in land use, ecological benefits, and compatibility with
1059 adjacent land uses. When compensating for impacts to marine resources, the location of the
1060 compensatory mitigation site should be chosen to replace lost functions and services within the
1061 same marine ecological system (e.g., reef complex, littoral drift cell). Compensation for impacts
1062 to aquatic resources in coastal watersheds (watersheds that include a tidal water body) should
1063 also be located in a coastal watershed where practicable. Compensatory mitigation projects
1064 should not be located where they will increase risks to aviation by attracting wildlife to areas
1065 where aircraft-wildlife strikes may occur (e.g., near airports).

1066 (2) Mitigation bank credits. When permitted impacts are located within the service area of an
1067 approved mitigation bank, and the bank has the appropriate number and resource type of

Procedures for Discharges of Dredged or Fill Material to Waters of the State

1068 credits available, the permittee's compensatory mitigation requirements may be met by securing
1069 those credits from the sponsor. Since an approved instrument (including an approved mitigation
1070 plan and appropriate real estate and financial assurances) for a mitigation bank is required to be
1071 in place before its credits can begin to be used to compensate for authorized impacts, use of a
1072 mitigation bank can help reduce risk and uncertainty, as well as temporal loss of resource
1073 functions and services. Mitigation bank credits are not released for debiting until specific
1074 milestones associated with the mitigation bank site's protection and development are achieved,
1075 thus use of mitigation bank credits can also help reduce risk that mitigation will not be fully
1076 successful. Mitigation banks typically involve larger, more ecologically valuable parcels, and
1077 more rigorous scientific and technical analysis, planning and implementation than permittee-
1078 responsible mitigation. Also, development of a mitigation bank requires site identification in
1079 advance, project-specific planning, and significant investment of financial resources that is often
1080 not practicable for many in-lieu fee programs. For these reasons, the permitting authority
1081 should give preference to the use of mitigation bank credits when these considerations are
1082 applicable. However, these same considerations may also be used to override this preference,
1083 where appropriate, as, for example, where an in-lieu fee program has released credits available
1084 from a specific approved in-lieu fee project, or a permittee-responsible project will restore an
1085 outstanding resource based on rigorous scientific and technical analysis.

1086 (3) In-lieu fee program credits. Where permitted impacts are located within the service area of
1087 an approved in-lieu fee program, and the sponsor has the appropriate number and resource
1088 type of credits available, the permittee's compensatory mitigation requirements may be met by
1089 securing those credits from the sponsor. Where permitted impacts are not located in the service
1090 area of an approved mitigation bank, or the approved mitigation bank does not have the
1091 appropriate number and resource type of credits available to offset those impacts, in-lieu fee
1092 mitigation, if available, is generally preferable to permittee-responsible mitigation. In-lieu fee
1093 projects typically involve larger, more ecologically valuable parcels, and more rigorous scientific
1094 and technical analysis, planning and implementation than permittee-responsible mitigation.
1095 They also devote significant resources to identifying and addressing high-priority resource
1096 needs on a watershed scale, as reflected in their compensation planning framework. For these
1097 reasons, the permitting authority should give preference to in-lieu fee program credits over
1098 permittee-responsible mitigation, where these considerations are applicable. However, as with
1099 the preference for mitigation bank credits, these same considerations may be used to override
1100 this preference where appropriate. Additionally, in cases where permittee-responsible
1101 mitigation is likely to successfully meet performance standards before advance credits secured
1102 from an in-lieu fee program are fulfilled, the permitting authority should also give consideration
1103 to this factor in deciding between in-lieu fee mitigation and permittee-responsible mitigation.

1104 (4) Permittee-responsible mitigation under a watershed approach. Where permitted impacts
1105 are not in the service area of an approved mitigation bank or in-lieu fee program that has the
1106 appropriate number and resource type of credits available, permittee-responsible mitigation is
1107 the only option. Where practicable and likely to be successful and sustainable, the resource
1108 type and location for the required permittee-responsible compensatory mitigation should be
1109 determined using the principles of a watershed approach as outlined in paragraph (c) of this
1110 section.

1111 (5) Permittee-responsible mitigation through on-site and in-kind mitigation. In cases where a
1112 watershed approach is not practicable, the permitting authority should consider opportunities to

Procedures for Discharges of Dredged or Fill Material to Waters of the State

1113 offset anticipated aquatic resource impacts by requiring on-site and in-kind compensatory
1114 mitigation. The permitting authority must also consider the practicability of on-site
1115 compensatory mitigation and its compatibility with the proposed project.

1116 (6) Permittee-responsible mitigation through off-site and/or out-of-kind mitigation. If, after
1117 considering opportunities for on-site, in-kind compensatory mitigation as provided in paragraph
1118 (b)(5) of this section, the permitting authority determines that these compensatory mitigation
1119 opportunities are not practicable, are unlikely to compensate for the permitted impacts, or will be
1120 incompatible with the proposed project, and an alternative, practicable off-site and/or out-of-kind
1121 mitigation opportunity is identified that has a greater likelihood of offsetting the permitted
1122 impacts or is environmentally preferable to on-site or in-kind mitigation, the permitting authority
1123 should require that this alternative compensatory mitigation be provided.

1124 (c) Watershed approach to compensatory mitigation.

1125 (1) The permitting authority must use a watershed approach to establish compensatory
1126 mitigation requirements in Orders as described in the main text of the Procedures. Where a
1127 watershed plan is available, the permitting authority will determine whether the plan meets the
1128 definition of watershed plan in the Procedures and therefore is appropriate for use in the
1129 watershed approach for compensatory mitigation. In cases where the permitting authority
1130 determines that an appropriate watershed plan is available, the watershed approach should be
1131 based on that plan. Where no such plan is available, the watershed approach should be based
1132 on information provided by the project sponsor or available from other sources. The ultimate
1133 goal of a watershed approach is to maintain and improve the abundance, diversity, and
1134 condition of aquatic resources within watersheds through strategic selection of compensatory
1135 mitigation sites.

1136 (2) Considerations.

1137 (i) A watershed approach to compensatory mitigation considers the importance of condition,
1138 landscape position and resource type of compensatory mitigation projects for the
1139 sustainability of aquatic resource functions within the watershed. Such an approach
1140 considers how the condition, types, and locations of compensatory mitigation projects will
1141 provide the desired aquatic resource functions, and will continue to function over time in a
1142 changing landscape. It also considers the habitat requirements of important species, habitat
1143 loss or conversion trends, sources of watershed impairment, and current development
1144 trends, as well as the requirements of other regulatory and non-regulatory programs that
1145 affect the watershed, such as storm water management or habitat conservation programs. It
1146 includes the protection and maintenance of terrestrial resources, such as non-wetland
1147 riparian areas and uplands, when those resources contribute to or improve the overall
1148 ecological functioning of aquatic resources in the watershed. Compensatory mitigation
1149 requirements determined through the watershed approach should not focus exclusively on
1150 specific functions (e.g., water quality or habitat for certain species), but should provide, where
1151 practicable, the suite of functions typically provided by the affected aquatic resource.

1152 (ii) Locational factors (e.g., hydrology, surrounding land use) are important to the success of
1153 compensatory mitigation for impacted habitat functions and may lead to siting of such
1154 mitigation away from the project area. However, consideration should also be given to

Procedures for Discharges of Dredged or Fill Material to Waters of the State

1155 functions and services (e.g., water quality, flood control, shoreline protection) that will likely
1156 need to be addressed at or near the areas impacted by the permitted impacts.

1157 (iii) A watershed approach may include on-site compensatory mitigation, off-site
1158 compensatory mitigation (including mitigation banks or in-lieu fee programs), or a
1159 combination of on-site and off-site compensatory mitigation.

1160 (iv) A watershed approach to compensatory mitigation should include, to the extent
1161 practicable, inventories of historic and existing aquatic resources, including identification of
1162 degraded aquatic resources, and identification of immediate and long-term aquatic resource
1163 needs within watersheds that can be met through permittee-responsible mitigation projects,
1164 mitigation banks, or in-lieu fee programs. Planning efforts should identify and prioritize
1165 aquatic resource restoration, establishment, and enhancement activities, and preservation of
1166 existing aquatic resources that are important for maintaining or improving ecological functions
1167 of the watershed. The identification and prioritization of resource needs should be as specific
1168 as possible, to enhance the usefulness of the approach in determining compensatory
1169 mitigation requirements.

1170 (v) A watershed approach is not appropriate in areas where watershed boundaries do not
1171 exist, such as marine areas. In such cases, an appropriate spatial scale should be used to
1172 replace lost functions and services within the same ecological system (e.g., reef complex,
1173 littoral drift cell).

1174 (3) Information Needs.

1175 (i) In the absence of a watershed plan determined by the permitting authority under
1176 paragraph (c)(1) of this section to be appropriate for use in the watershed approach, the
1177 permitting authority will use a watershed approach based on analysis of information
1178 regarding watershed conditions (as identified in the watershed profile) and needs, including
1179 potential sites for aquatic resource restoration activities and priorities for aquatic resource
1180 restoration and preservation. Such information includes: Current trends in habitat loss or
1181 conversion; cumulative impacts of past development activities, current development trends,
1182 the presence and needs of sensitive species; site conditions that favor or hinder the success
1183 of compensatory mitigation projects; and chronic environmental problems such as flooding or
1184 poor water quality.

1185 (ii) This information may be available from sources such as wetland maps; soil surveys; U.S.
1186 Geological Survey topographic and hydrologic maps; aerial photographs; information on rare,
1187 endangered and threatened species and critical habitat; local ecological reports or studies;
1188 and other information sources that could be used to identify locations for suitable
1189 compensatory mitigation projects in the watershed.

1190 (iii) The level of information and analysis needed to support a watershed approach must be
1191 commensurate with the scope and scale of the proposed impacts requiring an Order, as well
1192 as the functions lost as a result of those impacts.

1193 (4) **Watershed Scale.** The size of watershed addressed using a watershed approach should not
1194 be larger than is appropriate to ensure that the aquatic resources provided through

Procedures for Discharges of Dredged or Fill Material to Waters of the State

- 1195 compensation activities will effectively compensate for adverse environmental impacts resulting
1196 from activities authorized by Orders. The permitting authority should consider relevant
1197 environmental factors and appropriate locally-developed standards and criteria when
1198 determining the appropriate watershed scale in guiding compensation activities.
- 1199 (d) Site selection.
- 1200 (1) The compensatory mitigation project site must be ecologically suitable for providing the
1201 desired aquatic resource functions. In determining the ecological suitability of the compensatory
1202 mitigation project site, the permitting authority must consider, to the extent practicable, the
1203 following factors:
- 1204 (i) Hydrological conditions, soil characteristics, and other physical and chemical
1205 characteristics;
- 1206 (ii) Watershed-scale features, such as aquatic habitat diversity, habitat connectivity, and
1207 other landscape scale functions;
- 1208 (iii) The size and location of the compensatory mitigation site relative to hydrologic sources
1209 (including the availability of water rights) and other ecological features;
- 1210 (iv) Compatibility with adjacent land uses and watershed management plans;
- 1211 (v) Reasonably foreseeable effects the compensatory mitigation project will have on
1212 ecologically important aquatic or terrestrial resources (e.g., shallow sub-tidal habitat, mature
1213 forests), cultural sites, or habitat for federally- or state-listed threatened and endangered
1214 species; and
- 1215 (vi) Other relevant factors including, but not limited to, development trends, anticipated land
1216 use changes, habitat status and trends, the relative locations of the impact and mitigation
1217 sites in the stream network, local or regional goals for the restoration or protection of
1218 particular habitat types or functions (e.g., re-establishment of habitat corridors or habitat for
1219 species of concern), water quality goals, floodplain management goals, and the relative
1220 potential for chemical contamination of the aquatic resources.
- 1221 (2) Permitting authorities may require on-site, off-site, or a combination of on-site and off-site
1222 compensatory mitigation to replace permitted losses of aquatic resource functions and services.
- 1223 (3) Applicants should propose compensation sites adjacent to existing aquatic resources or
1224 where aquatic resources previously existed.
- 1225 (e) Mitigation type.
- 1226 (1) In general, in-kind mitigation is preferable to out-of-kind mitigation because it is most likely to
1227 compensate for the functions and services lost at the impact site. For example, tidal wetland
1228 compensatory mitigation projects are most likely to compensate for unavoidable impacts to tidal
1229 wetlands, while perennial stream compensatory mitigation projects are most likely to

Procedures for Discharges of Dredged or Fill Material to Waters of the State

1230 compensate for unavoidable impacts to perennial streams. Thus, except as provided in
1231 paragraph (e)(2) of this section, the required compensatory mitigation shall be of a similar type
1232 to the affected aquatic resource.

1233 (2) If the permitting authority determines, using the watershed approach in accordance with
1234 paragraph (c) of this section that out-of-kind compensatory mitigation will serve the aquatic
1235 resource needs of the watershed, the permitting authority may authorize the use of such out-of-
1236 kind compensatory mitigation. The basis for authorization of out-of-kind compensatory
1237 mitigation must be documented in the administrative record for the Order action.

1238 (3) For difficult-to-replace resources (e.g., bogs, fens, springs, streams, vegetated seasonal
1239 wetlands, slope and seep wetlands, vernal pools, and wet meadows) if further avoidance and
1240 minimization is not practicable, the required compensation should be provided, if practicable,
1241 through in-kind rehabilitation, enhancement, or preservation since there is greater certainty that
1242 these methods of compensation will successfully offset permitted impacts.

1243 (f) Amount of compensatory mitigation.

1244 (1) If the permitting authority determines that compensatory mitigation is necessary to offset
1245 unavoidable impacts to aquatic resources, the amount of required compensatory mitigation
1246 must be, to the extent practicable, sufficient to replace lost aquatic resource functions. In cases
1247 where appropriate functional or condition assessment methods or other suitable metrics are
1248 available, these methods should be used where practicable to determine how much
1249 compensatory mitigation is required.

1250 (2) The permitting authority must require a mitigation ratio greater than one-to-one where
1251 necessary to account for the method of compensatory mitigation (e.g., preservation), the
1252 likelihood of success, differences between the functions lost at the impact site and the functions
1253 expected to be produced by the compensatory mitigation project, temporal losses of aquatic
1254 resource functions, the difficulty of restoring or establishing the desired aquatic resource type
1255 and functions, and/or the distance between the affected aquatic resource and the compensation
1256 site. The rationale for the required replacement ratio must be documented in the administrative
1257 record for the Order action.

1258 (3) If an in-lieu fee program will be used to provide the required compensatory mitigation, and
1259 the appropriate number and resource type of released credits are not available, the permitting
1260 authority must require sufficient compensation to account for the risk and uncertainty associated
1261 with in-lieu fee projects that have not been implemented before the permitted impacts have
1262 occurred.

1263 (g) Use of mitigation banks and in-lieu fee programs. Mitigation banks and in-lieu fee programs
1264 may be used to compensate for impacts to aquatic resources authorized by general Orders and
1265 individual Orders in accordance with the preference hierarchy in paragraph (b) of this section.
1266 Mitigation banks and in-lieu fee programs may also be used to satisfy requirements arising out of an
1267 enforcement action, such as supplemental environmental projects.

1268 (h) Preservation.

Procedures for Discharges of Dredged or Fill Material to Waters of the State

- 1269 (1) Preservation may be used to provide compensatory mitigation for activities authorized by
1270 Orders when all the following criteria are met:
- 1271 (i) The resources to be preserved provide important physical, chemical, or biological
1272 functions for the watershed;
- 1273 (ii) The resources to be preserved contribute significantly to the ecological sustainability of
1274 the watershed. In determining the contribution of those resources to the ecological
1275 sustainability of the watershed, the permitting authority must use appropriate quantitative
1276 assessment tools where available;
- 1277 (iii) Preservation is determined by the permitting authority to be appropriate and practicable;
- 1278 (iv) The resources are under threat of destruction or adverse modifications; and
- 1279 (v) The preserved site will be permanently protected through an appropriate real estate or
1280 other legal instrument (e.g., easement, title transfer to state resource agency or land trust).
- 1281 (2) Where preservation is used to provide compensatory mitigation, to the extent appropriate
1282 and practicable the preservation shall be done in conjunction with aquatic resource restoration,
1283 establishment, and/or enhancement activities. This requirement may be waived by the
1284 permitting authority where preservation has been identified as a high priority using a watershed
1285 approach described in paragraph (c) of this section, but compensation ratios shall be higher.
- 1286 (i) Buffers. The permitting authority may require the restoration, establishment,
1287 enhancement, and preservation, as well as the maintenance, of riparian areas and/or buffers
1288 around aquatic resources where necessary to ensure the long-term viability of those
1289 resources. Buffers may also provide habitat or corridors necessary for the ecological
1290 functioning of aquatic resources. If buffers are required by the permitting authority as part of
1291 the compensatory mitigation project, compensatory mitigation credit will be provided for those
1292 buffers, as provided in section IV B.5 (c).
- 1293 (j) Relationship to other federal, tribal, state, and local programs.
- 1294 (1) Compensatory mitigation projects for Orders may also be used to satisfy the environmental
1295 requirements of other programs, such as tribal, state, or local wetlands regulatory programs,
1296 other federal programs such as the Surface Mining Control and Reclamation Act, Corps civil
1297 works projects, and Department of Defense military construction projects, consistent with the
1298 terms and requirements of these programs and subject to the following considerations:
- 1299 (i) The compensatory mitigation project must include appropriate compensation required by
1300 the Order for unavoidable impacts to aquatic resources authorized by that Order.
- 1301 (ii) Under no circumstances may the same credits be used to provide mitigation for more than
1302 one permitted activity. However, where appropriate, compensatory mitigation projects,
1303 including mitigation banks and in-lieu fee projects, may be designed to holistically address
1304 requirements under multiple programs and authorities for the same activity.

Procedures for Discharges of Dredged or Fill Material to Waters of the State

1305 (2) Except for projects undertaken by federal agencies, or where federal funding is specifically
1306 authorized to provide compensatory mitigation, federally-funded aquatic resource restoration or
1307 conservation projects undertaken for purposes other than compensatory mitigation, such as the
1308 Wetlands Reserve Program, Conservation Reserve Program, and Partners for Wildlife Program
1309 activities, cannot be used for the purpose of generating compensatory mitigation credits for
1310 activities authorized by Orders. However, compensatory mitigation credits may be generated by
1311 activities undertaken in conjunction with, but supplemental to, such programs in order to
1312 maximize the overall ecological benefits of the restoration or conservation project.

1313 (3) Compensatory mitigation projects may also be used to provide compensatory mitigation
1314 under the federal and state Endangered Species Act or for Natural Community Conservation
1315 Plans and Habitat Conservation Plans, as long as they comply with the requirements of
1316 paragraph (j)(1) of this section.

1317 (k) Order conditions.

1318 (1) The compensatory mitigation requirements for an Order, including the amount and type of
1319 compensatory mitigation, must be clearly stated in the special conditions of the individual Order
1320 or authorization to use the general Order. The special conditions must be enforceable.

1321 (2) For an Order that requires permittee-responsible mitigation, the special conditions must:

1322 (i) Identify the party responsible for providing the compensatory mitigation;

1323 (ii) Incorporate, by reference, the final or draft mitigation plan approved by the permitting
1324 authority;

1325 (iii) State the objectives, performance standards, and monitoring required for the
1326 compensatory mitigation project, unless they are provided in the approved final mitigation
1327 plan; and

1328 (iv) Describe any required financial assurances or long-term management provisions for the
1329 compensatory mitigation project, unless they are specified in the approved final mitigation
1330 plan.

1331 (4) If a mitigation bank or in-lieu fee program is used to provide the required compensatory
1332 mitigation, the special conditions must indicate whether a mitigation bank or in-lieu fee program
1333 will be used, and specify the number and resource type of credits the permittee is required to
1334 secure. In the case of an individual Order, the special condition must also identify the specific
1335 mitigation bank or in-lieu fee program that will be used. For authorizations to use a general
1336 Order, the special conditions may either identify the specific mitigation bank or in-lieu fee
1337 program, or state that the specific mitigation bank or in-lieu fee program used to provide the
1338 required compensatory mitigation must be approved by the permitting authority before the
1339 credits are secured.

1340 (l) Party responsible for compensatory mitigation.

Procedures for Discharges of Dredged or Fill Material to Waters of the State

1341 (1) For permittee-responsible mitigation, the special conditions of the Order must clearly
1342 indicate the party or parties responsible for the implementation, performance, and long-term
1343 management of the compensatory mitigation project.

1344 (3) If use of a mitigation bank or in-lieu fee program is approved by the permitting authority to
1345 provide part or all of the required compensatory mitigation for an Order, the permittee retains
1346 responsibility for providing the compensatory mitigation until the appropriate number and
1347 resource type of credits have been secured from a sponsor and the permitting authority has
1348 received documentation that confirms that the sponsor has accepted the responsibility for
1349 providing the required compensatory mitigation. This documentation may consist of a letter or
1350 form signed by the sponsor, with the Order number and a statement indicating the number and
1351 resource type of credits that have been secured from the sponsor. Copies of this
1352 documentation will be retained in the administrative records for both the Order and the
1353 instrument. If the sponsor fails to provide the required compensatory mitigation, the permitting
1354 authority may pursue measures against the sponsor to ensure compliance.

1355 (m) Timing. Implementation of the compensatory mitigation project shall be, to the maximum extent
1356 practicable, in advance of or concurrent with the activity causing the authorized impacts. The
1357 permitting authority shall require, to the extent appropriate and practicable, additional compensatory
1358 mitigation to offset temporal losses of aquatic functions that will result from the permitted activity.

1359 (n) Financial assurances.

1360 (1) The permitting authority shall require sufficient financial assurances to ensure a high level of
1361 confidence that the compensatory mitigation project will be successfully completed, in
1362 accordance with applicable performance standards. In cases where an alternate mechanism is
1363 available to ensure a high level of confidence that the compensatory mitigation will be provided
1364 and maintained (e.g., a formal, documented commitment from a government agency or public
1365 authority) the permitting authority may determine that financial assurances are not necessary for
1366 that compensatory mitigation project.

1367 (2) The amount of the required financial assurances must be determined by the permitting
1368 authority, in consultation with the project sponsor, and must be based on the size and
1369 complexity of the compensatory mitigation project, the degree of completion of the project at the
1370 time of project approval, the likelihood of success, the past performance of the project sponsor,
1371 and any other factors the permitting authority deems appropriate. Financial assurances may be
1372 in the form of performance bonds, escrow accounts, casualty insurance, letters of credit,
1373 legislative appropriations for government sponsored projects, or other appropriate instruments,
1374 subject to the approval of the permitting authority. The rationale for determining the amount of
1375 the required financial assurances must be documented in the administrative record for either the
1376 Order or the instrument. In determining the assurance amount, the permitting authority shall
1377 consider the cost of providing replacement mitigation, including costs for land acquisition,
1378 planning and engineering, legal fees, mobilization, construction, and monitoring.

1379 (3) If financial assurances are required, the Order must include a special condition requiring the
1380 financial assurances to be in place prior to commencing the permitted activity.

Procedures for Discharges of Dredged or Fill Material to Waters of the State

- 1381 (4) Financial assurances shall be phased out once the compensatory mitigation project has
1382 been determined by the permitting authority to be successful in accordance with its performance
1383 standards. The Order or instrument must clearly specify the conditions under which the
1384 financial assurances are to be released to the permittee, sponsor, and/or other financial
1385 assurance provider, including, as appropriate, linkage to achievement of performance
1386 standards, adaptive management, or compliance with special conditions.
- 1387 (5) A financial assurance must be in a form that ensures that the permitting authority will receive
1388 notification at least 120 days in advance of any termination or revocation. For third-party
1389 assurance providers, this may take the form of a contractual requirement for the assurance
1390 provider to notify the permitting authority at least 120 days before the assurance is revoked or
1391 terminated.
- 1392 (6) Financial assurances shall be payable at the direction of the permitting authority to his
1393 designee or to a standby trust agreement. When a standby trust is used (e.g., with performance
1394 bonds or letters of credit) all amounts paid by the financial assurance provider shall be
1395 deposited directly into the standby trust fund for distribution by the trustee in accordance with
1396 the permitting authority's instructions.
- 1397 (o) Compliance with applicable law. The compensatory mitigation project must comply with all
1398 applicable federal, state, and local laws. The Order, mitigation banking instrument, or in-lieu fee
1399 program instrument must not require participation by the permitting authority in project
1400 management, including receipt or management of financial assurances or long-term financing
1401 mechanisms, except as determined by the permitting authority to be consistent with its statutory
1402 authority, mission, and priorities.
- 1403 § 230.94 Planning and documentation.
- 1404 (a) Pre-application consultations. Potential applicants for Orders are encouraged to participate in
1405 pre-application meetings with the permitting authority and appropriate agencies to discuss potential
1406 mitigation requirements and information needs.
- 1407 (c) Mitigation plan.
- 1408 (1) Preparation and Approval.
- 1409 (i) For individual Orders, the permittee must prepare a draft mitigation plan and submit it to
1410 the permitting authority for review prior to issuing the Order. After addressing any comments
1411 provided by the permitting authority, the permittee must prepare a final mitigation plan, which
1412 must be approved by the permitting authority prior to commencing work in waters of the state.
1413 The approved final mitigation plan must be incorporated into the individual Order either as an
1414 attachment or by reference. The final mitigation plan must include the items described in
1415 paragraphs (c)(2) through (c)(14) of this section, but the level of detail of the mitigation plan
1416 should be commensurate with the scale and scope of the impacts. As an alternative, the
1417 permitting authority may determine that it would be more appropriate to address any of the
1418 items described in paragraphs (c)(2) through (c)(14) of this section as Order conditions,
1419 instead of components of a compensatory mitigation plan. For permittees who intend to fulfill
1420 their compensatory mitigation obligations by securing credits from approved mitigation banks
1421 or in-lieu fee programs, their mitigation plans need include only the items described in
1422 paragraphs (c)(5) and (c)(6) of this section, and the name of the specific mitigation bank or
1423 in-lieu fee program to be used.

Procedures for Discharges of Dredged or Fill Material to Waters of the State

- 1424 (2) Objectives. A description of the resource type(s) and amount(s) that will be provided, the
1425 method of compensation (i.e., restoration, establishment, enhancement, and/or preservation),
1426 and the manner in which the resource functions of the compensatory mitigation project will
1427 address the needs of the watershed, ecoregion, physiographic province, or other geographic
1428 area of interest.
- 1429 (3) Site selection. A description of the factors considered during the site selection process.
1430 This should include consideration of watershed needs, on-site alternatives where applicable,
1431 and the practicability of accomplishing ecologically self-sustaining aquatic resource restoration,
1432 establishment, enhancement, and/or preservation at the compensatory mitigation project site.
1433 (See [§ 230.93\(d\)](#).)
- 1434 (4) Site protection instrument. A description of the legal arrangements and instrument, including
1435 site ownership, that will be used to ensure the long-term protection of the compensatory
1436 mitigation project site (see [§ 230.97\(a\)](#)).
- 1437 (5) Baseline information. A description of the ecological characteristics of the proposed
1438 compensatory mitigation project site and, in the case of an application for an Order, the impact
1439 site. This may include descriptions of historic and existing plant communities, historic and
1440 existing hydrology, soil conditions, a map showing the locations of the impact and mitigation
1441 site(s) or the geographic coordinates for those site(s), and other site characteristics appropriate
1442 to the type of resource proposed as compensation. The baseline information should also
1443 include a delineation of waters of the state on the proposed compensatory mitigation project
1444 site. A prospective permittee planning to secure credits from an approved mitigation bank or in-
1445 lieu fee program only needs to provide baseline information about the impact site, not the
1446 mitigation bank or in-lieu fee project site.
- 1447 (6) Determination of credits. A description of the number of credits to be provided, including a
1448 brief explanation of the rationale for this determination. (See [§ 230.93\(f\)](#).)
- 1449 (i) For permittee-responsible mitigation, this should include an explanation of how the
1450 compensatory mitigation project will provide the required compensation for unavoidable
1451 impacts to aquatic resources resulting from the permitted activity.
- 1452 (ii) For permittees intending to secure credits from an approved mitigation bank or in-lieu fee
1453 program, it should include the number and resource type of credits to be secured and how
1454 these were determined.
- 1455 (7) Mitigation work plan. Detailed written specifications and work descriptions for the
1456 compensatory mitigation project, including, but not limited to, the geographic boundaries of the
1457 project; construction methods, timing, and sequence; source(s) of water, including connections
1458 to existing waters and uplands; methods for establishing the desired plant community; plans to
1459 control invasive plant species; the proposed grading plan, including elevations and slopes of the
1460 substrate; soil management; and erosion control measures. For stream compensatory
1461 mitigation projects, the mitigation work plan may also include other relevant information, such as
1462 planform geometry, channel form (e.g., typical channel cross-sections), watershed size, design
1463 discharge, and riparian area plantings.
- 1464 (8) Maintenance plan. A description and schedule of maintenance requirements to ensure the
1465 continued viability of the resource once initial construction is completed.
- 1466 (9) Performance standards. Ecologically-based standards that will be used to determine
1467 whether the compensatory mitigation project is achieving its objectives. (See [§ 230.95](#).)

Procedures for Discharges of Dredged or Fill Material to Waters of the State

1468 (10) Monitoring requirements. A description of parameters to be monitored in order to
1469 determine if the compensatory mitigation project is on track to meet performance standards and
1470 if adaptive management is needed. A schedule for monitoring and reporting on monitoring
1471 results to the permitting authority must be included. (See [§ 230.96.](#))

1472 (11) Long-term management plan. A description of how the compensatory mitigation project will
1473 be managed after performance standards have been achieved to ensure the long-term
1474 sustainability of the resource, including long-term financing mechanisms and the party
1475 responsible for long-term management. (See [§ 230.97\(d\).](#))

1476 (12) Adaptive management plan. A management strategy to address unforeseen changes in
1477 site conditions or other components of the compensatory mitigation project, including the party
1478 or parties responsible for implementing adaptive management measures. The adaptive
1479 management plan will guide decisions for revising compensatory mitigation plans and
1480 implementing measures to address both foreseeable and unforeseen circumstances that
1481 adversely affect compensatory mitigation success. (See [§ 230.97\(c\).](#))

1482 (13) Financial assurances. A description of financial assurances that will be provided and how
1483 they are sufficient to ensure a high level of confidence that the compensatory mitigation project
1484 will be successfully completed, in accordance with its performance standards (see [§ 230.93\(n\).](#))

1485 (14) Other information. The permitting authority may require additional information as
1486 necessary to determine the appropriateness, feasibility, and practicability of the compensatory
1487 mitigation project.

1488 § 230.95 Ecological performance standards.

1489 (a) The approved mitigation plan must contain performance standards that will be used to assess
1490 whether the project is achieving its objectives. Performance standards should relate to the
1491 objectives of the compensatory mitigation project, so that the project can be objectively evaluated to
1492 determine if it is developing into the desired resource type, providing the expected condition or
1493 functions, and attaining any other applicable metrics (e.g., acres).

1494 (b) Performance standards must be based on attributes that are objective and verifiable. Ecological
1495 performance standards must be based on the best available science that can be measured or
1496 assessed in a practicable manner. Performance standards may be based on variables or
1497 measures of functional capacity or condition as described in assessment methodologies,
1498 measurements of hydrology or other aquatic resource characteristics, and/or comparisons to
1499 reference aquatic resources of similar type and landscape position. The use of reference aquatic
1500 resources to establish performance standards will help ensure that those performance standards
1501 are reasonably achievable, by reflecting the range of variability exhibited by the regional class of
1502 aquatic resources as a result of natural processes and anthropogenic disturbances. Performance
1503 standards based on measurements of hydrology should take into consideration the hydrologic
1504 variability exhibited by reference aquatic resources, especially wetlands. Where practicable,
1505 performance standards should take into account the expected stages of the aquatic resource
1506 development process, in order to allow early identification of potential problems and appropriate
1507 adaptive management.

1508 § 230.96 Monitoring.

1509 (a) General.

1510 (1) Monitoring the compensatory mitigation project site is necessary to determine if the project is
1511 meeting its performance standards, and to determine if measures are necessary to ensure that

Procedures for Discharges of Dredged or Fill Material to Waters of the State

1512 the compensatory mitigation project is accomplishing its objectives. The submission of
1513 monitoring reports to assess the development and condition of the compensatory mitigation
1514 project is required, but the content and level of detail for those monitoring reports must be
1515 commensurate with the scale and scope of the compensatory mitigation project, as well as the
1516 compensatory mitigation project type. The mitigation plan must address the monitoring
1517 requirements for the compensatory mitigation project, including the parameters to be monitored,
1518 the length of the monitoring period, the party responsible for conducting the monitoring, the
1519 frequency for submitting monitoring reports to the permitting authority, and the party responsible
1520 for submitting those monitoring reports to the permitting authority.

1521 (2) The permitting authority may conduct site inspections on a regular basis (e.g., annually)
1522 during the monitoring period to evaluate mitigation site performance.

1523 (b) Monitoring period. The mitigation plan must provide for a monitoring period that is sufficient to
1524 demonstrate that the compensatory mitigation project has met performance standards, but not less
1525 than five years. A longer monitoring period must be required for aquatic resources with slow
1526 development rates (e.g., forested wetlands, bogs). Following project implementation, the permitting
1527 authority may reduce or waive the remaining monitoring requirements upon a determination that the
1528 compensatory mitigation project has achieved its performance standards. Conversely the
1529 permitting authority may extend the original monitoring period upon a determination that
1530 performance standards have not been met or the compensatory mitigation project is not on track to
1531 meet them. The permitting authority may also revise monitoring requirements when remediation
1532 and/or adaptive management is required.

1533 (c) Monitoring reports.

1534 (1) The permitting authority must determine the information to be included in monitoring reports.
1535 This information must be sufficient for the permitting authority to determine how the
1536 compensatory mitigation project is progressing towards meeting its performance standards, and
1537 may include plans (such as as-built plans), maps, and photographs to illustrate site conditions.
1538 Monitoring reports may also include the results of functional, condition, or other assessments
1539 used to provide quantitative or qualitative measures of the functions provided by the
1540 compensatory mitigation project site.

1541 (2) The permittee or sponsor is responsible for submitting monitoring reports in accordance with
1542 the special conditions of the Order or the terms of the instrument. Failure to submit monitoring
1543 reports in a timely manner may result in compliance action by the permitting authority.

1544 (3) Monitoring reports must be provided by the permitting authority to interested federal, tribal,
1545 state, and local resource agencies, and the public, upon request.

1546 § 230.97 Management.

1547 (a) Site protection.

1548 (1) The aquatic habitats, riparian areas, buffers, and uplands that comprise the overall
1549 compensatory mitigation project must be provided long-term protection through real estate
1550 instruments or other available mechanisms, as appropriate. Long-term protection may be
1551 provided through real estate instruments such as conservation easements held by entities such
1552 as federal, tribal, state, or local resource agencies, non-profit conservation organizations, or
1553 private land managers; the transfer of title to such entities; or by restrictive covenants. For
1554 government property, long-term protection may be provided through state or federal facility
1555 management plans or integrated natural resources management plans. When approving a
1556 method for long-term protection of non-government property other than transfer of title, the

Procedures for Discharges of Dredged or Fill Material to Waters of the State

1557 permitting authority shall consider relevant legal constraints on the use of conservation
1558 easements and/or restrictive covenants in determining whether such mechanisms provide
1559 sufficient site protection. To provide sufficient site protection, a conservation easement or
1560 restrictive covenant should, where practicable, establish in an appropriate third party (e.g.,
1561 governmental or non-profit resource management agency) the right to enforce site protections
1562 and provide the third party the resources necessary to monitor and enforce these site
1563 protections.

1564 (2) The real estate instrument, management plan, or other mechanism providing long-term
1565 protection of the compensatory mitigation site must, to the extent appropriate and practicable,
1566 prohibit incompatible uses (e.g., clear cutting or mineral extraction) that might otherwise
1567 jeopardize the objectives of the compensatory mitigation project. Where appropriate, multiple
1568 instruments recognizing compatible uses (e.g., fishing or grazing rights) may be used.

1569 (3) The real estate instrument, management plan, or other long-term protection mechanism
1570 must contain a provision requiring 60-day advance notification to the permitting authority before
1571 any action is taken to void or modify the instrument, management plan, or long-term protection
1572 mechanism, including transfer of title to, or establishment of any other legal claims over, the
1573 compensatory mitigation site.

1574 (4) For compensatory mitigation projects on public lands, where state or Federal facility
1575 management plans or integrated natural resources management plans are used to provide long-
1576 term protection, and changes in statute, regulation, or agency needs or mission results in an
1577 incompatible use on public lands originally set aside for compensatory mitigation, the public
1578 agency authorizing the incompatible use is responsible for providing alternative compensatory
1579 mitigation that is acceptable to the permitting authority for any loss in functions resulting from
1580 the incompatible use.

1581 (5) A real estate instrument, management plan, or other long-term protection mechanism used
1582 for site protection of permittee-responsible mitigation must be approved by the permitting
1583 authority in advance of, or concurrent with, the activity causing the authorized impacts.

1584 (b) Sustainability. Compensatory mitigation projects shall be designed, to the maximum extent
1585 practicable, to be self-sustaining once performance standards have been achieved. This includes
1586 minimization of active engineering features (e.g., pumps) and appropriate siting to ensure that
1587 natural hydrology and landscape context will support long-term sustainability. Where active long-
1588 term management and maintenance are necessary to ensure long-term sustainability (e.g.,
1589 prescribed burning, invasive species control, maintenance of water control structures, easement
1590 enforcement), the responsible party must provide for such management and maintenance. This
1591 includes the provision of long-term financing mechanisms where necessary. Where needed, the
1592 acquisition and protection of water rights must be secured and documented in the Order conditions
1593 or instrument.

1594 (c) Adaptive management.

1595 (1) If the compensatory mitigation project cannot be constructed in accordance with the
1596 approved mitigation plans, the permittee or sponsor must notify the permitting authority. A
1597 significant modification of the compensatory mitigation project requires approval from the
1598 permitting authority.

1599 (2) If monitoring or other information indicates that the compensatory mitigation project is not
1600 progressing towards meeting its performance standards as anticipated, the responsible party
1601 must notify the permitting authority as soon as possible. The permitting authority will evaluate

Procedures for Discharges of Dredged or Fill Material to Waters of the State

1602 and pursue measures to address deficiencies in the compensatory mitigation project. The
1603 permitting authority will consider whether the compensatory mitigation project is providing
1604 ecological benefits comparable to the original objectives of the compensatory mitigation project.

1605 (3) The permitting authority, in consultation with the responsible party (and other federal, tribal,
1606 state, and local agencies, as appropriate), will determine the appropriate measures. The
1607 measures may include site modifications, design changes, revisions to maintenance
1608 requirements, and revised monitoring requirements. The measures must be designed to ensure
1609 that the modified compensatory mitigation project provides aquatic resource functions
1610 comparable to those described in the mitigation plan objectives.

1611 (4) Performance standards may be revised in accordance with adaptive management to
1612 account for measures taken to address deficiencies in the compensatory mitigation project.
1613 Performance standards may also be revised to reflect changes in management strategies and
1614 objectives if the new standards provide for ecological benefits that are comparable or superior to
1615 the approved compensatory mitigation project. No other revisions to performance standards will
1616 be allowed except in the case of natural disasters.

1617 (d) Long-term management.

1618 (1) The Order conditions or instrument must identify the party responsible for ownership and all
1619 long-term management of the compensatory mitigation project. The Order conditions or
1620 instrument may contain provisions allowing the permittee or sponsor to transfer the long-term
1621 management responsibilities of the compensatory mitigation project site to a land stewardship
1622 entity, such as a public agency, non-governmental organization, or private land manager, after
1623 review and approval by the permitting authority. The land stewardship entity need not be
1624 identified in the original Order or instrument, as long as the future transfer of long-term
1625 management responsibility is approved by the permitting authority.

1626 (2) A long-term management plan should include a description of long-term management
1627 needs, annual cost estimates for these needs, and identify the funding mechanism that will be
1628 used to meet those needs.

1629 (3) Any provisions necessary for long-term financing must be addressed in the original Order or
1630 instrument. The permitting authority may require provisions to address inflationary adjustments
1631 and other contingencies, as appropriate. Appropriate long-term financing mechanisms include
1632 non-wasting endowments, trusts, contractual arrangements with future responsible parties, and
1633 other appropriate financial instruments. In cases where the long-term management entity is a
1634 public authority or government agency, that entity must provide a plan for the long-term
1635 financing of the site.

1636 (4) For permittee-responsible mitigation, any long-term financing mechanisms must be
1637 approved in advance of the activity causing the authorized impacts.