

DOCKETED

Docket Number:	09-AFC-07C
Project Title:	Palen Solar Power Project - Compliance
TN #:	200861
Document Title:	2012 DRAFT Rangelwide Mojave Desert Tortoise Monitoring
Description:	USFWS report
Filer:	Ileene Anderson
Organization:	Center for Biological Diversity
Submitter Role:	Intervenor
Submission Date:	10/15/2013 11:48:14 PM
Docketed Date:	10/16/2013

**RANGE-WIDE MONITORING OF
THE MOJAVE DESERT
TORTOISE (*GOPHERUS
AGASSIZII*):**

2012 ANNUAL REPORT

**PREPARED BY LINDA ALLISON
DESERT TORTOISE MONITORING COORDINATOR
U.S. FISH AND WILDLIFE SERVICE**

DECEMBER 2012

DRAFT

Recommended Citation: U.S. Fish and Wildlife Service. 2012. DRAFT Range-wide Monitoring of the Mojave Desert Tortoise (*Gopherus agassizii*): 2012 Annual Report. Report by the Desert Tortoise Recovery Office, U.S. Fish and Wildlife Service, Reno, Nevada.

TABLE OF CONTENTS

Executive Summary	5
Introduction.....	8
Methods.....	9
Study areas and transect locations	9
Distance sampling transect completion	10
Proportion of tortoises available for detection by line distance sampling, G_0	15
Field observer training	16
Telemetry training.....	16
Distance sampling training	16
Data management including quality assurance and quality control.....	23
Tortoise encounter rate and development of detection functions	24
Proportion of available tortoises detected on the transect centerline, $g(0)$	24
Estimates of tortoise density	26
Estimating the area of each stratum sampled and the number of tortoises in that area	27
Modification of previous procedures	28
Debriefing to describe strengths and weaknesses of project preparation and execution..	28
Results.....	29
Field observer training	29
Proportion of tortoises detected at distances from the transect centerline.....	29
Quality assurance and quality control.....	33
Transect completion.....	34
Burrow encounter rate.....	39
Tortoise encounter rates and detection functions.....	39
Proportion of tortoises available for detection by line distance sampling, G_0	42
Proportion of available tortoises detected on the transect centerline, $g(0)$	42
Estimates of tortoise density	43
Area of each stratum sampled and the number of tortoises in that area	45
Evaluating transect classification.....	45
Number of tortoises based on density and area sampled	46
Possibility of using burrow counts to estimate tortoise abundance at smaller scales	46
Debriefing to identify strengths and weaknesses in preparation for future years.....	47
Earlier review of the first field data	47
Training improvements to make more effective use of same time period.....	47
Discussion.....	48
Quality control and quality assurance.....	48
Improving ability to detect trends in desert tortoise abundance	48
Consequences of insufficient transects	48
Literature Cited	50

LIST OF TABLES

Table 1. Training schedule for 2012.....	18
Table 2. Proportion of tortoise models detected by teams within 1-, 2-, or 5-m of the transect centerline. Values that scored below the target of 0.90 at 1- and 2-m are highlighted.....	30
Table 3. Diagnostics for individual teams after training.....	31
Table 4. Number and type of transects in each stratum.....	34
Table 5. Availability of tortoises (G_0) during the period in 2012 when transects were walked in each group of neighboring strata.....	42
Table 6. Recovery unit and stratum-level encounters and densities in 2012 for tortoises with $MCL \geq 180\text{mm}$	44
Table 7. Estimated density of desert tortoises in monitored areas of each recovery unit in the Mojave and Colorado deserts in 2012.	45
Table 8. Transects completed other than as planned and any resulting reclassification.....	45
Table 9. Estimated tortoise abundance in sampled areas of each stratum.....	46

LIST OF FIGURES

Figure 1. Sampled areas 2012.....	11
Figure 2. Planned (dotted lines) and reflected transect paths at administrative boundaries, now also applied to stratum edges. A) One-corner reflection. B) Two-corner reflection.	14
Figure 3. Data flow from collection through final products.	23
Figure 4. Relationship between single-observer detections (by the leader, p) and dual-observer (team) detections, $g(0)$	25
Figure 5. Process for developing density estimates in 2012. For each type of estimate, the full set of data was subdivided appropriately, as indicated by columns.	27
Figure 6. Detection curves for each of the 2012 GBI teams during training. Curves are based on 16 km trials with approximately 100 detections.	32
Figure 7. Detection curves for each of the 2012 IWS trainee teams. Curves are based on 16 km trials with approximately 100 detections.	32
Figure 8. Detection curves for each of the 2012 Kiva trainee teams. Curves are based on 16 km trials with approximately 100 detections.	33
Figure 9. Distribution of distance sampling transects and live tortoise observations in the Northeastern Mojave Recovery Unit (Coyote Springs Valley, Mormon Mesa, Beaver Dam Slope, and Gold Butte-Pakoon monitoring strata).	35
Figure 10. Distribution of distance sampling transects and live tortoise observations in the Eastern Mojave Recovery Unit (Eldorado Valley and Ivanpah monitoring strata) and the eastern part of the Colorado Desert Recovery Unit (Fenner, Piute Valley, and Chemehuevi monitoring strata).....	36

Figure 11. Distribution of distance sampling transects and live tortoise observations in the Western Mojave Recovery Unit (Fremont-Kramer, Superior-Cronese, and Ord-Rodman monitoring strata).....	37
Figure 12. Distribution of distance sampling transects and live tortoise observations in the western portion of the Colorado Desert Recovery Unit (Pinto Mountains, Joshua Tree, Chocolate Mountain, and Chuckwalla monitoring strata).	38
Figure 13. Observed detections (histogram) and the resulting detection function (smooth curve) for live tortoises with $MCL \geq 180\text{mm}$ found by GBI. This curve uses only the 151 observations found within 21 m of the line.	40
Figure 14. Observed detections (histogram) and the resulting detection function (smooth curve) for live tortoises with $MCL \geq 180$ mm found by IWS. This curve uses only the 28 observations found within 15 m of the line.	41
Figure 15. Observed detections (histogram) and the resulting detection function (smooth curve) for live tortoises with $MCL \geq 180\text{mm}$ found by Kiva. This curve uses only the 73 observations found within 14 m of the line.	41
Figure 16. Detection pattern for the leader (p) and by the team ($g(0)$) based on all observations out to a given distance (x) from the centerline in 2012. Note convergence of $g(0)$ on 1.0 as x goes to 0.	43
Figure 17. Scatter plot of stratum densities and burrow encounter rate.	47

ACKNOWLEDGEMENTS

Funding or support-in-kind was provided by Fort Irwin National Training Center, the National Park Service portion of Grand Canyon-Parashant National Monument, Joshua Tree National Park, and the Arizona Strip Office of the Bureau of Land Management.

The original design for this project and considerations for optimizing it based on new information and experience were first set out in Anderson and Burnham (1996) and Anderson et al. (2001). Estimation methods were further refined during a 2008 workshop for distance sampling held by S. Buckland, L. Thomas, T. Marques, E. Rexstad, and D. Harris in Marshall, California.

Personnel from Kiva Biological Consulting (California) led by I. Daly, from the Institute for Wildlife Studies (California and Nevada) led by N. Gregory, and from the Great Basin Institute (Nevada, Arizona, and the Beaver Dam Slope of Utah) led by T. Christopher conducted the field surveys. The field monitors from these teams and from Joshua Tree National Park who did the hard work of collecting and verifying the data were: A. Allan, L. Baierl, M. Bassett, A. Bennett, B. Benson, Z. Cava, I. Daly, A. d'Eprenesnil, L. Drake, L. Driscoll, T. Faust, K. Foley, P. Fuchs, M. Grode, J. Hamner, K. Harding, M. Immel, A. Johnson, G. Johnson, S. Johnson, M. Jolliff, K. Lalumiere, D. Leach, J. Macmillan, J. MacNaughton, C. Martin, C. McClurg, C. Michaud, N. Mickle, L. Mjos, B. O'Brien, C. O'Leary, A. Paulson, T. Rode, S. Root, J. Ross, R. Salonikios, D. Sanderson, W. Schultz, J. Scott, C. Sherwood, B. Sparks, A. Sturgill, T. Sullivan, D. Summers, C. Taylor, S. Vaghini, M. Vamstad, J. Werba, K. Wickwire, D. Wolfson.

R. Patil (University of Nevada, Reno); M. Brenneman (Topoworks); J. Johnson (Arizona Exotic Animal Hospital); K. Dutcher (San Diego Zoo); J. Stegmeier and R. Parr (Las Vegas Field Office, BLM); and T. Christopher and B. Sparks (GBI) provided specialized training instruction for field crews. D. Zeff, R. Schultz, and L. Powell (Mojave Desert Ecosystem Program) provided preseason GIS support as well as independent review and post-processing of data submitted by both field groups. R. Patil (University of Nevada, Reno) updated the electronic data-collection forms and procedures used in 2012. M. Brenneman developed GIS tools for correctly reflecting transect paths and tracking completion of assigned transects. She also developed the final databases.

EXECUTIVE SUMMARY

The recovery program for desert tortoises in the Mojave and Colorado deserts (USFWS, 2011a) requires range-wide, long-term monitoring to determine whether recovery goals are met. Specifically, will population trends within recovery units increase for a period of 25 years? In 1999, the Desert Tortoise Management Oversight Group endorsed the use of line distance sampling (Buckland et al., 2001) as the method for estimating range-wide desert tortoise density. From 2001 to 2005, and again from 2007 through 2012, desert tortoise populations in 4 of the 5 recovery units have been part of a coordinated, range-wide monitoring program using line distance sampling. (The Upper Virgin River Recovery Unit is monitored by Utah Division of Wildlife Resources (McLuckie et al., 2012).) The first 5 years of monitoring culminated in a summary report (USFWS, 2006) that included eleven recommendations, seven of which were tied to functioning of the monitoring program and are paraphrased here:

1. The range-wide monitoring program should continue under a formal study plan subject to scientific review.
2. Refine [line distance sampling] techniques to improve sampling efficiency and estimates of trends.
3. Evaluate the spatial scale of the monitoring program.
4. Improve training lines.
5. Evaluate the use of independent field teams to improve data consistency and quality.
6. Refine and formalize/document the QA/QC process.
7. Identify and assess options for securing continued funding for range-wide population monitoring.

This report describes the full set of quality assurance steps and final results for the 2012 monitoring effort. The above issues continue to drive review and improvement of the program, so that reporting also addresses these aspects of the annual effort. The range-wide monitoring effort is directed each year at 16 strata that will be used to describe long-term trends. Data were collected on transects by field personnel working with three different groups, Kiva Biological (10 personnel), the Institute for Wildlife Studies (11 personnel), and Great Basin Institute (26 personnel). Four personnel from Joshua Tree National Park also collected telemetry data in the Park. After an intensive, 12-day specialized training session, crews completed 631 transects (7102 km) between 13 March and 22 May. In the course of these surveys, they reported 236 live tortoises.

Training is provided each year so that field crews are familiar with the specifics of distance sampling. Training also ensures consistency between the many crews collecting data. Inexperienced crews as well as those with prior experience participated in preseason training and testing provided by the USFWS. Crews were passed after demonstrating appropriate detection

patterns (including detection proportion on the transect line), measurement accuracy from tortoise models to the transect line, and other skills.

Four parameter estimates contribute to final reported tortoise densities in each monitoring stratum. The basis for distance sampling is the estimation of the number of tortoises detected at increasing distances from the walked transect. As surveyors look farther from the transect centerline, they will detect fewer and fewer of the tortoises that are actually there, so describing the way detections decrease with distance allows for estimation of the proportion that were present but not detected within a given distance of the centerline. Second, an estimate is made of the proportion above ground or visible in their burrows and available to be detected on transects. Third, the first two estimates are combined with the number of tortoises encountered per kilometer walked to provide the actual density in each stratum. Finally, the proportion detected on the line must be estimated. Unless all tortoises were detected on the centerline, the density estimate must be adjusted to account for the occurrence of these additional tortoises.

Separate detection curves were used to describe the decreasing ability of each team to see tortoises that were farther from the walked transect line. These detection curves will capture any differences between teams in application of the protocol, but are mostly expected to reflect the terrain as well as the extent to which vegetation obscures the view in different parts of the range, since the curves account for tortoises that were present in the same area but not seen. Kiva crews detected 39.7% of tortoises within 14 m of the transect centerline, GBI detected 37.9% out to 21 m, and IWS detected 43.2% to 15 m. The proportion of tortoises that were visible to be counted (G_0) varied in different parts of the range, which were surveyed at different times during the spring season. Visibility was as high as 97.7% in the Coyote Springs Valley telemetry site during 2 weeks in April. During the same weeks, visibility at the Superior-Cronese telemetry sites was also very high, 93.5%. Visibility in Joshua Tree National Park during the last 2 weeks of March and first 2 of April was 94.5%. Unfortunately, during this time visibility was as low as 35.6% in Chemehuevi, and was persistently low during the usual weeks of activity in the eastern part of the range. Visibility is related to encounter rates, which combined with a lack of funds from the BLM in Nevada and California to result in few observations in the central part of the Mojave Desert. On average, crews walked 30.1 km for each tortoise that was observed, but this number varied considerably from one monitoring stratum to the next. Although densities in the Northeastern Mojave Recovery Unit had been estimated at less than $2/\text{km}^2$ in some previous years, the density was estimated at $3.4/\text{km}^2$, similar to the past 4 years. The Western and Eastern Mojave recovery units had few transects and therefore few observations in some strata; density estimates in 4 strata were based on 1 or 2 observations. These strata had extremely low density estimates which were presumably an artifact of the sampling regime but nonetheless depressed recovery-unit level density estimates as well. Note that funding commitments in California were withdrawn 2 weeks before crews were on the ground so it was not possible to alter the sampling design that was already in place.

To enable field crews to complete transects in previously unsampled areas within strata, a set of guidelines was implemented in 2008 and 2009 for shortening transects in areas with rugged terrain or other obstacles (USFWS 2012a). These rules did enable crews to sample entire strata in a more representative way; however, in areas of California with lower funding, the resulting substrata never had enough transects or tortoise observations to separately evaluate tortoise densities in flat compared to rugged terrain. For this reason, in 2010, a more relaxed protocol was implemented in California, under which transects were to be completed to the extent possible along the original 12 km path. Mountainous terrain in the path was circumnavigated without searching for tortoises, then the path was resumed when possible. This method samples walkable areas representatively and also allows the proportion of unwalkable terrain to be estimated; in 2012 it was implemented throughout the range. Our density estimates are applicable to the estimated 88.3% of terrain in critical habitat and other areas managed for tortoises that is also walkable.

RANGE-WIDE MONITORING OF THE MOJAVE DESERT TORTOISE 2012

INTRODUCTION

The Mojave Desert population of the desert tortoise (*Gopherus agassizii*) was listed as threatened under the Endangered Species Act in 1990. This group of desert tortoises north and west of the Colorado River are now recognized as the species *G. agassizii*, separate from *G. morafkai* south and east of the Colorado River (Murphy et al., 2011). The revised recovery plan (USFWS, 2011a) designates five recovery units and to which decisions about continued listing status should be applied. The recovery plan specifies that consideration of delisting should only proceed when populations in each recovery unit have increased for at least one tortoise generation (25 years), and the only means to determine trend is by a rigorous program of long-term monitoring. Before the tortoise was listed, populations were monitored either using strip transects (Luckenbach, 1982) where indications of tortoise presence (live or dead tortoises, scats, burrows, or tracks) were converted to tortoise abundance categories based on calibration transects conducted in areas of better-known tortoise density, or by using capture-recapture population estimates on a limited number of (usually) 1-mi² study plots (Berry and Nicholson, 1984). Although data have continued to be collected on transects and study plots in recent years, these methods suffer statistical deficiencies and/or logistical constraints that render them unsuited for monitoring trends in abundance applicable to entire recovery units (Corn, 1994; Anderson et al., 2001; Tracy et al., 2004). In 1999 the Desert Tortoise Management Oversight Group endorsed the use of line distance sampling (Buckland et al., 2001) for estimating range-wide desert tortoise density.

Distance sampling methods use measurements taken from the center of the transect lines to tortoises to model detection as a function of distance from the walked path; tortoises farther from the travelled path have a lower probability of detection. In order to anchor the curve and estimate the true (not relative) proportion of tortoises detection within a given distance from the center of the transect, all tortoises must be detected on the transect centerline (Anderson et al., 2001; Buckland et al., 2001). There are additional assumptions in distance analysis – that distance is measured to the point where the animal was first detected and that distance is measured accurately – but these are easily satisfied in line distance sampling of desert tortoises. The assumption that detection at the centerline of the transect is perfect, however, can be violated during line distance sampling of tortoises, but the use of two observers minimizes the probability that tortoises are missed on the centerline and provides a correction factor in the form of an estimate of the number of tortoises on the line that were missed (USFWS, 2009).

Distance methods have been applied to estimate abundance of Sonoran Desert Tortoises (*G. morafkai*) since 2000 (Swann et al., 2002; Averill-Murray and Averill-Murray, 2005) and for *G. agassizii* in the Upper Virgin River Recovery Unit in Utah since a pilot study in 1997 (McLuckie

et al., 2010). The USFWS used line distance sampling to estimate abundance of tortoises in the remaining five recovery units for *G. agassizii* in Utah, Arizona, Nevada, and California starting in 2001 (USFWS 2006, 2009, 2012a, 2012b, 2012c). This report includes results of training exercises for field crews, describes implementation of monitoring, and presents the analysis of desert tortoise density in 2012.

METHODS

Study areas and transect locations

Long-term monitoring strata (Figure 1) will be used over the life of the project to describe population trends in areas managed to conserve tortoises (“tortoise conservation areas,” TCAs). Generally each critical habitat unit (CHU) is treated as one monitoring stratum, although the portion of Mormon Mesa CHU that is associated with Coyote Springs Valley is treated as a separate stratum. Chuckwalla CHU is also treated as dual monitoring strata, with potentially unequal sampling effort in the areas managed by the Department of Defense (Chocolate Mountain Aerial Gunnery Range, CMAGR) and by the Bureau of Land Management (BLM). New recovery units were established under the revised recovery plan (USFWS, 2011), so while making the corresponding changes to our databases we also separated the Piute and Eldorado Valleys into 2 distinct strata; they are in different recovery units. The Joshua Tree stratum does not encompass all suitable habitat for desert tortoises in Joshua Tree National Park (JTNP). The national park designation and current boundaries just post-date the designation of CHUs, so some of the Pinto Mountains and Chuckwalla CHUs (and monitoring strata) are in the current JTNP.

The optimal number of transects in a monitoring stratum was determined by evaluating how these samples would contribute to the precision of the annual density estimate for a given recovery unit (Anderson and Burnham, 1996). Power to detect an increasing population size is a function of 1) the magnitude of the increasing trend, 2) the “background noise” against which the trend operates, and 3) the length of time the trend is followed (even a small annual population increase will result in a noticeably larger population size if the increase continues for many years).

The magnitude of the population trend is a function of recovery activities and the population dynamics of the tortoise – neither of these elements are affected by monitoring design and sample size. The second contributor to the power to detect a trend – the level of background variability in the density estimates – is directly affected by the number, length, and placement of transects in the monitoring strata. Anderson and Burnham (1996) recommended that transect number and length be chosen to target precision reflected in a coefficient of variation (CV) of 10-15% for the estimate of importance (here, density for tortoise conservation areas in each recovery unit). The CV describes the standard deviation (a measure of variability) as a

proportion of the mean and is often converted to a percentage. Since recovery criteria target trends within recovery units (USFWS, 2011a), precision in that density estimate was the focus. The target CV is achieved based on the number of tortoises that might be encountered there (some strata currently have higher densities than others), as well as the area of the stratum – its proportional contribution to the recovery unit density estimate (Buckland et al., 2001).

The actual number of transects assigned in each stratum was a function of the optimal numbers described above, as well as on available funding. Once the number of transects in a stratum was determined, these were laid out systematically across strata, with a random origin for the lattice of transects. In strata with more assigned transects, nested lattices with smaller spacing (3 km) were used to ensure sufficient transects. In strata with fewer transects, lattices 9 km spacing were used. Systematic placement provides more even coverage of the entire stratum, something that may not occur when strictly random placement of transects is used. In both cases, transects are located at random with respect to the location of desert tortoises.

Distance sampling transect completion

One adaptation that tortoises have for living in the desert is to restrict surface activity to fairly narrow windows of time during the year. In general, tortoises emerge from deep within shelters (burrows) from mid-March through mid-May and then again (less predictably) in the fall. These periods coincide with flowering of their preferred food plants (in spring) and with annual mating cycles (in fall). The annual range-wide monitoring effort is scheduled to match the spring activity period for tortoises.

During this season, not all tortoises are above ground or visible in burrows. To encounter as many tortoises as possible, monitoring is scheduled for early in the day and to be completed before the hottest time of day. Because tortoises are located visually, monitoring is restricted to daylight hours. Based on past experience, we expect tortoises to become most active after 7am at the beginning of April (it is usually too cool before this time), but to emerge earlier and earlier until their optimal activity period includes sunrise by the beginning of May. In May, we also expect daytime temperatures to limit tortoise above-ground activity as the morning progresses to afternoon.




Figure 1. Sampled areas 2012.

Field crews complete transects during this optimal period each day. Start times are decided a week in advance, so crews arrive at transects at similar times on a given morning. However, completion times will be more variable, as a consequence of terrain, air temperature, number of tortoises encountered, etc. Under normal conditions, each team walked one 12 km square transect each day. Teams were comprised of 2 field personnel who switched lead and follow positions at each corner of each transect, so they each spent an equal amount of time in the leader and follower positions. The leader walked on the designated compass bearing while pulling a 25 m length of durable cord; the walked path is also the transect centerline and was indicated by the location of the cord. The length of cord also spaced the two independent observers, guiding the path of the follower; when the cord was placed on the ground after a tortoise or carcass was detected, it facilitated measurement of the local transect bearing. The walked length of each transect was calculated as the straight-line distance between GPS point coordinates that were recorded at 500 m intervals (waypoints) along the transect and/or whenever the transect bearing changed.

Both leader and follower scanned for tortoises independently without leaving the centerline, and the role of the crew member finding each tortoise was recorded in the data. Although the leader saw most of the tortoises, the role of the follower was to see any remaining tortoises near the centerline, so the follower role is crucial to unbiased estimation of tortoise densities.

Distance sampling requires that distance from the transect centerline to tortoises is measured accurately. When a tortoise was observed, crews 1) used a compass to determine the local transect bearing based on the orientation of the 25 m centerline, 2) used a compass to determine the bearing from the point of observation to the tortoise, and 3) used a measuring tape to determine the distance from the observer to the tortoise. These data are sufficient to calculate the perpendicular distance from the observed tortoise to the local transect line. If the tortoise was outside of a burrow, was handled enough to measure midline carapace length (MCL), to determine its sex, assess its body condition (USFWS 2011b), and to apply a small numbered tag to one scute. If a tortoise could not be measured because it was in a burrow, because temperatures precluded handling, or for any other reason, crews attempted to establish by other means whether the animal was larger than 180 mm MCL, the criterion for including animals in density estimates.

Because transects are 3 km on one side, it is not unusual for that path to cross through varied terrain or even be blocked by an obstacle such as an interstate highway. In the first years of this program, smaller transects in inconvenient locations were shifted or replaced, but this compromised the representative nature of the sample. Since 2007, the basic rules for modifying transects involve 1) reflecting or elongating transects to avoid obstacles associated with human infrastructure (large roads, private inholdings, etc.), or 2) shortening transects in rugged terrain. Substrate and access to transects can also make it difficult to complete transects during the

optimal period of times, so 3) transects could be shortened to enable completion before 4pm each day.

In 2008 and 2009, the rules for shortening transects were made more restrictive. Crews had the option to complete transects that were 12 km long (in low-relief terrain) or 6 km long (where higher-relief terrain precluded completion of 12 km in a working day). In the latter case, to avoid crews selecting particular terrain, the only way to shorten the transect was to walk it in the southwestern quadrant of the intended 12 km square. If the southwestern quadrant was judged too rugged to be completed safely by transect walkers, the final option was to not complete the transect at all.

In 2010 and 2011, the same option to shorten transects to 6km in rugged terrain were made available to crews in the eastern portion of the range where there was sufficient funding to substratify encounter rates for flat and rugged terrain. In the western part of the range, crews shortened transects by following as much of the planned 12 km route as was possible. This technique was implemented in 2012 for the entire range. If it was anticipated that fewer than 4 km could be walked, the transect should be replaced instead with a transect from the alternate list. I assumed that the proportion of the area that is unwalkable is the same as the proportion of total planned kilometers (12 X number of planned transects) that were unwalkable. As in previous years, unwalked transects were replaced from the list of alternates. More situations were anticipated by additional rules in 2010, as described below.

Transects that crossed stratum boundaries into public lands had previously been walked as planned (squares). Although this added sampling just outside the stratum, it seemed reasonable to assume the land management and tortoise fate would be similar on each side of the invisible boundary. Walking in a square is also less likely to introduce other problems compared to reflecting the transect. Nonetheless, starting in 2010, the protocol used to modify transects that intersected private lands or interstates since 2007 was applied to the portion of any transect that crossed out of monitoring strata, reflecting that portion into the stratum. Whether the segments of those transects outside the boundaries were walked outside the stratum or as a mirror image inside the stratum, the same length of transect was walked at the same distance from the stratum boundary, avoiding undersampling of areas on stratum boundaries (Figure 2). The impetus for this change was the recent large scale development and construction on public lands, often just along the borders of critical habitat, especially for renewable energy facilities and transmission lines.

Specifics of how transect paths were to be modified for rugged terrain (shortened) or for administrative boundaries (reflected) can be found in the *2012 Desert Tortoise Monitoring Handbook* (USFWS 2012d).




Figure 2. Planned (dotted lines) and reflected transect paths at administrative boundaries, now also applied to stratum edges. A) One-corner reflection. B) Two-corner reflection.

Proportion of tortoises available for detection by line distance sampling, G_0

Although we have general expectations about when tortoises are most active each day, and plan our sampling to match the best season and time of day, the fact remains that basing our density estimates only on the tortoises that are visible will result in density estimates that are consistently underestimated (biased low). Instead, we use telemetry to estimate the proportion of tortoises available for sampling, G_0 (“gee-sub-zero”), which is incorporated in the equation for estimating tortoise density and is used to correct this bias.

Telemetry allows us to locate radio-equipped tortoises that are visible as well as those that are otherwise undetectable in deep burrows or well hidden in dense vegetation. To quantify the proportion that were available for detection (“visible”) in 2012, telemetry technicians used a VHF radio receiver and directional antenna to locate 9-19 radio-equipped G_0 tortoises in each of 10 sites throughout the Mojave and Colorado deserts (Fig. 1).

Each time a transmittered tortoise was located, the observer determined whether the tortoise was visible (*yes* or *no*). Through careful coordination, observers at telemetry sites monitored visibility during the same daily time period when field crews were walking transects in the same region of the desert. Observers completed a survey circuit of all focal animals as many times as possible during the allotted time, recording visibility each time. Bootstrapped estimates of G_0 started by selecting one visibility record at random for each tortoise on each day it was located. The average visibility of all tortoise observations at a site on a given day was calculated and used to estimate the mean and variance of G_0 at that site. When there was more than one site in a given area, the G_0 estimate was calculated as the grand mean of all G_0 sites in the group. One thousand bootstrap samples were generated in PASW Statistics (release 18.0.2; SPSS, Inc., 2 April 2010) to estimate G_0 and its standard error.

Use of radio transmitters/receivers to locate tortoises is a technique that is very different from the method used to detect tortoises on line transects. Therefore, in addition to stating whether any part of the tortoise is visible when located, since 2008 behavioral observers and transect walkers have categorized all “visible” tortoises and burrows (when tortoises are found in burrows) as low, medium, or high visibility based on the ability to see part of the tortoise or its burrow from any angle of approach. For the telemetry observers it is a matter of locating a tortoise (visible or not) after they have determined its general location aurally, whereas transect walkers are not searching with certainty of locating a tortoise – they rely only on visual cues. We would therefore not be surprised if the distance sampling method results in detection of a higher proportion of “high” visibility and a lower proportion of “low” visibility tortoises/burrows than when tortoises are located using telemetry. If the odds of being detected differ not only by distance from the line but also a combination of method of detection used (visual or radio receiver) and visibility, we should be able to describe this difference and be able to modify our calculation of visibility following radio-receiver information to more accurately match the visibility to transect walkers.

Field observer training

Training for careful data collection and consistency between crews is fundamental part of quality assurance for this project. This training includes instruction as well as required practice time on skills such as tortoise handling, walking practice transects, and developing detection and distance-measuring techniques. The latter skills include practice on a training course with tortoise models (Table 1). The monitoring handbook developed in 2008 was comprehensive, and serves as a training manual and documentation of training that is provided. Chapters are updated each year as needed and printed for training. They are also posted to the Desert Tortoise Recovery Office website (http://www.fws.gov/nevada/desert_tortoise/reports).

In 2012, three teams of field observers participated. Kiva Biological (Kiva) supplied crews for monitoring in the West Mojave and the eastern portion of the Colorado Desert recovery units. The Institute for Wildlife Studies (IWS) monitored in the western portion of the Colorado Desert and in the Eastern Mojave recovery units. Great Basin Institute (GBI) supplied crews for monitoring in the Northeastern Mojave Recovery Unit. Eight of 10 personnel for Kiva had previous experience with this monitoring program, as did 2 of the 11 personnel for IWS, and 12 of the 26 personnel for GBI were returnees. The three teams were trained in 2 overlapping periods, and to enhance consistency in application of field protocols across teams, where possible the same trainers were used in both training sessions and across teams. Also, for small-group training, experienced personnel from each team worked with the trainees from other teams.

Telemetry training

The primary goal of G_0 training is successful implementation of the G_0 protocol by telemetry crews. This includes correct use of telemetry equipment, understanding G_0 data collection fields, observation of as many radio-equipped tortoises as possible during the day, and covering a window of observation that overlaps the day's transect observation period for each sampling area. Although all telemetry crews had some prior telemetry experience, performance on this project differs from others that do not require confirmation of the exact location of the tortoise. Unless the exact location is determined, its visibility cannot be accurately recorded. Beyond instruction and testing on use of the equipment in desert terrain, several days of practice were compulsory to be able to troubleshoot locating the tortoise and confirming the location when it could not be seen. In addition, some instruction for telemetry and transect crews overlapped to help each group better understand the purpose their data serve and how separate data types are related to the final density estimate.

Distance sampling training

Transect walkers were given classroom instruction, field demonstrations, practice transects to complete, and ultimately each team was evaluated based on performance on a field arena outfitted with a high density of polystyrene tortoise models placed in measured locations (Anderson et al., 2001).

Polystyrene desert tortoise models were set out on the training course each year using placement instructions (vegetation or open placement, distance along training line, and distance perpendicular from training line). This course was used to determine whether 1) individual teams are able to detect all models on the transect centerline, 2) whether their survey techniques yield useful detection functions, and 3) whether they can accurately report the distance of each model from the transect centerline. For each purpose, many opportunities must be provided, so the course is populated at a very high density of models (410/km²).

Crews were sent on transects and training lines as paired, independent observers. That is, the follower was 25 m behind the leader, with the opportunity to detect models not found by the leader. If the leader detects 80% of all tortoises that are found, the assumption is that the follower detects 80% of the tortoises that are missed by the leader. If this assumption is true, in this example, the pair together will detect $0.80 + (0.80 \times (1 - 0.80)) = 0.96$ of all tortoises on the centerline. Because the location of all models was known, data from training lines were also used to 1) assess the dual-observer assumption that all models were equally detectable (detections attributed to the follower occur at the same rate as original detection rate by leader), and 2) to estimate the detection rate using this technique for tortoises elsewhere in the Mojave Desert. These data on models were used to evaluate and correct crew performance before the field season, but were not used in any way to estimate densities of live tortoises once field surveys began.

Table 1. Training schedule for 2012

Day/ Date	Trainees I (IWS, Kiva)			Trainees II (GBI)		
	Activity	Location	Trainer	Activity	Location	Trainer
WEEK 1						
Monday, 12-Mar	Transect methods overview 6 km transect (paper forms)	DTCC	Allison Experienced crews	Same as Trainees I		
Tuesday, 13-Mar	Desert Tortoise Recovery Program Working on Public Lands Distance Sampling Tortoise Activity/G ₀ Transect methods Non-standard transects Juno, Pendragon Database Field crew quality control procedures Compass & GPS for distance sampling	GBI field station	Allison BLM LE Allison “ “ “ Patil “ Allison	Same as Trainees I		
Wednesday, 14-Mar	Tortoise biology and handling 1 Tortoise handling - small groups Compass Exercise Training line preview and crew QA/QC	DTCC " " "	Christopher GBI Staff Crew Allison	Full transects (12km)	LSTS	Allison/ Data spec
Thursday, 15-Mar	Training Lines (practice, 8km) Begin data download from RDAs	DTCC	Allison/AM	Tortoise biology and handling 1 Tortoise handling - small groups Training line preview and crew QAQC Compass Exercise Begin data download from RDAs	DTCC " " "	Woodman, Dutcher GBI Staff Allison/PM Crew
Friday, 16-Mar	Full transects (12km)	LSTS	Allison/ Data specialist	Training Lines (practice, 8km)	DTCC	Allison

DRAFT: Range-wide Monitoring of the Mojave Desert Tortoise: 2012

Day/Date	Trainees I (IWS/Kiva)			Trainees II (GBI)		
	Activity	Location	Trainer	Activity	Location	Trainer
WEEK 2						
Monday, 19-Mar	Tortoise handling 2 Training line debriefing	DTCC “	Veterinarian Allison	G0 / activity observation	River Mtns	Sparks
Tuesday, 20-Mar	Training Lines (evaluation, 8km)	DTCC Mgmt Area	Crew	Tortoise handling 2 Training line debriefing	DTCC “	Veterinarian Allison
Wednesday, 21-Mar	Training Lines (evaluation, 8km)	DTCC Mgmt Area	Crew	Full transects (rugged or administrative)	LSTS	
Thursday, 22-Mar	Full transects (rugged)	LSTS		Training Lines (evaluation, 8km)	DTCC	Crew
Friday, 23-Mar	G0 / activity observation	River Mtns	Sparks	Training Lines (evaluation, 8km)	DTCC	Crew
WEEK 3						
Monday, 26-Mar	Full transects (administrative obstacle)	LSTS		Tortoise handling 3 Training line debriefing	DTCC “	Staff Allison
Tuesday, 27-Mar	Tortoise handling 3 Training line debriefing	DTCC “	Staff Allison	Repeat training lines as needed <i>Begin field data collection</i>		
Wednesday 28-Mar	Repeat training lines as needed <i>Begin field data collection</i>					

DRAFT: Range-wide Monitoring of the Mojave Desert Tortoise: 2012

Day/ Date	Trainees III (experienced Kiva)			QAQC Specialists			G ₀ Trainees		
	Activity	Location	Trainer	Activity	Location	Trainer	Activity	Location	Trainer
WEEK 0									
Wed, 7-Mar							Intro to desert, tortoises, test of telemetry skills		Sparks
Thursday, 8-Mar				Datasheet QA/QC, visibility	GBI field sta.	Allison	G0 datasheets, daily protocols & QA/QC, visibility	GBI field station	Allison
				Walkorder review I		“	Beginning G0 instruction	Piute	Sparks
Friday, 9-Mar	Review new data fields	DTCC	Allison						
	Training line practice	DTCC							
Saturday, 10 Mar	Training lines (evaluation)	DTCC							
Sunday, 11 Mar	Training lines (evaluation)	DTCC							

DRAFT: Range-wide Monitoring of the Mojave Desert Tortoise: 2012

Day/Date	Trainees III (experienced Kiva)	QAQC Specialists	G ₀ Trainees
WEEK 1			
Monday, 12-Mar	[Training for CMAGR]		G0 assessment course River Mtns Sparks / AM
Tuesday, 13 Mar	[CMAGR south]	Desert Tortoise Recovery Program Working on Public Lands Distance Sampling Tortoise Activity/G ₀ Transect methods lecture Non-standard transects RDA/BT GPS, Pendragon Database QA/QC procedures for field crews Compass and GPS for Distance Sampling	Same as Data Specialists GBI field station Allison BLM LE Allison Allison Allison Allison Patil Patil Allison
Wed, 14-Mar	[CMAGR south]	Submit walk order for first 3 weeks	Tortoise biology & handling DTCC Christopher Tortoise handling " GBI Staff Compass Exercise " Crew Training line preview " Allison
Thursday, 15-Mar	[CMAGR south]	Data transfer and QA/QC process Begin data download from RDAs	Training Lines (practice, 8km) DTCC Allison Patil/Brenne-man
Friday, 16-Mar	[CMAGR south]	Initial QAQC by specialists Paper review with data download of LSTS, G ₀	G ₀ - practice 1 + data transfer CS Sparks Brenne-man DTRO
Saturday, 17 Mar	[CMAGR south]	Data upload by specialist Team1	
Sunday, 18 March		Data upload by specialist Team2	

DRAFT: Range-wide Monitoring of the Mojave Desert Tortoise: 2012

Day/Date	Trainees III (experienced Kiva)			QAQC Specialists			G ₀ Trainees		
	Activity	Location	Trainer	Activity	Location	Trainer	Activity	Location	Trainer
WEEK 2									
Monday, 19-Mar	Tortoise handling 2 Training line debriefing	DTCC "	Johnson Allison	Training line data upload			GBI: G0 and activity observation	River Mtns	Sparks
Tuesday, 20 Mar	Full transects (rugged)	LSTS	Crew				IWS: Tortoise handling 2	DTCC	
Wed, 21 Mar	Repeat training lines as needed <i>Field data collection</i>	LSTS		Walk order review	GBI station	Allison	All: G0 - practice 3	IV camp in PI	Sparks
Thursday, 22-Mar							All: G0 - practice 3 continued	CM	Sparks
Friday, 23-Mar							IWS: G0 and activity observation	River Mtns	Sparks
Saturday, 24 Mar				Data upload: transects, training lines, 2 days G0					
WEEK 3									
Monday, 26-Mar							Preliminary GB locations	GB	GBI
							IWS: Full transects	LSTS	
WEEK 4									
Monday, 9-Apr				Deliver QA/QC'd training data					

Data management including quality assurance and quality control

Two sets of data tables are maintained through the field season, organizing data collected on transects and at the G_0 sites. Collection data forms, sheets, applications, and databases are designed to minimize data entry errors and facilitate data verification and validation. Data were collected in both electronic and paper formats by the two survey organizations, then combined and processed in a series of phases to create final database products. Data quality assurance and quality control (data QA/QC, also known as verification and validation) is performed during the data collection, data integration, and data finalization phases. During the second, data integration phase, after combining data from separate groups, some attribute fields are added and all fields are formatted for final processing. The third phase, data finalization, involves consolidation, resolution of data inconsistencies, and generation of final spatial and non-spatial data products used for analysis. After data analysis and reporting are completed, electronic data are actively hosted for download from the internet through http://www.mojavedata.gov/deserttortoise_gov/recovery/data.php. Figure 3 describes the overall data flow.




Figure 3. Data flow from collection through final products.

Tortoise encounter rate and development of detection functions

The number of tortoises seen in each stratum and their distances from the line are used to estimate the encounter rate (tortoises seen per kilometer walked), the detection rate (proportion of available tortoises that are detected out to a certain distance from the transect centerline), and their respective variances. Detection function estimation is “pooling robust” under most conditions (Buckland et al., 2001). This property holds as long as factors that cause variability in the curve shape are represented proportionately (Marques et al., 2007). Factors that can affect curve shape include vegetation that differentially obscures vision with distance, and different detection protocols used by individual crews (pairs). Field teams (IWS, GBI, Kiva) typically walk different number of transects. For this reason, after the field season I expected to develop at least one curve for each field team, which also corresponds to different regions of the desert. The encounter rate is less sensitive to small sample sizes, so it was estimated for each stratum separately.

I used Program DISTANCE, Version 6, Release 2 (Thomas et al., 2010) to fit appropriate detection functions, to estimate the encounter rate of tortoises in each stratum, and to calculate the associated variances. One record was created for each transect, with additional records for each additional tortoise on that transect. Analysis was applied to all live tortoises larger than 180 mm MCL. Transects were packaged into monitoring strata (“regions” in Program DISTANCE).

I truncated observations to improve model fit as judged by the simplicity (reasonableness) of the resulting detection function estimate (Buckland et al., 2001:15-16). Using truncated data, I used the Akaike Information Criterion (AIC) to compare detection-function models (uniform, half normal, and hazard-rate) and key function/series expansions (none, cosine, simple polynomial, hermite polynomial) recommended in Buckland et al. (2001).

Proportion of available tortoises detected on the transect centerline, $g(0)$

Transects were conducted by 2-person crews using the method adopted beginning in 2004 (USFWS, 2006). Transects were walked in a continuous fashion, with the lead crew member walking a straight line on a specified compass bearing, trailing about 25 m of line, and the second crew member following at the end of the line. This technique involves little lateral movement off the transect centerline, where attention is focused. Use of two observers allows estimation of the proportion of tortoises detected on the line; this provides a test of the assumption is that all tortoises on the transect centerline are recorded ($g(0) = 1$). The capture probability (p) for tortoises within increasing distances from the transect centerline was estimated as for a 2-pass removal estimator (White et al., 1982): $p = (\text{lead} - \text{follow}) / \text{lead}$, where lead = the number of tortoises first seen by the observer in the leading position and follow = the number of tortoises seen by the observer in the follower position. The corresponding proportion detected on the line by two observers was estimated by $g = 1 - q^2$, where $q = 1 - p$. Figure 4 graphs the relationship between the single-observer detection rate (p) and the corresponding

dual-observer detection rate ($g(0)$; “gee at zero”). The actual proportion detected can be estimated, but to avoid the necessity of compensating for imperfect detection, during training field crews (pairs) are expected to detect 96% of all models within 1 m of the transect centerline. This corresponds to the leader being responsible for at least 80% of the team’s detections near on the centerline in order to meet this standard (Fig. 4) and is the basis for one of the training metrics (see Table 3).

Few or no tortoises are located exactly on the line, and even examining a small interval (such as 1 m on each side of the transect line) results in few observations to precisely estimate $g(0)$. Instead, my test of the assumption involves examination of the lead and follow proportions starting with counts of tortoises in larger intervals from the line, moving to smaller intervals centered on the transect centerline. As the intervals get smaller the sample sizes also get smaller, but the estimates are more relevant to the area right at the transect centerline. The expectation is that the estimates should converge on $g(0) = 1.0$.

If the test does not indicate that all tortoises were seen on the transect centerline, the variance of p can be estimated as the binomial variance = $q(1 + q)/np$ (White et al., 1982), where n = the estimated number of tortoises within 1 m of the transect centerline, and the variance of $g(0)$ is estimated as twice the variance of p .




Figure 4. Relationship between single-observer detections (by the leader, p) and dual-observer (team) detections, $g(0)$.

Estimates of tortoise density

Each year, the density of tortoises is estimated at the level of the recovery unit. The calculation of these densities starts with estimates of the density of tortoises in each stratum from Program DISTANCE, as well as their variance estimates:

$$D = \frac{n}{2wLP_aG_0g(0)},$$

where L is the total length of kilometers walked in each stratum and w is the distance to which observations are truncated, so $2wL$ is the area searched in each stratum. This is a known quantity (not estimated). P_a is the proportion of desert tortoises detected within w meters of the transect centerline and was estimated using detection curves in Program DISTANCE. The encounter rate (n/L) and its variance were estimated in Program DISTANCE for each stratum. Calculation of D requires estimation of n/L , P_a , G_0 , and $g(0)$. This means that the variance of D depends on the variance of these quantities as well.

For desert tortoise densities, the encounter rate (n/L) is estimated independently for each stratum (“unpooled”), whereas proportion of available tortoises and proportion of available tortoises detected on the transect centerline are estimated jointly for all strata ($g(0)$) or for all strata in the recovery unit (G_0). The detection function, which comes into the above equation as P_a , may be estimated jointly or separately for each team, depending on the number and quality of observations. In 2012, separate detection curves were created for each team (GBI, Kiva, and IWS, pooled across all strata surveyed by that team). A schematic of the process leading to density estimates is given in Figure 5. Contributing estimates in the four left-hand columns are listed with the subsets of the data on which they are based. These estimates combined from left to right to generate stratum and recovery unit density estimates.

Whereas the number of tortoises in the set of strata representing a recovery unit can simply be added together, the variance must be arrived at by accounting for whether this involves pooled or unpooled estimates. As described above, three of the four estimates that contribute to calculating density in a stratum were based on data “pooled” from other strata as well, so when encounter rate and detection probability data from these strata are combined, the correlated nature of the detection probability variances has to be accounted for, for instance. Specifically, the method described in Buckland et al. (2001:89) was used to combine density variances correctly and arrive at the variance (and confidence intervals and CV) for the recovery unit. Pooled and unpooled variance estimates cannot currently be combined as needed in Program DISTANCE, so final construction of density mean and variance estimates from the above components was completed without specialized software.

Tortoise encounter rate	Proportion that are visible, G_0	Detection rate, P_a	Proportion seen on the line, $g(\theta)$	Density	Density	
<i>Stratum or substratified</i>	<i>Neighboring G_0 sites</i>	<i>Data collection group</i>	<i>Overall</i>	<i>Stratum</i>	<i>Recovery unit</i>	
FK	Ord Rodman + Superior Cronese	Kiva	Full set of tortoise observations	FK	Western Mojave	
SC				SC		
OR				OR		
AG	Joshua Tree NP + Chuckwalla			AG	Colorado Deserts	
JT				JT		
PT				PT		
CK				CK		
PV	Piute + Chemehuevi + Ivanpah			PV		
CM				CM		
FE		FE				
IV	Piute + Chemehuevi + Ivanpah	IWS		IV		Eastern Mojave
EV				EV		
GB	Gold Butte			GBI	GB	Northeastern Mojave
BD	Halfway Wash				BD	
MM					MM	
CS			Coyote Springs		CS	

Figure 5. Process for developing density estimates in 2012. For each type of estimate, the full set of data was subdivided appropriately, as indicated by columns.

Estimating the area of each stratum sampled and the number of tortoises in that area

Before the 2008 field season, based on experience in 2007 and visual examination of DEM overlays, all assigned transects were classified as possible for completion as 12k, shortened, or as unwalkable (USFWS 2012a). These classifications before the field season are advisory only, because exact ground conditions, weather, and crew condition all affect the ability to complete a transect. If a non-standard transect (not 12 km square) is walked, crews indicate the obstacles they encountered that forced the change in protocol. In addition to the above named factors, substrate that is very loose on a steep slope or that includes large boulders can make progress so slow or treacherous that crews modify the transect.

Each year, some transects are repeated, providing new information on ground conditions, and new transects are attempted. At the end of each field season, transects that were completed differently from expected are evaluated. At that point, a decision is made whether to reclassify the transect. The classification is used to advise future transect completion, but also to estimate the proportion of each monitoring stratum that is actually represented by the walked transects. Proportions used in this report reflect experience with this set of transects through November

2012, as well as untested transects that were classified using topographic overlays as part of planning for 2013.

Crews completed all transects using the 12 km square path, completing as much of that path as possible. The calculation of unwalkable area in these strata based on the proportion of unwalkable kilometers, not unwalkable transects. Transects completed as 12 km represent the 100% completion option. The total area of the stratum that is walkable is estimated as:

$$\textit{Proportion walkable} = \frac{\textit{transect kilometers walked}}{12 \times \textit{transects completed}}$$

If a given stratum covers 5000 km², but only 90% was walkable and represented by our sampling design, then the density estimate applies to 4500 km², and can be used to generate an estimate for the number of tortoises in those 4500 km². Using these area estimates adds another source of imprecision, so abundance estimates are slightly less precise than the density estimates they derive from. The additional error of this estimate is calculated as the error for a binomial proportion.

Modification of previous procedures

Because tortoises are predictably but not reliably above-ground for parts of the year, there has been continuing interest in use of surveys for sign to estimate number of live animals. The uses and limitations of this approach have been discussed at length, with the performance of live tortoise surveys being preferred. However, on smaller areas of land, few tortoises are expected to be found, so estimates based on live tortoise estimates will be imprecise. Krysik (2002) and Duda et al. (2002) explored the correlation between tortoises and sign using 2 approaches. Although the correlation between sign and live animals was not stable, the interest of land managers in developing better estimates of tortoise numbers in areas smaller than an entire management entity (such as a national park) continue. In 2012, transect walkers counted all burrows that they examined, entering these tallies at the next waypoint. We continued estimating tortoise density using our usual approach, but also provide counts and locations of tortoise burrows so that interested parties can focus on particular areas, interpolating from the critical habitat unit to the smaller parcel based on the relative number of burrows at each scale. This comparison is expected to mirror the relative number of tortoises at each scale, with caveats. Because distance sampling surveyors may not have experience identifying all types and conditions of tortoise shelter sites, tallies were made of any soil or rock openings that was examined for live tortoises and was at least 2 tortoise-lengths deep.

Debriefing to describe strengths and weaknesses of project preparation and execution

At the end of each field season, a debriefing meeting was held to review tasks and responsibilities, strengths and weaknesses of the program, and to plan for the next field season. Because the field teams had disbanded by then, field crew members were surveyed prior to the

end of the field season to nonetheless gather their direct input as we identified training and logistical issues to target for improvement before the next field season. Although issues and/or tasks may be ascribed to individual entities, this meeting is most beneficial in identifying where centralized and/or coordinated response is required to improve the quality of the program.

RESULTS

Field observer training

The smaller Kiva and IWS groups trained alongside one another and mostly separate from GBI, although experienced crews worked between all three teams. Training started on 7 March and continued through 27 March (Table 1). Final tests of field detection abilities occurred toward the end of this period.

Proportion of tortoises detected at distances from the transect centerline

Table 2 reports the proportion of models that were available and were detected by each team at 1-, 2-, and 5-m from the transect centerline. Teams were tested after a trial run on the detection lines or after returning crews walked practice transects to refresh the search pattern (3 of 4 Kiva crews). Detection on the centerline should be 100%, and many crews achieved this. Teams 21 and 43 had perfect detection within a meter of the line during their practice trial, so their performance is not a concern.

Table 3 reports further statistics for each team after collecting data on 16 km on the evaluation lines. Measurement accuracy reported in Table 3 gives the average absolute difference between the expected and measured perpendicular distances from the model to the walked line. All measurements for all models during the 2-day trial are used for this estimate, and capture inaccuracies from 1) using a compass and measuring tape to record distances to the models, plus 2) inaccurately following the trajectory of the transect. The latter source of error does not occur on monitoring transects, because the walked transect is the true transect. On training lines, measurement error increases if crew paths diverge from the measured line that was used to place the models. The “Available Models Detected by Leader” column reports the proportion of all models that were found first by the leader. During training, this number is easily calculated and is used to identify crews in which one of the observers is not finding at least 80% of all detected. With an 80% detection rate for the leader, a 96% detection rate is expected for the team.

Although some individual metrics were below-par (gray cells in Tables 2 and 3), all teams performed well overall and no further changes were made. During training, detection curves were fit to each crew’s set of tortoise model observations. In no case was the best-fitting model a negative exponential one without a “shoulder” describing detections near the centerline. Detection curves reflecting such a model would have led to additional practice. The best-fitting of the 3 remaining basic types of models were then fit to the data to generate density estimates in Table 3. In Figure 6 to Figure 8, all of the crew detection curves for each field team are overlaid.

Crews were not evaluated on their ability to match teammates; such overlays were used to focus field personnel on an additional level of conformity they could work toward. Distance sampling and development of a single detection curve from many observers is robust to the effects of pooling across observations from crews with variable search patterns, when observers contribute proportionally to the overall pattern (Marques et al., 2007).

Table 2. Proportion of tortoise models detected by teams within 1-, 2-, or 5-m of the transect centerline. Values that scored below the target of 0.90 at 1- and 2-m are highlighted.

Team Number	1m	2m	5m
1	1.00	0.96	0.93
2	0.94	0.96	0.87
3	1.00	0.93	0.93
4	1.00	0.97	0.81
21	0.86	0.92	0.90
22	0.92	0.93	0.88
23	0.94	0.93	0.92
24	0.94	0.96	0.91
25	1.00	0.86	0.89
41	0.93	0.93	0.86
42	0.86	0.81	0.87
43	0.88	0.89	0.91
44	0.94	0.96	0.91
45	0.93	0.90	0.90
46	0.93	0.93	0.90
47	0.94	0.85	0.89
48	1.00	1.00	0.91
49	1.00	0.96	0.93
50	1.00	0.96	0.88
GBI	0.94	0.92	0.90
Kiva	0.99	0.96	0.89
IWS	0.93	0.92	0.90
Overall	0.95	0.93	0.89

The Kiva teams had very similar detection patterns by the time the field season began (Fig. 8). Within the GBI crews, teams 44, 48, and 50 had the most anomalous curves (broadest shoulders) in Figure 6. The usual concern when crews are successful searching farther from the line is that they will focus less close to the line; this was not the case for any of these 3 teams. They were coached on tightening their search pattern to better match other teams, but the patterns were not of concern. The extra detections by some IWS team 23 past 10m (the single curve with an extra pair of inflections), are of most concern when this indicates that one of the searchers is not detecting tortoises at the line. However, Team 23 was otherwise judged to have fairly healthy

diagnostics. The testing arena differs from the normal field setting in ways that make it easier to evaluate trainees (higher encounter rate, for example), but also make it more difficult to use as an absolute standard for search patterns.

Table 3. Diagnostics for individual teams after training

Team	Available models detected		Measured v. exact model distance (m)	Estimated abundance	95% confidence interval	
	Within 2m of centerline by leader	Within 2m of centerline by team			Lower limit	Upper limit
1	0.93	0.96	1.23	499	416.2	598.6
2	0.93	0.96	0.81	440	359.4	538.5
3	0.93	0.93	0.72	470	402.4	549.2
4	0.93	0.97	0.78	472	384.8	580.0
21	0.88	0.92	0.86	377	312.4	454.1
22	0.89	0.93	1.08	389	328.9	459.8
23	0.85	0.93	0.89	435	340.7	555.9
24	0.92	0.96	1.60	386	304.7	489.7
25	0.86	0.86	1.19	360	309.2	418.2
41	0.86	0.93	1.12	410	351.5	477.1
42	0.74	0.81	0.77	410	371.5	451.9
43	0.89	0.89	0.90	425	344.7	525.1
44	0.89	0.96	0.83	436	348.2	545.3
45	0.79	0.90	0.78	422	363.0	491.4
46	0.89	0.93	0.82	411	353.5	476.8
47	0.74	0.85	0.83	402	332.8	486.0
48	0.92	1.00	0.72	424	388.7	461.9
49	0.96	0.96	0.76	409	336.4	496.9
50	0.82	0.96	1.17	356	316.1	400.1
Target	>0.80	>0.90	<1	410		
GBI	0.85	0.92	0.87	410.4		
Kiva	0.93	0.96	0.89	470.4		
IWS	0.88	0.92	1.12	389.3		
Overall	0.87	0.93	0.94	417.5		




Figure 6. Detection curves for each of the 2012 GBI teams during training. Curves are based on 16 km trials with approximately 100 detections.




Figure 7. Detection curves for each of the 2012 IWS trainee teams. Curves are based on 16 km trials with approximately 100 detections.




Figure 8. Detection curves for each of the 2012 Kiva trainee teams. Curves are based on 16 km trials with approximately 100 detections.

Quality assurance and quality control

There were 17,654 transect records and 8216 G_0 records associated with the monitoring effort in 2012. After data specialists with the field teams had finished verifying and validating the information in these 2 databases, there were 461 cases where the data were inconsistent with constraints and expectations. (Note that many more issues are addressed each year by data specialists for field crews before the field data are submitted.) Relatively few (378) were errors created by the field crews (sometimes faulty equipment, other times data entry error), of which all were corrected by later phases of QA/QC with recourse to paper datasheets or were identified as correct but unusual data entries (not errors). Another 140 errors were “processing” errors that were identified and corrected before the final database products were created. Processing steps are associated with correcting other errors (perhaps the correct entry is mis-entered), with adding new fields, or any other manipulation that occurs after the data have been collected. When files of scanned datasheets are missing pages, this is a processing error. Other entries violate QA/QC rules but are extreme or explicable entries; there were 301 of these exceptions which require time to research and are one of the costs of QA/QC. Nonetheless, this year of data had relatively few errors, comparable to the previous year of improved QA/QC, and reflecting the improved processes to limit data entry and processing errors.

Transect completion

Table 4 reports the number of assigned and completed transects in each stratum. For the second year in a row, no transects were inadvertently completed twice by different crews. Kiva was assigned 160 transects and walked an additional transect. They consistently replaced any assigned but unwalkable transects with alternates in the same strata. All assigned transects were completed or replaced by IWS, although one unwalkable transect in the Eldorado Valley was replaced by an alternate in neighboring Piute Valley. The Great Basin Institute completed 380 transects, replacing any assigned but unwalkable transects with alternates in the same strata. Great Basin Institute used base-camping in route-less areas to provision crews with supplies, including water, to enable crews to complete 17 transects without returning over large or difficult distances to their vehicles.

Table 4 indicates the number of assigned transects that could be completed as standard square 12km transects or by reflecting around property boundaries and infrastructure (column 4). These transects represent flatter topography. An additional number (column 5) were shortened and represent more rugged terrain. Finally, some transects were considered unwalkable (column 6). Figures 9 through 12 show locations of transects and observations of live tortoises.

Table 4. Number and type of transects in each stratum.

Stratum	Assigned transects	Assigned and alternate transects completed*	Assigned, completed 12k	Assigned, completed shortened	Assigned, judged unwalkable
AG	33	33	20	5	8
BD	70	70	58	11	1
CK	20	20	8	5	7
CM	15	15	12	2	1
CS	89	89	59	17	13
EV	29	29	19	6	4
FE	14	14	14	0	0
FK	16	16	15	0	1
GB	100	100	59	20	21
IV	27	27	23	3	1
JT	18	18	5	7	6
MM	110	106	72	27	7
OR	15	15	7	5	3
PT	15	15	2	7	6
PV	15	15	7	3	5
SC	45	45	37	4	4
Total	631	627	417	122	88
GBI	369	365	248	75	42
IWS	100	100	75	14	11
Kiva	162	162	94	33	35

*Assigned transects that were not walked were supposed to be replaced by alternates.




Figure 9. Distribution of distance sampling transects and live tortoise observations in the Northeastern Mojave Recovery Unit (Coyote Springs Valley, Mormon Mesa, Beaver Dam Slope, and Gold Butte-Pakoon monitoring strata).




Figure 10. Distribution of distance sampling transects and live tortoise observations in the Eastern Mojave Recovery Unit (Eldorado Valley and Ivanpah monitoring strata) and the eastern part of the Colorado Desert Recovery Unit (Fenner, Piute Valley, and Chemehuevi monitoring strata).




Figure 11. Distribution of distance sampling transects and live tortoise observations in the Western Mojave Recovery Unit (Fremont-Kramer, Superior-Cronese, and Ord-Rodman monitoring strata).




Figure 12. Distribution of distance sampling transects and live tortoise observations in the western portion of the Colorado Desert Recovery Unit (Pinto Mountains, Joshua Tree, Chocolate Mountain, and Chuckwalla monitoring strata).

Burrow encounter rate

We reported tallies of examined burrows at each waypoint, representing a path 500 m long. Crews examined 12,817 burrows, inspecting as many as 30 in a 500 m path between waypoints.

Tortoise encounter rates and detection functions

In 2012, all pairs worked together from the beginning to the end of the season. Each Kiva crew walked on average 41 transects and overall they detected 77 tortoises over 180 mm MCL; GBI crews walked a median of 37 transects and detected 162 tortoises; and IWS crews had considerably lower effort, with a median of 22 transects (days of work). However, each IWS observer subsequently walked 1-person distance transects in a separate study at the Large Scale Translocation Site near Jean, Nevada; altogether IWS personnel completed 31 days of work and the team observed 30 tortoises larger than 180 mm MCL, so the LSTS encounters were used to model the overall IWS detection probability illustrated in Fig. 14. Even when crews have similar levels of effort, if there are sufficient observations to generate separate detection curves for each team, that is preferred to also account for differences in vegetation and other regional differences that may alter detection probabilities.

Figures 13 to 15 are histograms of the observed number of tortoises seen at increasing distance from the transect centerline. There is one histogram for each team (Kiva, IWS, and GBI). These observations were used to model detection curves, overlaid in the same figures. Based on detection function behavior, it is typical to discard a few observations in the tails of the histograms in order to build a more robust model (Buckland et al. 2001). Each figure indicates the customized truncation distance that was applied. Any observations that are not used to estimate detection functions will also not be used to estimate the encounter rate (tortoises detected per kilometer walked). In distance sampling applications for many other species, encounter rate can be estimated with relatively high precision, but tortoise encounter rates are low enough that this becomes a factor in considering how to truncate observations to develop detection functions. Truncation was conservative to maximize the number of observations per stratum.

Kiva detected only 77 tortoises from which a detection curve could be built. All truncation distances considered resulted in approximately the same overall density estimate, so it was preferable to use as many observations as possible to keep precision as high as possible. Use of only observations within 14 m allowed a model with no extra inflections and with all but 5% of the observations. At this distance, the hazard-rate model with no adjustments performed best (Fig. 15). The best-fitting negative exponential model was 3rd best overall.

For GBI, truncation at 21 m required discarding just under 7% of the available observations but provided a fairly simple model fit including $g(w) = 10\%$. The 21 m truncation distance was used

(Fig. 13), for which the hazard rate model with 2 parameter estimates but no adjustments fit the data best, and indicate a shoulder of about 2 m – just about 10% of the width.

For IWS, the 5% truncation would be at about 14 m. The model for data truncated at 15 m is a better fit, however, and the hazard rate model with no adjustments performed better than half-normal or uniform models. The shoulder is very narrow, and in fact the best negative exponential model had an AIC slightly less than the selected hazard rate model.

The area below the curves in Figs. 13-15 is the proportion of tortoises that were detected, P_a , estimated as far as the truncation distance (the farthest distance on the x-axis in each figure). Based on these curves, GBI detected 37.9% of the visible tortoises within 21 m of the centerline (CV=0.121). The corresponding estimate of P_a for strata surveyed by IWS was 43.2% (CV=0.263) within 15 m, and for Kiva was 39.7% (CV=0.145) within 14 m.




Figure 13. Observed detections (histogram) and the resulting detection function (smooth curve) for live tortoises with MCL \geq 180mm found by GBI. This curve uses only the 151 observations found within 21 m of the line.




Figure 14. Observed detections (histogram) and the resulting detection function (smooth curve) for live tortoises with $MCL \geq 180$ mm found by IWS. This curve uses only the 28 observations found within 15 m of the line.




Figure 15. Observed detections (histogram) and the resulting detection function (smooth curve) for live tortoises with $MCL \geq 180$ mm found by Kiva. This curve uses only the 73 observations found within 14 m of the line.

Proportion of tortoises available for detection by line distance sampling, G_0

In general, telemetry sites and associated transects were completed sequentially, from south to north. This pattern corresponds to the expected timing of tortoise activity; peaking first in the south, later in the north. In 2012, to accommodate access dates on Chocolate Mountain, this stratum was completed before and during training. The remaining strata in the eastern part of the Colorado Desert were started 12 days after this, but visibility from the Chuckwalla and Joshua Tree telemetry sites were high during the entire period, so a single G_0 estimate was used for all of these strata and for the entire 4 week period. Tortoise activity in the eastern part of the range was very delayed compared to typical patterns, and this is reflected in unusually low G_0 estimates for Gold Butte, Halfway Wash, Chemehuevi, Piute-Mid, and Ivanpah, despite the fact that these strata were monitored in April and May. Dates, total days monitored, and G_0 estimates are given in Table 5.

Strata in the western part of the range were apparently sampled during a very active period, reflected in high G_0 estimates. In the eastern part of the range, G_0 remained relatively lower, only increasing slightly in Coyote Springs Valley which was monitored later in the field season.

Table 5. Availability of tortoises (G_0) during the period in 2012 when transects were walked in each group of neighboring strata.

G_0 sites	Strata	Dates	Days	G_0 (Std Error)
Gold-Butte	Gold Butte	27 Mar – 6 Apr	11	0.52 (0.119)
Halfway Wash	Beaver Dam Slope, Mormon Mesa	23 Apr – 21 May	29	0.54 (0.116)
Coyote Springs Valley	Coyote Springs	9 Apr – 22 Apr	14	0.98 (0.049)
Piute/ Ivanpah/ Chemehuevi	Chemehuevi, Eldorado, Fenner, Ivanpah, Piute	28 Mar – 3 May	37	0.52 (0.163)
Joshua Tree/ Chuckwalla	Chocolate Mtn, Chuckwalla, Joshua Tree, Pinto Mtns	13 Mar – 9Apr	28	0.86 (0.107)
Ord-Rodman/ Superior Cronese	Fremont-Kramer, Ord- Rodman, Superior Cronese	10 Apr – 29 Apr	20	0.87 (0.138)

Proportion of available tortoises detected on the transect centerline, $g(0)$

Because they are cryptic, even tortoises that are visible (not covered by dense vegetation or out of sight in a burrow) may not be detected. For 25 detections of tortoises within 1 m of the transect centerline, 21 were found by the observer in the lead position and 4 by the follower, so that the probability of detection by single observer, $p = 0.810$, and the proportion detected using the dual observer method, $g(0 \text{ to } 1 \text{ m}) = 0.964$ (SE = 0.11). Figure 16 shows that $g(0)$ was converging on 1.0, indicating the assumption of perfect detection on the centerline was met; consequently, no adjustment was made to the final density estimate. The curves since dual

observers were first used in 2004 have all supported the premise that complete detection on the transect line was achieved for years in which the dual-observer method was used (USFWS 2009, 2012a, 2012b).




Figure 16. Detection pattern for the leader (p) and by the team ($g(0)$) based on all observations out to a given distance (x) from the centerline in 2012. Note convergence of $g(0)$ on 1.0 as x goes to 0.

Estimates of tortoise density

Density estimates were generated separately for each monitoring stratum (Table 6), then weighted by stratum area to arrive at average density in the monitored area of each recovery unit (Table 7). Although encounter rates were estimated separately for each stratum, and have independent variances, the detection function and G_0 were estimated jointly (pooling data from multiple strata), so these variances are not independent (Fig. 5 illustrates how estimates were pooled for 2012).

When the annual estimates are imprecise, it should not be expected that there will be a close match from one year to the next. Over a period of many years, however, we expect any underlying trend in the number of tortoises to be obvious through this “background noise.” Some density estimates in 2012 were based on so few observations that the estimates and very high CVs are beyond the expected 20-40% CV. This arose from last-minute funding withdrawals that manifested as sequestration rather than strategic alteration of the sampling design.

Table 6. Recovery unit and stratum-level encounters and densities in 2012 for tortoises with MCL \geq 180mm.

Recovery Unit	Stratum	Area (km ²)	Number of Transects	Total transect length (km)	Sampling Dates		Field Observers	<i>n</i> (tortoises observed)	<i>CV</i> (<i>n</i>)	<i>Density</i> (/km ²)	<i>CV</i> (<i>Density</i>)
					Begin	End					
Colorado Desert		13530	130	1403	30-Mar	24-Mar		44		2.4	26.0
Chocolate Mountain	AG	755	33	363	13-Mar	24-Mar	Kiva	21	24.7	6.1	31.26
Chuckwalla	CK	3509	20	213	14-Apr	22-Apr	Kiva	8	37.5	3.9	42.10
Chemehuevi	CM	4038	15	176	30-Mar	28-Apr	IWS	1	99.9	0.8	107.85
Fenner	FE	1841	14	168	30-Mar	27-Apr	IWS	1	100.0	0.9	107.95
Joshua Tree	JT	1567	18	183	14-Apr	22-Apr	Kiva	6	44.5	3.4	48.45
Pinto Mountains	PT	751	15	140	14-Apr	22-Apr	Kiva	5	38.5	3.7	42.97
Piute Valley	PV	1070	15	159	30-Mar	28-Apr	IWS	2	67.4	1.8	78.76
Eastern Mojave		3720	56	639	30-Mar	28-Apr		8		2.2	60.5
Eldorado Valley	EV	1153	29	320	30-Mar	28-Apr	IWS	2	69.1	0.9	80.19
Ivanpah	IV	2567	27	318	30-Mar	28-Apr	IWS	6	49.9	2.8	64.41
Northeastern Mojave		4889	369	4184	6-Apr	27-May		164		3.4	20.1
Beaver Dam Slope	BD	828	70	819	20-Apr	10-May	GBI	38	17.0	5.4	29.93
Coyote Springs Valley	CS	1117	89	996	13-May	27-May	GBI	45	18.3	2.9	22.55
Gold Butte-Pakoon	GB	1977	100	1116	6-Apr	20-Apr	GBI	21	20.5	2.3	32.89
Mormon Mesa	MM	968	110	1253	20-Apr	13-May	GBI	47	17.2	4.3	30.03
Western Mojave		6873	76	876	25-Mar	11-Apr		33		3.6	28.8
Fremont-Kramer	FK	2417	16	193	25-Mar	9-Apr	Kiva	4	44.8	2.2	49.68
Ord-Rodman	OR	1124	15	158	27-Mar	10-Apr	Kiva	7	41.2	4.6	46.54
Superior-Cronese	SC	3332	45	525	25-Mar	11-Apr	Kiva	22	24.3	4.4	32.49



Table 7. Estimated density of desert tortoises in monitored areas of each recovery unit in the Mojave and Colorado deserts in 2012.

Recovery Unit	Sampled area (km ²)	Kilometers walked	Tortoises detected	Density (/km ²)	Lower limit 95% CI (Density)	Upper limit 95% CI (Density)	%CV (Density)
Colorado Desert	13530	1403	44	2.4	1.45	3.95	26.0
Eastern Mojave	3720	639	8	2.2	0.75	6.67	60.5
Northeastern Mojave	4889	4184	164	3.4	2.29	5.00	20.1
Western Mojave	6873	876	33	3.6	2.05	6.21	28.8

Area of each stratum sampled and the number of tortoises in that area

Evaluating transect classification

In 2012, 149 of the 627 walked and 88 unwalked transects were not completed or addressed as predicted. Table 8 summarizes conclusions after examining these transects. Sixty were reclassified based on crew experience. In some cases, this reflects discrepancy between interpretation of terrain from imagery; in others, classification is ambiguous because over the course of a 12 km transect, terrain is so variable to that it was not a simple matter to evaluate the ability of a typical crew to complete it. The remaining 89 anomalous transects were not reclassified, because earlier experience indicated that most crews would use the original completion strategy. The 60 transects that were reclassified represent only 2.1% of the 2786 transects evaluated since 2008, so this had very little impact on our estimate of the proportion of each stratum that is walkable.

Table 8. Transects completed other than as planned and any resulting reclassification

Previous substratum	Situation	New substratum	Number of transects
12k	Shortened in 2012, contradicting previous experience. In many cases, shortened on military base for non-terrain issues	12k	30
12k	On-the-ground observation differs from imagery	Shortened	25
12k	Crew attempted the transect but couldn't walk at least 6km	Unwalkable	1
Shortened	On-the-ground experience indicates walkable though rugged	12k	21
Shortened	Lengthened or replaced as unwalkable in 2012, but not enough evidence to override previous crew experience	Shortened	59
Shortened	Crew planners considered unwalkable; reexamination of topo maps matches their interpretation	Unwalkable	6
Unwalkable	On-the-ground experience contradicted imagery	12k	1
Unwalkable	On-the-ground experience contradicted imagery	Shortened	6

Number of tortoises based on density and area sampled

The proportion of each stratum represented by distance sampling is calculated as the proportion of planned kilometers that can be walked (Column 3 in Table 9). This proportion is calculated based on all transects evaluated since 2008. Table 9 reports the area of each stratum, the proportion covered by our density estimates, and the associated estimate of tortoise abundance.

Table 9. Estimated tortoise abundance in sampled areas of each stratum

Stratum	Area (km ²)	Proportion sampled	SE(Prop. Sampled)	Sampled area	N (number of tortoises)	95% Confidence Interval	
						Lower Limit	Upper Limit
AG	755	0.94	0.027	706	4287	2350.3	7818.4
BD	828	0.91	0.029	756	4059	2279.1	7230.2
CK	3509	0.80	0.031	2818	11123	5024.3	24623.4
CM	4038	0.93	0.023	3763	3142	561.5	17582.5
CS	1117	0.86	0.031	960	2789	1793.3	4339.1
EV	1153	0.87	0.030	999	916	230.0	3649.6
FE	1841	0.97	0.017	1782	1561	278.8	8744.8
FK	2417	0.97	0.010	2347	5060	2015.2	12707.6
GB	1977	0.82	0.026	1623	3675	1955.5	6906.4
IV	2567	0.95	0.019	2447	6777	2134.8	21512.8
JT	1567	0.74	0.034	1152	3962	1605.0	9780.4
MM	968	0.87	0.033	844	3661	2049.1	6539.8
OR	1124	0.73	0.040	821	3768	1573.5	9023.3
PT	751	0.68	0.052	508	1900	838.1	4308.0
PV	1070	0.87	0.031	927	1714	438.7	6696.5
SC	3332	0.93	0.014	3088	13432	7215.1	25004.1
Total	29012	0.880		25541	71827	46684.8	110508.5

Possibility of using burrow counts to estimate tortoise abundance at smaller scales

Crews examined 12,817 burrows as they collected tortoise observations to arrive at this estimate of N. Because there was different effort (kilometers walked) in each stratum, Figure 17 plots the relationship between cover sites and tortoise abundance as their standardized measures (burrows per kilometer walked *versus* tortoise density). Although some association is apparent it is not a linear one; without the cluster of poorly sampled strata in the lower left of Fig. 17, no association is apparent at all. This is not entirely surprising, although it is worth exploring whether the association is accurate at smaller spatial scales. In the Colorado Desert, Krysik (2002) has shown that a strong association exists for 9 km² plots in a square study area that is 7 km on a side, but is broken when the plots are 64 km apart. Meanwhile, in areas where burrows are in caliche, not dirt, the mechanism for any association (e.g., more desert tortoises construct and maintain more burrows) does not exist, so for either small or large areas with predominantly caliche caves, the associations are not expected to hold up. This is clearly a topic that should be addressed with additional data and at different scales.




Figure 17. Scatter plot of stratum densities and burrow encounter rate.

Debriefing to identify strengths and weaknesses in preparation for future years

This meeting was held on 31 May, about 1 week after the last field work was completed. Crews were generally well prepared and appropriately addressed any complications in transect and telemetry work. The following issues were identified to be addressed by coordinated effort rather than by efforts of single parties.

Earlier review of the first field data

This was a consistent complaint of crews – that they did not get it in a timely manner, at all, or it did not address their work. There were issues with specialists and USFWS schedules that created unique timing problems this year, but in any year it would be helpful to circulate copies of the assessments during training so crews know ahead of time the type of review they will see.

Training improvements to make more effective use of same time period

In general, crews felt they received the appropriate amount of preparation (list of goals) for each training day and their work was reviewed effectively. However, 2 or 3 days of practice transects were completed at the LSTS and crews felt the transects should have been reviewed as if it were the regular field season.

To avoid conflicting advice from experienced personnel attempting to fill in gaps in the written information, it was suggested that a FAQ and/or step-through list be created for 2013 G₀. For instance, the book and training should be clear that seeing more tortoises is a priority over having more observations of a small number of tortoises. What should be reported when the tortoise's behavior was apparently in response to seeing the technician first? How often should recently transmittered tortoises be tracked and how should this be documented in our G₀ database so that these animals are not included in analyses?

Photos taken on transects should be incorporated in training to illustrate terrain and burrow types not encountered during the pre-season.

DISCUSSION

Quality control and quality assurance

Electronic data collection devices were purchased before the 2011 field season, reducing equipment-related errors, and these were still performing as well or better in 2012 due to greater care with batteries. In addition, the project continued to rely on extensive and intensive feedback to crews and to data specialists during training and the field season. The training emphasis was on review of paper datasheets and standardization of rules for modifying records and documenting these changes. Data managers involved in the final stages of data processing were used to train crews involved in data collection. When project rely on large numbers of relatively independent data collection crews, this sort of standardization reduces identifiable and unidentifiable errors in the final database. The 2012 database products were comparable to those in 2011, with a marked decrease in the number of errors passed through from the field teams to independent reviewers.

Improving ability to detect trends in desert tortoise abundance

The primary goal of the monitoring program is to provide population estimates that are relevant to the recovery plan criteria (USFWS, 2011). The priority for this and every field season is therefore to improve ability to detect trends in desert tortoise abundance at the recovery unit level.

Consequences of insufficient transects

Estimates of tortoise visibility (G_0) were relatively low in the eastern part of the range; associated with this, encounter rates were the lowest that have been reported on this project (under 4 per 100 km walked in all recovery units), but adequate funding in some portions of the range allowed for sufficient transects to nonetheless detect several tortoises per stratum. In 10 of the 16 surveyed strata, fewer than 10 tortoises were detected; this is a rough lower limit for the number of observations to develop an adequate density estimate (Buckland et al. 2001). None of

the stratum estimates are meant to stand on their own, and will either be used in combination with other stratum estimates in the same recovery unit (in annual reports such as this one) or with other years of estimates from the same stratum to describe trends; however, the dramatic loss of funding for transects BLM lands in California resulted in such reduced effort and detections that these stratum estimates should probably not be used in any aggregated for either. One way to better utilize available funds is to refrain from sampling in areas without at least a minimum level of funding. Instead of using annual recovery unit estimates to develop trends, description of recovery unit trends will be based on average trends within strata in that recovery unit. Missing years of information should be avoided if it is an important goal to describe annual recovery unit abundance, requiring representation of all associated strata. However, infrequent years of missing data do not prevent trends in individual strata from being described.

Although Anderson and Burnham (1996) targeted CVs between 10 and 15% in each recovery unit, this would be unusually high precision for such an estimate of wildlife abundance. In 2012, 3 recovery units had CVs between 20 and 29% of the density estimate – satisfactory for an annual estimate. The Eastern Mojave Recovery Unit had a density estimate of much lower precision (60.5 %). This is not surprising given the lack of investment in these strata, but is of less concern than the small number of tortoise observations ($n=8$) on which the density estimate for these 3700 km² is based.

LITERATURE CITED

- Anderson, D.R., and K.P. Burnham. 1996. A monitoring program for the desert tortoise. Report to the Desert Tortoise Management Oversight Group.
- Anderson, D.R., K.P. Burnham, B.C. Lubow, L. Thomas, P.S. Corn, P.A. Medica, and R.W. Marlow. 2001. Field trials of line transect methods applied to estimation of desert tortoise abundance. *Journal of Wildlife Management* 65:583-597.
- Averill-Murray, R.C., and A. Averill-Murray. 2005. Regional-scale estimation of density and habitat use of the desert tortoise (*Gopherus agassizii*) in Arizona. *Journal of Herpetology* 39:65–72.
- Berry, K.H., and L.L. Nicholson. 1984. The distribution and density of desert tortoise populations in California in the 1970's. Chapter 2 in K.H. Berry (ed.), *The status of the desert tortoise (Gopherus agassizii) in the United States*. Desert Tortoise Council Report to the U.S. Fish and Wildlife Service. Order No. 11310-0083-81.
- Buckland, S.T., D.R. Anderson, K.P. Burnham, J.L. Laake, D.L. Borchers, and L. Thomas. 2001. *Introduction to Distance Sampling: Estimating Abundance of Biological Populations*. Oxford Univ. Press, Oxford. 432 pp.
- Corn, P. S. 1994. Recent trends of desert tortoise populations in the Mojave Desert. *Fish and Wildlife Research* 13:85-93.
- Duda, J.J., Krzysik, A.J. and J.M. Meloche. 2002. Spatial organization of desert tortoises and their burrows at a landscape scale. *Chelonian Conservation and biology* 4:387-397.
- Krzysik, A.J. 2002. A landscape sampling protocol for estimating distribution and density patterns of desert tortoises at multiple spatial scales. *Chelonian Conservation and biology* 4:366-379.
- Luckenbach, R.A. 1982. Ecology and management of the desert tortoise (*Gopherus agassizii*) in California. In R.B. Bury (ed.). *North American Tortoises: Conservation and Ecology*. U.S. Fish and Wildlife Service, Wildlife Research Report 12, Washington, D.C.
- Marques, T.A., L. Thomas, S.G. Fancy, and S. T. Buckland. 2007. Improving estimates of bird density using multiple-covariate distance sampling. *The Auk* 124(4) 1229-1243.

- McLuckie, A.M., M. Ratchford, and R.A. Fridell. 2012. Regional desert tortoise monitoring in the Red Cliffs Desert Reserve, 2011. Salt Lake City: Utah Division of Wildlife Resources, Publication Number 12-13. 54 pp.
- Murphy, RW, KH Berry, T Edwards, AE Leviton, A Lathrop, JD Riedle, 2011. The dazed and confused identity of Agassiz's land tortoise, *Gopherus agassizii* (Testudines, Testudinidae) with the description of a new species, and its consequences for conservation. *ZooKeys* 113: 33-71. doi: 10.3897/zookeys.113.1353.
- SPSS, Inc. 2010. PASW Statistics, Rel. 18.0.2. Chicago: SPSS Inc.
- Swann, D.E., R.C. Averill-Murray, and C.R. Schwalbe. 2002. Distance sampling for Sonoran desert tortoises. *Journal of Wildlife Management* 66:969–975.
- Thomas, L, S.T. Buckland, E.A. Rexstad, J.L. Laake, S. Strindberg, S.L. Hedley, J.R.B. Bishop, T.A. Marques, and K.P. Burnham. 2010. Distance software: design and analysis of distance sampling surveys for estimating population size. *Journal of Applied Ecology* 47:5-14.
- Tracy, C.R., R.C. Averill-Murray, W.I. Boarman, D. Delehanty, J.S. Heaton, E.D. McCoy, D.J. Morafka, K.E. Nussear, B.E. Hagerty, and P.A. Medica. 2004. Desert Tortoise Recovery Plan Assessment. Report to the U.S. Fish and Wildlife Service, Reno, Nevada.
- [USFWS] U.S. Fish and Wildlife Service. 1994. Desert Tortoise (Mojave Population) Recovery Plan. U.S. Fish and Wildlife Service, Portland, Oregon.
- [USFWS] U.S. Fish and Wildlife Service. 2006. Range-wide Monitoring of the Mojave Population of the Desert Tortoise: 2001-2005 Summary Report. Report by the Desert Tortoise Recovery Office, U.S. Fish and Wildlife Service, Reno, Nevada.
- [USFWS] U.S. Fish and Wildlife Service. 2009. Range-wide Monitoring of the Mojave Population of the Desert Tortoise: 2007 Annual Report. Report by the Desert Tortoise Recovery Office, U.S. Fish and Wildlife Service, Reno, Nevada.
- [USFWS] U.S. Fish and Wildlife Service. 2011a. Revised recovery plan for the Mojave Population of the desert tortoise (*Gopherus agassizii*). U.S. Fish and Wildlife Service, Pacific Southwest Region, Sacramento, California. 222 pp.
- [USFWS] U.S. Fish and Wildlife Service. 2011b. Health Assessment Procedures for the Desert Tortoise (*Gopherus agassizii*): A Handbook Pertinent to Translocation. Desert Tortoise

Recovery Office, U.S. Fish and Wildlife Service, Reno, Nevada. Accessible through:
http://www.fws.gov/nevada/desert_tortoise/index.html

[USFWS] U.S. Fish and Wildlife Service. 2012a. Range-wide Monitoring of the Mojave Desert Tortoise: 2008 and 2009 Reporting. Report by the Desert Tortoise Recovery Office, U.S. Fish and Wildlife Service, Reno, Nevada.

[USFWS] U.S. Fish and Wildlife Service. 2012b. Range-wide Monitoring of the Mojave Desert Tortoise: 2010 Reporting. Report by the Desert Tortoise Recovery Office, U.S. Fish and Wildlife Service, Reno, Nevada.

[USFWS] U.S. Fish and Wildlife Service. 2012c. DRAFT Range-wide Monitoring of the Mojave Desert Tortoise: 2011 Reporting. Report by the Desert Tortoise Recovery Office, U.S. Fish and Wildlife Service, Reno, Nevada.

[USFWS] U.S. Fish and Wildlife Service. 2012d. Desert Tortoise Monitoring Handbook. Desert Tortoise Recovery Office, U.S. Fish and Wildlife Service, Reno, Nevada. Version: 6 March 2012. http://www.fws.gov/nevada/desert_tortoise/reports.

White, G.C., D.R. Anderson, K.P. Burnham, and D.L. Otis. 1982. Capture-recapture and removal methods for sampling closed populations. LA-87-87-NERP. Los Alamos National Laboratory, Los Alamos, NM. 235pp.