



# Paleoseismic features as indicators of earthquake hazards in North Coastal, San Diego County, California, USA

Gerald G. Kuhn

1031 Palm Avenue, #26, Carlsbad, CA 92008, USA

Received 11 October 2004; received in revised form 8 February 2005; accepted 8 April 2005  
Available online 4 June 2005

**DOCKET**

**11-IEP-1J**

DATE

RECD. AUG 02 2011

## Abstract

New road cut and mass-grading excavations in the north coastal area of San Diego County, California expose heretofore generally unrecognized, probable late Holocene tsunami deposits and paleoseismically deformed sediments. Remnant tsunami deposits occur up to 100 + m in elevation around the margins of modern coastal lagoons and estuaries and, combined with local mima mounds of possible sand blow origin, provide indirect but compelling evidence for the late Quaternary activity of onshore and offshore faults in the immediate study area. Probable paleoliquefaction features are regionally widespread and range from fissures filled with sediments derived from overlying marine terrace sand and soil, to the more traditional sand-filled injection dikes and sills, lateral spreads, and filled craterlets. The source of most liquefied sediment is underlying Tertiary “bedrock sand” and local, Quaternary marine-terrace deposits. A paleoseismic liquefaction origin rather than soft-sediment loading is deduced for these features based on morphology, internal stratigraphy, field setting, and near proximity to known seismogenic sources.

Some paleoseismic events impacted late Holocene Indian middens and burial sites. The last seismic event probably occurred within the past 1 to 3 ka, and possibly even records the historic earthquakes of either November 22, 1800 or May 27, 1862. The liquefaction features also affect marine terrace sediments tens of meters above modern regional water levels, inferentially “recording” paleoseismic events in this Mediterranean-type climatic region during winter rains when high-level, perched water saturates the several meter thick source sediments.

Based on their regional extent, the paleoseismic features were likely caused by  $M \sim 7+$  tectonic events inferentially generated by the nearby offshore Newport–Inglewood/Rose Canyon fault system, or possibly by smaller, recently exposed, related and localized faults. Accordingly, the seismic hazard of the north coastal area of San Diego County may be substantially higher than previously assumed, and hence of concern owing to the rapid ongoing and projected population increase.

© 2005 Elsevier B.V. All rights reserved.

**Keywords:** Earthquake indicators; Tsunami deposits; Holocene; California; Paleoseismic events

## 1. Introduction

The population of southern California is now approximately 20 million and increasing. Much

E-mail address: [gkuhn@nethere.com](mailto:gkuhn@nethere.com).




Fig. 1. Map showing location of paleoseismic features along the north San Diego County coast. Los Angeles (LA), San Diego (SD) and the study area (hatched) are shown for reference.

new growth is taking place along the coastal area of northern San Diego County (Fig. 1). Here, from the shoreline to about 18 km inland, thousands of new houses and commercial centers have been, and are being, constructed. Combined with related road and infrastructure excavation, the new exposures now reveal hundreds of heretofore unrecognized paleotsunami and seismically-induced liquefaction features. Detailed mapping, both in vertical cuts and in continuous, grading-excavations, shows that these features are regionally widespread and recurrent. Accordingly, they provide stratigraphic evidence of late Quaternary, prehistoric earthquakes and possible harbingers of future earthquakes and risk in this part of California. The north coastal area of San Diego County has a typical Mediterranean climate: average annual precipitation is approximately 250 mm; mostly occurring during the winter months of November through March. Much natural vegetation has long been removed, and is

now supplanted by exotic introduced species such as eucalyptus, palms, and a host of other decorative plants associated with rapid, post-WW II urban spread.

The presence of people and active faults often form a deadly combination, well demonstrated in California by recent, high-magnitude earthquakes and related damage in the 1971 San Fernando earthquake (Oakshott, 1975), and the 1994 Northridge earthquake (Woods and Seiple, 1995). Neotectonic investigations in California traditionally focus on on-shore surface rupture associated with geomorphically well defined fault systems as documented by many workers for the San Andreas and its various splays (Table 1). New investigations now show that many seismic sources are “so-called” blind faults, which can generate high-magnitude earthquakes and yet have only subtle or broadly distributed surface geomorphic expression (Namson and Davis, 1988; Davis et al., 1989; Shaw and Shearer, 2000). Similar blind thrusts and other seismic sources occur immediately offshore the southern California coast as deduced from interpretation of geophysical data (Bohannon and Geist, 1998; Rivero et al., 2000; Figs. 1 and 2; Table 2). Because of their offshore location, the earthquake history of these faults is poorly known; however, it now appears that at least some are recorded by on-shore tsunami deposits and related paleoliquefaction features.

Table 1  
Local and regional onshore geology

Blake, 1856a,b; Goodyear, 1888; Fairbanks, 1893; Ellis and Lee, 1919; Hertlein and Grant, 1944, 1954; Larson, 1948; Emery, 1950a; Wilson, 1972; Hannan, 1973; Moyle, 1973; Barrows, 1974; Crowell, 1974; Kennedy, 1975; Kennedy et al., 1975; Shepard and Kuhn, 1977; Sieh, 1978; Guptill and Heath, 1981; Emery and Kuhn, 1982; Weber, 1982; Hall, 1984; Kuhn and Shepard, 1984; Eisenberg, 1985; Weldon and Sieh, 1985; Tan, 1986; Harden and Matti, 1989; U.S. Geological Survey [USGS], 1990a; Lajoie et al., 1991; Prentice and Schwartz, 1991; Aydin et al., 1992; Kern and Rockwell, 1992; USGS, 1992, 1998; USGS and Southern California Earthquake Center [SCEC], 1994; Lindvall and Rockwell, 1995; Sims and Garvin, 1995; Tan and Kennedy, 1996; Grant et al., 1999, 2002; Vaughan et al., 1999; Baldwin et al., 2000; Franklin and Kuhn, 2000; Kuhn, 2000; Kuhn et al., 2000, 2004; Lienkaemper, 2001; Grant and Rockwell, 2002; Stone et al., 2002.




Fig. 2. Major fault zones in the immediate offshore and onshore area of the southern California coast from Los Angeles on the north to near Ensenada in Baja, California on the south. Modified from Legg (1985).

The north coastal paleoliquefaction features are particularly well preserved on flights of marine terraces and within their underlying regressive sands (Figs. 3 and 4). These have been mapped, albeit discontinuously, from approximately San Diego on the south to north of Los Angeles (Fig. 2). Locally, the wave-cut platforms and overlying marine deposits are capped by beach ridges, former dune fields now somewhat lithified. Recent mass-grading (excavations for new houses and commercial development) now exposes hundreds of “anomalous” sedimentary features, which are rea-

sonably judged to be of paleoseismic origin. Accordingly, this paper initially sets forth the geomorphic and neotectonic setting of the study area. This is followed by description of the inferred paleoliquefaction features and the reasons why they are not a result of local loading and soft-sediment deformation. Also discussed is indirect evidence supporting the paleoseismic hypothesis; namely, the inferred seismic origin of local mima mounds, and the character and presence of tsunami deposits. This is followed by interpretation of data concerning paleoseismic recurrence and magni-

Table 2

## Offshore geology

Shepard and Emery, 1941; Shepard, 1948; Emery, 1960; Hand and Emery, 1963; Moore, 1969; Moore, 1972; Western Geophysical, 1972; Legg and Kennedy, 1979; 1991; Field and Richmond, 1980; Kennedy et al., 1980a,b, 1985, 1987; Graham and Bachman, 1983; Clarke et al., 1985, 1987, Legg, 1985, 1987, 1991; Darigo and Osborne, 1986; Greene and Kennedy, 1987; Fischer and Mills, 1991, Fischer et al., 1992; Legg et al., 1992, 1994, 2003, 2004; Crouch and Suppe, 1993; Astiz and Shearer, 2000; Rivero et al., 2000; Sliter et al., 2001; Legg and Kamerling, 2003.

pointed to and support the paleoseismic hypothesis for the now-documented sand blows, sand dikes, craterlets, sand laccoliths, filled fissures (that is, fissures filled with materials from overlying sediments), and lateral spreads. These reports and other documents bearing upon neotectonics and earthquake hazards in the area are cited in specific categories pertaining to local onshore and offshore geology, liquefaction, paleoseismites, tsunamigenic, and engineering implications (Tables 1–7).

tude, seismic source area, and earthquake-hazard impact in this part of California.

This paper also summarizes local technical reports and other papers of limited distribution that initially

## 2. Geomorphic and tectonic setting

The geomorphic and tectonic setting of the north coastal, San Diego County area has been described




Fig. 3. Marine terraces in the study area. Terrace 1 is topographically lowest and youngest; terrace 4 is highest and oldest. Cross-section A-A' shown on Fig. 4 (after Tan and Kennedy, 1996; topographic base map from U. S. Geological Survey, 1901).






Fig. 4. Schematic cross-section near Encinitas, California (Fig. 3) showing major marine terraces and wave-cut platforms ( $Qt_1$  through  $Qt_4$ ). Approximate age of terrace sediments deduced from association with the marine-isotope stage chronology (Shackleton and Opdike, 1976), substage 5a through stage 11. Basal regressive marine sand and cobbles deposited denoted by “dots”; fault zone (FZ); terrace thickness exaggerated.

and refined over the past 100 years (Table 1). In addition to published papers and informal guidebook articles, there is a plethora of pertinent unpublished reports and data compilations, many pertaining to neotectonism and earthquake hazards in the area (Table 6). Some data were originally proprietary, obtained for the mid-1970s geologic investigations at and near the San Onofre Nuclear Generating Station in the northwest corner of the Camp Pendleton Marine

Corps Base (Fig. 1). Many associated reports dealt with the origin, relative age and deformation history of coastal marine terraces (Table 6), the same as those in the San Diego County north-coastal area that harbor the postulated paleoseismic features described herein (Table 1).

In 1991, LaJoie et al. summarized marine terrace evolution for this part of California, and dated several erosion platforms using amino-acid and uranium-series techniques. Tan and Kennedy (1996) compiled prior geological mapping for the study area and applied local names for the lower four marine terraces that range in elevation from about 3 to 130 m. Based on association with the marine, oxygen-isotope stage chronology, on local radiocarbon and uranium-series dating, and on relative soil (pedogenic) profile development, the terraces are designated and inferentially dated as:  $Qt_1$  (lowest), ~80–125 ka (marine, oxygen-isotope substages 5a through 5e, respectively);  $Qt_2$ , ~200 ka (stage 7);  $Qt_3$ , ~320 ka (stage 9), and  $Qt_4$ , ~440 ka (stage 11; Figs. 3 and 4).

Table 3  
Liquefaction issues

Atwater et al., 2001; Obermeier and Dickenson, 2000. [Regarding U. S. Northwest Cascadian Earthquake of 1700 A.D.]
Fuller, 1912; Morse, 1941; Saucier, 1987; 1991a,b; Obermeier, 1987, 1989, 1996a,b; USGS, 1990b; Obermeier et al., 1991, 1993, 2004; Marple and Schweig, 1992; Su and Follmer, 1992; Munson et al., 1995; Li et al., 1996; Tuttle et al., 1996; Obermeier and Pond, 1999; Cox et al., 2001; Hough, 2001. [Regarding Midwestern U. S. Earthquakes.]
Dutton, 1889; Obermeier et al., 1985, 1990, 2002; Talwani and Cox, 1985; Peters and Hermann, 1986; Amick and Gelinis, 1991; Marple and Talwani, 2000. [Regarding Charleston, South Carolina Earthquakes.]
Lawson, 1908; Holzer et al., 1989; 2004; Holzer and Clark, 1993; Meisling, 1979; Jachens et al., 2002; Michael et al., 2002; Rymer et al., 2002a,b; Sylvester et al., 2002. [Regarding California Earthquakes.]
Harp et al., 2003. [Regarding Denali, Alaska Earthquake of 2002.]
Rajendran and Rajendran, 2003. [Regarding Kachchh Region, India Earthquakes.]

Table 4  
Paleoseismites

Lamont, 1936; Galli and Ferrel, 1995; Landuzzi et al., 1995; Lucci, 1995; Michetti et al., 1995; Reiter, 1995; Bartholomew et al., 2002; Etensohn et al., 2002; Greb and Dever, 2002; Mariotti et al., 2002; Merriam and Forster, 2002; Moretti et al., 2002; Obermeier et al., 2002; Stewart et al., 2002; Wheeler, 2002.
--

Table 5  
Tsunamis

Emery, 1950b; Shepard et al., 1950; Natland and Kuenen, 1951; Reimnitz and Marshall, 1965; Coleman, 1968; Eiby, 1982; Foster et al., 1991; Shi et al., 1993; Daag et al., 1995; Dawson, 1994, 1995, 1999; Minoura et al., 1994; Bondevik et al., 1995; Bourgeois, 1995; Carver and Gilpin, 1995; Kelsey et al., 1995; Moore and Lewis, 1995; Nishimura and Miyaji, 1995; Peterson and Priest, 1995; Tuttle et al., 1995; Dawson et al., 1996; Morner, 1996; Borrero et al., 1997; Imamura et al., 1997; Synolakis et al., 1997; Ortiz et al., 1998; Clague et al., 1999; Hindson and Andrade, 1999; Kawata et al., 1999; Takashimizu and Masuda, 2000; Atwater et al., 2001; Carey et al., 2001; Zitellini et al., 2001; Okal et al., 2002a,b; Pratt, 2002; Legg et al., 2003, 2004; Scheffers and Kelletat, 2003; Brookfield, 2004; Costa, 2004.

The marine abrasion platforms are typically sequentially overlain by ~1-m-thick veneer of cobbles (former beach gravel), several meters of regressive marine sand, and locally by prograding continental deposits. Locally, the terrace surfaces are covered by “beach ridges,” former coastal dunes now topographically rounded and often relatively lithified. These beach ridge dune sands and the underlying marine sands are prone to seismically-induced liquefaction when saturated, this typically occurring during winter rains when local perched water levels are common (Kuhn et al., 2000, 2004).

From a tectonic standpoint, several prominent aerial-photographic lineaments trend across the study area (Kuhn et al., 2004). These lineaments form a general rhombohedral pattern, with prominent axes oriented northwest and northeast, respectively (Fig. 5). Some lineaments coincide with NW-trending valleys that define the back edge of the terraces; these are now demonstrably fault related as exposed in road and grading cuts and locally in the sea cliffs (Seitz, 1983; Shepard and Kuhn, 1977; Kuhn and Shepard, 1984). The lineaments and related fault patterns are now identified as possible major ancient tectonic shear zones (Slosson et al., 2000), that are probably re-activated. Additionally, new offshore,

Table 6  
Unpublished reports

Euge et al., 1972; Western Geophysical, 1972; Fugro, 1975a,b, 1977a,b; Anderson et al., 1977; Ehlig, 1977; Shlemon, 1977, 1978a,b, 1979a,b,c,d.

Table 7  
Engineering-geology publications

Housner, 1958; Seed and Lee, 1966; Lee and Seed, 1967; Seed, 1968; Scott and Zuckerman, 1973; Youd, 1973, 1984, 1985; Lowe, 1975, 1976; Youd et al., 1978; Seed and Idress, 1983; National Research Council, 1985; Owen, 1987; Holzer et al., 1989; Meier, 1993; Nichols, 1995.

seismic reflection profiles document the nearby presence and transtensional and transpressional character of the Newport–Inglewood/Rose Canyon fault (NIRC), a major component of the San Andreas fault system (Fig. 2). The new urban exposures also show that many onshore lineaments are aligned with concentrations of sand dikes and boils, filled fissures and lateral spreads, features herein interpreted to be of paleoliquefaction origin.

### 3. Description of interpreted paleoliquefaction features

In this section, North San Diego County’s liquefaction-induced features are discussed according to their morphology and genesis. The most widespread and common are dikes, sills, and laccoliths. These intrusions occur on marine terraces  $Qt_1$  through  $Qt_4$ . Craterlets are common, but principally occur only on two terraces. Likewise abundant are lateral spreads and filled fissures. One terrace has a very irregular and undulating surface topography caused by liquefaction at depth (Fig. 5).

#### 3.1. Sand dikes, sills, and laccoliths

##### 3.1.1. Field observations

Railroad and urban-excavation cuts now expose many dikes filled with fine sand that pierce terrace sediments (Figs. 6 and 7), and such terrace sediments range in age from 100 to over 300 ka. The dikes vary from about a few mm to ~10-cm wide, and often taper upward and locally form cross-cutting patterns (Fig. 8). A trench across prominent aerial lineaments on the lowest coastal terrace (Fig. 5) exposed unmatched soil and stratigraphic units, suggesting that this particular lineament is fault controlled (Franklin and Kuhn, 2000; Fig. 9). The trench also exposed paleoliquefaction features such



Fig. 5. 1953 vertical aerial view of the coastal lagoons and terraces at Carlsbad, California. The circular features located in the center of the photograph are “mima mounds;” many are now confirmed to be “sand blow” deposits. Thus several lineaments include lateral spread surfaces or fissure fills (photograph from U.S. Department of Agriculture Photograph #AXN-8M-100, taken on 11 April, 1953).

as sand dikes, lateral spreads, and a sand laccolith. Locally, the source sands, as exposed in the trench, are traced to the underlying Tertiary marine sand (Fig. 10).

Of particular relevance to origin is the relationship of the sand laccolith and the overlying topography (Fig. 11). Here the sand laccolith is demonstrably fed by several sand dikes, which in turn had locally pierced and or otherwise given rise to an irregular, micro-topography. The overlying surface bears a soil likely no more than about 2 to 3 ka old profile and perhaps much younger (Shlomon, 1999, pers. com.) based on the relative degree of development. The liquefied sediments exposed in this trench vary in age from late Holocene to prob-

ably late Pleistocene, and demonstrate that two and perhaps several seismic events occurred in this area. The Tertiary source sands here are still relatively cemented and therefore prone to liquefy under seismic loading (Fig. 10). Terrace sediments containing the dikes vary widely in origin, ranging from Pleistocene (?) regressive marine and clay-rich marsh and lagoon deposits to prograding continental and eolian deposits.

Other trenches on the same marine terrace exposed more sand sills, dikes, laccoliths, and warped argillic soil horizons (Fig. 12), as well as highly contorted lateral spreads that flowed over deformed, tilted sand-flat and marsh sediments (Fig. 13). Also observed were liquefied and offset channel-lag deposits in






Fig. 6. View (1994) of a railroad cut exposure showing fine grain, tan and white sand dikes cutting into and through continental deposits and into an overlying Bt argillic soil horizon at Carlsbad, California (15.24 m el. MSL).

which the adjacent horizontal bedding had been uplifted and eroded (Fig. 14).

Likewise evident are irregularly shaped clay beds with entrained sand, showing that the sand had moved laterally and upward. Implicitly the contorted clay and

sand probably moved plastically to nearby swales, ultimately giving rise to the undulating, modern ground topography.

On adjacent terraces, urban grading cuts exposed polygonal, reticulated, and sand-filled features (Fig.




Fig. 7. View (1994) of sand dikes cutting vertically through clay-rich estuarine deposits at Carlsbad, California (13.41 m el. MSL). Note: Numerous multi-colored “source sand sills” are confined by clay layers, forming sand dikes that fine upward.






Fig. 8. A vertical cut exposing sand-dikes cutting coastal terrace sands at Carlsbad, California (46.63 m el. MSL). Note: The scale is resting on a lateral spread surface which has cutoff the underlying sand dikes, which are injected from below.

15). Where exposed vertically the sand fillings proved to be tabular dikes, which often trended along pre-existing tectonically-related stress fractures. The dikes are filled with upward-fining, loosely-packed, clean

and cohesionless sands. And here, too, some sands are traced to their source in their underlying Tertiary marine beds. Also, as exposed in the various cuts, the dikes are horizontally sheared by lateral spreads (Fig. 8); and where coalescing, the sand dikes are injected into overlying coastal terrace sediments and often form sand laccoliths (Fig. 12).

Terrace  $Qt_2$  exposures show the local presence of gravel-bearing sands at, and near, the tops of vents (Fig. 16). These gravels are similarly traceable into the underlying source sediments, where they have been dragged upward and truncated by lateral spreads. The gravel at the top of the dike may well stem from gravel entrainment during eruption of the liquefied sediment, a phenomenon resulting from strong seismic shaking (Prentice et al., 1992; Meier, 1993; Yegian et al., 1994).

### 3.1.2. Source mechanism

The geometry and character of the sand-filled dikes observed in the study area is very similar to those reported to have originated as a result of seismic liquefaction as documented by Obermeier (1996b). Also, north San Diego County dikes are associated with sand-filled laccoliths, similar to the seismic-induced features of Obermeier (1996a). Accordingly, the sand dikes, sills, and laccoliths are most likely, heretofore, unrecognized paleoseismic features.




Fig. 9. 1994 color aerial photograph taken along the lowest coastal terrace at Carlsbad, California. Note: The strong, dark lineament coincides with depressions.




Fig. 10. Trench cut into a coastal terrace at Carlsbad, California exposing liquefied sediments (light layer) overlying the Eocene-age Santiago Formation (B member) with numerous sand dikes emanating from below.

### 3.2. Craterlets

Obermeier (1996a) pointed out that liquefaction-induced craterlets were extensively produced during the 1886 Charleston, South Carolina earthquake. Similar swarms of craterlets occur in the study area, particularly in terrace  $Qt_4$  (Figs. 3 and 17) likewise supporting a probable seismic origin.

In plan view, the filled craterlets are nearly circular throughout their height and vary in diameter from

about 0.3 to 2.0 m (Fig. 18). Their heights vary from about 2 to 10 m. At depth, their internal stratigraphy is characterized by vertical zones of upward-fining, clean, fine- to medium-grained sands, locally varying in color and texture (Fig. 19). These sands, as recorded during field observations, are injected through each other and locally into and through a central core. The upper part of most craterlets appears to be a collapse feature, creating depressions 0.5 to 1.3 m deep (Fig. 20). These craterlets are confined to an




Fig. 11. Closeup view of warped Holocene colluvial sediments and sand dikes exposed in a trench cut into a coastal terrace at Carlsbad, California. Note: Location is the same as Fig. 9.






Fig. 12. Exposure of a storm-drain cut into the lowest coastal terrace (15.54 m el. MSL) at Carlsbad, California. Note: Multi-colored sand dikes are traceable from marsh source sands (located at base of brush), are upward-fining, and cut through, offset, and warped “beta” (argillic) lamellae.

ancient beach ridge, where the dike sand demonstrably originated as fluidized sediment transported upward from underlying Tertiary marine sands. Locally, the internal stratigraphy of the craterlets shows that they were likely produced during two or more discrete episodes (Fig. 18).

The craterlets also occur in sediments filling linear swales along the back edge of terrace escarp-

ments and local beach ridges (Figs. 3, 4, and 17). Here the craterlets are filled by fluidized sediment, which later was replaced almost entirely by silica. These features occur near the base of the wave-cut platform and grade into a strongly cemented, 1- to 2-m-thick silcrete duripan. Based on local geomorphic and stratigraphic relations, it is deduced that the craterlets similarly formed by seismic liquefaction,




Fig. 13. A construction trench in the lowest coastal terrace (15.84 m el. MSL) at Carlsbad, California exposing tilted paleosand flat sands and marsh sediments, overlain by remnant v-shaped wedges of “beta” (argillic) lamellae, and capped by a clay-rich lateral spread, which flowed easterly, forming a depression. Note: These same sediments, as exposed in sea bluff exposures located 100 m to the west, dip to the west or are near-horizontal.






Fig. 14. A construction trench exposing marsh and sand flat sediments vertically cut by sand dikes at Carlsbad, California (19.50 m el. MSL). The liquefied, dragged, and offset channel-lag deposit and that the adjacent bedding has been eroded.

comparable to that reported by Twidale (1976), in Australia.

Elsewhere many craterlets in late Quaternary sediments are characterized by N30W to N70W filled

fissures. These craterlets were later offset by the filled fissure deposits which, themselves, trend from N–S to approximately E–W. These particular craterlets vary in diameter from about 1.2 to 1.5 m, have a central



Fig. 15. A horizontally-graded terrace surface at Carlsbad, California (46.32 m el. MSL) exposing polygonal-shaped, “reticulated” multi-colored sand-filled features. Note: Vertical cuts of these same polygonal features expose numerous, adjacent, coalescing, upward-fining sand dikes demonstrably injected from below and not the result of surface infill.




Fig. 16. Near-vertical exposure in a coastal terrace made for a major highway cut at Carlsbad, California (33.52–35.05 m el. MSL). Note: The gravel to boulder-bearing sands, deposited in a very fine, well-sorted dark mineral-rich matrix, are observed at the top of vents, and traceable to sand sills and dikes emanating from underlying marine source sands.




Fig. 17. Schematic vertical section of ancient barrier bar (beach ridge) showing sediment types, filled craterlets, and approximate abrasion platform contacts (former shorelines) at Encinitas, California (see Figs. 3 and 4). Adapted from Obermeier (1996b), his Fig. 13, p. 20.

core, and are formed during at least two intrusive episodes (Fig. 18).

### 3.2.1. Source mechanism

The craterlets in the study area are likewise interpreted as of paleoseismic liquefaction in origin, based mainly on the similarity of their internal stratigraphy, their morphology, and their occurrence in tectonic regimes where such craterlets have been previously described. For example, Dutton (1889), noted swarms of similar appearing craterlets, and deduced them to be liquefaction features associated with the 1886

Charleston, South Carolina earthquake of  $M \sim 7.2$ . More recently, similar liquefaction-induced craterlets were regenerated in the Kachchh Region of India following the  $M \sim 7.7$  earthquake in 2001 (Rajendran and Rajendran, 2003). A likely earthquake origin for the study area craterlets is also indicated by the sand-filled tabular fissures whose overall dimensions and shapes suggest that they are “incipient craterlets” (features described by Obermeier, 1996b).

Although yet uncertain, the source of the strong seismic shaking may well be “hidden” or “blind” faults directly beneath the site, locally enhanced by topo-




Fig. 18. Near-circular paleoliquefaction features (craterlets) exposed on the surface during construction grading on a coastal terrace at Encinitas, California (110.94 m el. MSL). Note: The boundary is outlined by scraping tools and scales.



graphic amplification, a phenomenon observed during the 1971  $M \sim 6.6$  San Fernando, California earthquake (Saul, 1975, p. 68–69), and during the 1886  $M \sim 7.2$  earthquake at Charleston, South Carolina (Obermeier, 1996a, p. 345).

Also suggestive of local faulting is the apparent displacement of terrace  $Qt_4$  (Fig. 17), which, to the east, dramatically increases in dip (Fig. 4). Exposures here show that  $Qt_4$  is apparently offset by a fault zone 3–4 m wide. Local faulting and regional tilting in this area has been postulated by others, although few urban and related grading cut exposures were then available (Wilson, 1972; Hannan, 1973; Eisenberg, 1985; Lajoie et al., 1991; Tan and Kennedy, 1996).

Conceivably, although unlikely, the craterlets may owe their origin to some unique, local artesian condition. This postulate, however, is rejected as a source mechanism because the study area craterlets are regionally extensive, and it occurs up-slope from dune field swales where local perched water would most likely occur. Additionally, the craterlets lack a regional hydrogeologic recharge/source area to supply the necessary head for their sole occurrence.




Fig. 19. Paleoliquefaction feature exposed in a vertical cut during construction grading on a coastal terrace at Encinitas, California (102.41 m el. MSL). Note: The multi-colored “flowering upward” feature, cuts through marine terrace sands and underlies the “craterlet” seen in Fig. 18. The blue hue on the right was made by the scraper blade on the terrace wall.




Fig. 20. Paleoliquefaction features exposed in a vertical escarpment along a coastal terrace at Encinitas, California (118.87 m el. MSL). Several episodes of upward-fining, multi-colored, tonal and textured sands are postulated to have been injected through each other, into and through a central core, and exhibit a “collapsed” infilled cone at the top.

### 3.3. Lateral spreads

#### 3.3.1. Field observations

The myriad of new road cuts and grading pads on terraces  $Qt_1$  through  $Qt_4$  expose many dikes and other ground failure features typically associated with seismically-induced lateral spreads, features well described by Seed (1968) and Youd (1984), and depicted graphically by Obermeier (1996a, Fig. 7, part 3).

The new cuts now provide cross-section plan views, thereby permitting above-average field documentation of lateral spread extent and morphology.

The lateral spreads are exemplified by large blocks that are bordered by dikes, which are nearly linear in plan view. The dike sidewalls and widths range from about ~5 to about 45 cm; many have a left or right-step, en-echelon plan-view pattern. In vertical cuts, horizontal ground shifting is apparent by the truncation and general displacement of sand dikes and filled fissures (Fig. 15).

The lateral spreads occur on  $1^\circ$ – $8^\circ$  slopes, although locally slopes may exceed  $10^\circ$  forming minor “landslide” topography (Fig. 13).






Fig. 21. Paleoliquefaction features (fissure fill and intruded sands) exposed by construction grading on coastal terrace at Carlsbad, California (16.45 m el. MSL). Note: The fissure fill, measuring 3.81–5.08 cm wide, consists of down-dropped colluvium in terrace sands, is near-vertical to vertical below the surface, and becomes wedge-shaped with depth.

### 3.3.2. Source mechanism

According to Youd (1984), lateral spreads can take place on slopes as low as  $0.1^\circ$  in the form of laterally moving landslides of non-liquefied sediments moving

atop a liquefied layer. Such lateral spreads may also form grabens at their headwall and thrusting shear zones at the toe (Seed, 1968; Obermeier, 1987). The study area features are morphologically similar to those described by Seed, Youd, and Obermeier and, because they are in direct association with tabular sand dikes, sand laccoliths, and filled fissures, they are therefore interpreted as similarly being of seismic liquefaction origin (Fig. 13).

### 3.4. Filled fissures

#### 3.4.1. Field observations

Hundreds of filled fissures are exposed in new excavations on marine terraces. These fissures are vertical to near-vertical fractures that widen near the ground surface. Most pinch out at depth. They range in height and width from about 5 to 35 cm and from 4 to 8 cm, respectively. Sediments filling the fissures have two sources; some demonstrably emanated from the ground surface and apparently moved downward through gravitational flow (Fig. 21); but other fissures are clearly filled with sand emplaced by fluidization of underlying Tertiary sediments. In sectional view many fissures are tabular and have laminations that parallel fissure walls. Also, many filled fissures cross-cut one another and thus may be highly contorted (Fig. 22).




Fig. 22. Paleoliquefied features (fissure fills and lateral spreads) exposed on a steeply dipping cut slope on a coastal terrace at Encinitas, California (71.32 m el. MSL). Note: Northwest, northeast, and east–west fissure fill sediments offset, drag, and even reverse themselves in a strike-slip mode. Also note that a laterally-spread liquefied layer at the top has cut off fissure fills.




Fig. 23. Exposure displaying 3 epochs of Quaternary terrace fissure fills (from 3.81 to 5.08 cm wide), down-dropped into the underlying “parent” Tertiary sandstone at Carlsbad, California (6.09–7.62 m el. MSL).

The filled fissures occur on a wide age range of sediments; and locally some appear to penetrate into the underlying Tertiary parent sandstone (Fig. 23), and even form wedge-shaped down-dropped blocks (Fig. 24).

#### 3.4.2. Source mechanism

Based on their widespread occurrence, on the multiple sources of fill, and on their association with lateral spreads, the study-area, filled fissures probably formed recurrently.

Locally some fissures may be associated with landslides, themselves possibly induced by seismic events. Indeed, it is also plausible that some filled fissures reflect faulting at depth, for the surface fissure pattern mimics bedrock fractures. Regional faulting is largely strike-slip (Sylvester, 1988; Weldon et al., 1996), and this style of deformation typically gives rise to reversal of throw, transtensional and transpressional fractures and thus to widespread complex ground fissuring (Obermeier et al., 2004).

#### 4. Basis for a liquefaction origin

A proposed seismic liquefaction origin for the filled fissures is similarly based on comparing the morphology of these features with those described elsewhere that are associated with documented seismicity (Tables 1–3). Specifically, the filled fissures in the study area are very similar to those described by Sims and Garvin (1995) who analyzed and described liquefaction features associated with the M~7.1, 1989 Loma Prieta, California earthquake.

Alternative mechanisms for origin of the filled fissures range from local artesian flow to wave-cutting along paleo-shorelines. But these hypotheses are untenable in the study area for artesian flow is not, and has not occurred given the local geomorphic, hydrogeologic, and stratigraphic setting, and wave-cutting is similarly ruled out owing to relative “recency” of liquefaction in marine terrace sediments now tens of meters above present sea level. Of particular interest, however, is the nearby presence of the active Newport–Inglewood/Rose Canyon fault zone, and a related blind thrust immediately offshore (Fig. 2). According to Rivero et al. (2000) and Legg (personal communication, 2004) the blind thrust extends directly under the study area, and may be capable of generating seismic events of M~7.0 or greater (Kuhn et al., 2000, 2004).




Fig. 24. Colluvial wedges; inferred to be large fissure fills exposed in Tertiary sandstone (fractured grey-white unit) at Carlsbad, California (6.09–7.62 m el. MSL).

## 5. Indirect paleoseismic evidence

Mima mound topography also characterizes much of the North San Diego County study area. Although these features alone do not provide incontrovertible evidence for local-area paleoseismicity, their presence compliments and otherwise supports major seismic events in the late Quaternary. Extensive mound fields were once in evidence (Orcutt, 1887) and visible on aerial photography taken along the North San Diego County coastal surfaces, prior to urbanization (USGS, 1947; USDA, 1953). Scattered mounds still occur in Carlsbad and Encinitas and on the Camp Pendleton Marine Base. As shown on aerial photographs and originally noted in soil surveys (USDA, 1918, 1973), continental and estuarine deposits located on coastal terraces  $Qt_1$  at Carlsbad are mantled by a local micro-relief or hummocky landscape of “mima mounds,” each of which is usually less than ~1 m high and 2 to 3 m in diameter (Fig. 5).

### 5.1. Field observations

Site-specific excavation of Carlsbad and Encinitas coastal mound sites show that some mound fields are aligned along fissure fills and laterally spread depressions called “vernal pools”. The mound topography and adjacent vernal pools are irregular. They appear concentrated along some lineaments, possibly fault or fracture zones, a phenomena reported elsewhere in California (Shlemon et al., 1973). The mounds are typically subrounded, are frequently confined by an expansive clay cap, and are often filled by clean white to tan sand, giving rise to a dramatic contrast in color and grain-size, when compared with the generally dark-colored clayey sediments characteristic of surrounding continental or estuarine deposits. As exposed in grading cuts, some mounds directly coincide with sand dikes and sand laccoliths. These sand dikes, which are tabular and tubular in plan view, are traceable to underlying sand source beds derived from Tertiary bedrock. Most terraces are burrowed, and the sand dikes, sills and source sands are often occupied by ground squirrels and gophers that indicate an ongoing biogenic maintenance by rodents. Similar-appearing mound complexes in the San Clemente area, about 48 km to the north, have been interpreted to be mainly constructional (bioturbation), in

origin (Shlemon et al., 1997), however a paleoseismic origin was not precluded.

### 5.2. Multiple origin of mounds

There are many hypotheses for the origin of mounds: eolian deposits (Barnes, 1879); water-deposited features (Dietz, 1945); water and glacial combined (Newcomb, 1952; Washburn, 1997); physical or chemical segregation (Ritchie, 1952); features created by human agency (Aten, 1981); ants, ground squirrels or pocket gophers (Dahlquist and Scheffer, 1942; Arkley and Brown, 1954; Cox, 1990, 1991); seismogenic (Berg, 1990; Riefner and Pryor, 1996), and “multiple origins” (Krnitzsky, 1949).

#### 5.2.1. Source mechanism

Some of the Carlsbad and Encinitas mounds evidently originated as sand blows, and were then later colonized by fossorial rodents. Also some the mounds apparently formed during genesis of sand laccoliths, which hydraulically deformed the overlying clay expansive cap where fluidized sand could not escape to the ground surface. Many of the associated vernal pools were in part created by lateral spreads which are linear in plan view (Kuhn et al., 2000).

The Carlsbad and Encinitas mima-mound topography, is therefore a probable important indicator of paleoseismicity because morphologically similar features are associated with tabular sand dikes, sand laccoliths, sand blow deposits, lateral spreads, filled fissures and plausible tsunami features (Kuhn et al., 1995a,b,c, 2000, 2004). This is contrary to observations from the extensively studied New Madrid Seismic Zone where seismicity apparently did not produce mima mounds (Saucier, 1991a). It should be noted that from aerial photos, the New Madrid sand blows are similar in form to mounds; however, agricultural leveling for almost 200 years has destroyed much of their relief.

## 6. Age of liquefaction features

### 6.1. Evidence for quaternary paleoseismic events

#### 6.1.1. Field observations

Features interpreted to be of seismic liquefaction origin in the Carlsbad to Encinitas area include



extensive swarms of craterlets, principally found on the ~440 ka terrace  $Qt_4$  (Fig. 4). Moreover, fissure fills, sand dikes, lateral spreads and mounds occur on all terraces, which vary in age from ~80 ka to 440 ka. A “classic” exposure at one location in Carlsbad shows that 11 epochs of paleoliquefied sand dikes, fissure fills, and lateral spreads probably occurred during the past 120 ka (Figs. 25 and 26).

## 6.2. Evidence for Holocene paleoseismic events

### 6.2.1. Field observations

The north coastal area of San Diego County contains widespread Paleo-Indian sites, some of

which are radiocarbon-dated at more than 8 ka (Breschini et al., 1992). These archaeological sites consist of burials, “kitchen middens,” and transitory camps that contain diverse artifacts dated mainly by shells and charcoal (Carter, 1957; Gallegos, 1987, 2002). Several early-to-late Holocene archaeological sites have been affected by paleoseismic features. For example, liquefied sand dikes and sills extend upward, intrude and deform midden deposits located on the  $Qt_1$ – $Qt_4$  terraces in the Carlsbad area (Fig. 3). Further, based on radiocarbon-dating, at least one major event took place about 6 to 8 ka bp (Smith, 1996). Some of these liquefaction-deformed archaeological sites occur on uplifted 60-m high, marine terrace deposits, well above Holocene regional water levels (Shlemon and Kuhn, 1997; Fig. 27). It is thus inferred that liquefaction likely took place during the winter rainy season when the high-level terrace sands contained perched water (Kuhn et al., 2000).

Other archaeological evidence suggests that seismically-induced liquefaction may have taken place as recently as 2 to 3 ka ago (Franklin and Kuhn, 2000). Indeed, an archaeological site at Batiquitos lagoon in Carlsbad (Gallegos and Associates, 1997) exposed an intact prehistoric hearth. One test pit in the center of the site (feature 97-1) exposed artifacts and shells that were apparently offset and locally dragged upward by fissure fill sediments (Fig. 28). The deformed artifact horizons yield radiocarbon dates of 0.9 to 1.3 ka, documenting incontrovertible evidence for paleoseismic liquefaction and ground deformations in late Holocene time (Fig. 29).




Fig. 25. Photograph of a vertical cut exposing sand dikes cutting coastal terrace sands at Carlsbad, California (47.24–51.81 m el. MSL). Note: The scale is resting on a lateral spread surface that has cut-off the underlying sand dikes, which were injected from below. At least 10 episodes of lateral spreads cut-off sand dikes. Measured sections of multiple lateral spread surfaces in vertical cuts are shown in Fig. 26.

## 7. Tsunamigenic features

The possible tsunamigenic features identified in this investigation are based on comparison with stratigraphic relations and internal characteristics of tsunami deposits recorded elsewhere (Table 5).

### 7.1. Field observations

Many abrupt, chaotic, convulsive sedimentary features now exposed in the north coastal area of San Diego County cannot be simply explained by non-seismic soft-sediment deformation, nor by variations




Fig. 26. Measured sections of vertical cuts made 15 m apart showing multiple lateral spread surfaces. Same location as seen in Fig. 25.

and cyclicity of storm deposits (“tempestites,” of Aigner, 1985). Rather, the features are similar to those formed by (1) ancient earthquakes [“seismites” (Seilacker, 1969, 1984)]; 2), turbidity current deposits [“turbidites” (SEPM, 1951; Kuenen, 1947, 1957, p. 231)], and; 3) modern tsunami deposits [“tsunamites” (Pratt, 2002)]. Many coastal features also show unusual chaotic variations here likewise inferred to be caused by paleoliquefaction. Typically, a locally generated tsunami deposit can also be deformed by large aftershocks.

The convulsive features in the north coastal Carlsbad–Encinitas region have characteristics that are in common with other tsunamis described elsewhere (Table 5). Specifically:

- (1) The presence of chaotic, turbid, sedimentary debris (Fig. 30), commonly mixed with wood

fragments, shells, clay balls and pods of sharp, angular pieces of terrace deposits and bedrock, and locally capped by peat (Fig. 31).

- (2) The initial event deposits grade from coarse to landward-fining, rather than landward-coarsening, typical of “tempestites” (storm deposits).
- (3) The pebbles are imbricated (Fig. 32).
- (4) The landward movement of sediment is “lobe form” (Fig. 33), similar to those in the Philippines Islands produced by the 1994 Mindoro earthquake (Daag et al., 1995).
- (5) The “v-shaped” chaotic features were inferentially caused by an abrupt “hydraulic jump” when tsunami waves reach a “bedrock high” or very resistant surface feature (Fig. 34), a phenomenon often associated with fluid-like features exposed in the overlying sediments (Morner, 1996; Fig. 35), and;




Fig. 27. Paleoliquefaction features exposed by grading deform, cut, and offset a shell (*Argopecton* sp., *Chione* sp.), charcoal, and artifact-rich, Indian midden site located on a 60 m high, now well-drained terrace at Carlsbad, California. Note: Arrows point to sand dikes, fissure fills (trending N20W and N80W), and lateral spreads that crosscut the terrace sands and offset each other.

- (6) The abrupt landward termination of sand wedges has an angle similar to, but steeper than dunes (Daag et al., 1995; Fig. 36).

Because of their similar internal stratigraphy and occurrence on former abrasion platforms on modern “bedrock highs,” the anomalous, chaotic, con-




Fig. 28. An archaeological site located along the north side of Batiquitos Lagoon at Carlsbad, California (5.18–5.48 m el. MSL). Note: Paleoliquefaction features (fissure fill and liquefied sediments) appear to offset, dragged, and disrupted fire-affected rocks and 0.9 to 1.3 ka year old, calibrated radiocarbon-dated shells (i.e. *Argopecton* sp., *Chione* sp.) and charcoal (photograph: Courtesy of Dennis Gallegos).






Fig. 29. Paleoliquefaction features exposed in a cut slope on the north side of Batiquitos Lagoon at Carlsbad, California (6.09–7.62 m el. MSL), located near the site seen in Fig. 26. Note: Shell horizons (*Argopecten* sp., *Chione* sp., and *Ostrea lurida*), charcoal, fire-affected rock fragments, and cobbles are offset, dragged, and liquefied upward into pre-existing fractures and into sand dikes.

vulsive, deposits in the Carlsbad to Encinitas area were likely produced by paleoseismic events, and hence are here deduced to be tsunamigenic in origin.

#### 7.2. Local tsunami mechanism

Although Emery (1960, p.124) indicated that southern California was not immune to tsunamis,




Fig. 30. View looking south of a storm drain trench wall cut into a coastal terrace exposing possible tsunami deposits at Carlsbad, California (15.54 m el. MSL). Note: Exposed are highly disturbed, contorted, chaotic, turbid sediments and debris overlying liquefied and dragged Bt horizons resting on a Tertiary sandstone abrasion platform. The upper half of this exposure interpreted as tsunamigenic in origin, for it also displays a distinct “hydraulic jump”, markedly different from documented paleoliquefaction features documented elsewhere.




Fig. 31. “Anomalous sediments” inferred to be tsunami deposits exposed in a vertical cut adjacent to a coastal lagoon at Carlsbad, California (6.70–9.14 m el. MSL). Note: The Jacob staff (scaled in 1' increments), rests in highly disturbed, mudflat sediments containing shells (*Chione* sp.), rock and clay fragments, overlain by chaotic, convulsive, turbid, sedimentary debris, including wood fragments, sharp angular pods of terrace deposits and Tertiary sandstone “bedrock”. These sediments are capped by lagoonal tidal flat organic-rich sands, “peat”, pieces of terrace sands and other chaotic sediments.

many considered that San Diego was relatively safe owing to the bordering, wide continental shelf that inhibits seismically-induced wave attack generated outside the region (Van Dorn, 1965). Recently, however, geophysical research shows that the southern California Continental Borderland is crossed by

many active faults with characteristics capable of producing large-scale, coseismic sea floor deformation during submarine earthquakes (Legg, 1991; Legg and Kennedy, 1991; Rivero et al., 2000; Grant and Rockwell, 2002). Additionally, local strong seismicity generated by any potential local earthquake sources,




Fig. 32. An abrupt, highly localized, disrupted, “chaotic” imbrication of marine gravels exposed in a vertical excavation in the lowest coastal terrace showing possible tsunamigenic and/or paleoliquefaction features at Carlsbad, California (3.04–3.35 m el. MSL). Note: Tape measure (in in.) for scale (1 in. = 2.54 cm).





Fig. 33. View of landward-fining sand layers forming lobes on on-lapping coastal terrace sands at Carlsbad, California (20.72–21.33 m el. MSL). Note: Putty knife rests against terrace sands for scale; also see irregular sharp contact with white sands at top of photo. These features are morphologically similar to tsunami deposits photographically documented during the 1994 Mindoro event in the Philippine Islands (Daag et al., 1995).

both onshore and offshore, could trigger large-scale slope failures and thereby generate local tsunamis (McCarthy et al., 1993; Borrero et al., 2001; Legg et al., 2003). Additionally, many large slope failures have been mapped in the southern California Continental Borderland (Field and Richmond, 1980; Clarke et al., 1985, 1987; Legg and Kamerling, 2003), even off Carlsbad and Encinitas (Greene and Kennedy, 1987; Clarke et al., 1987; Fig. 37). Paleoseismic investigations also show large coastal earthquakes occurred within the Holocene, and historically at least four measurable local tsunamis impacted the southern California region in 1812, 1862, 1927, and

1930 (McCulloch, 1985; Lander et al., 1993). Accordingly, evidence for paleotsunamis is abundant, and the potential for future impact is high (McCarthy et al., 1993; Legg et al., 2003, 2004).

## 8. Potential seismic sources

Based on the extent (730 km), character, and distribution of paleoseismic features and the residual evidence afforded by mima-mound topography and tsunamigenic deposits, it seems likely that the causative earthquakes were of at least  $M \sim 7.0$  (Kuhn et al.,




Fig. 34. North wall of a sewer-line trench cut into a terrace and exposing a possible paleotsunami deposit at Encinitas, California (93.26 m el. MSL). Note: The scale is located on a Tertiary-age sandstone abrasion platform, which is unconformably capped by a chaotic, turbid deposit containing small angular pieces of terrace and “parent” Tertiary sandstone and small rocks. Also note the “v-shaped upward climbing” chaotic features interpreted to result from an abrupt “hydraulic jump” created when tsunami waves reached a bedrock high.






Fig. 35. North wall of a cut slope exposing possible tsunamigenic and/or paleoliquefaction features at Encinitas, California (95.09 m el. MSL). Note: These features are the same as seen in Fig. 34 but are 3 m higher in the section.

2000, 2004). The specific fault(s) that triggered the liquefaction and likely secondary surface faulting is not well constrained. However, the likely seismic sources are the Newport–Inglewood/Rose Canyon Fault Zone (NIRC), ~4–6 km offshore and possibly under the study area (Kuhn et al., 2000, 2004), other Continental Borderland faults (Legg, 1991), and possible site-specific faults (Figs. 1 and 2).

#### 8.1. Newport–Inglewood/Rose Canyon fault zone

The Newport–Inglewood/Rose Canyon fault zone (NIRC) is the longest and most active in the north-coastal San Diego County area (Figs. 2 and 37). It is therefore the most likely seismogenic source for the observed paleoseismic features (Lindvall and Rockwell, 1995; Grant et al., 1999; Rivero et al., 2000;




Fig. 36. View looking southwest at abnormally tilted “sand flat” sediments exposed in a storm-drain trench cut into a coastal terrace at Carlsbad, California (28.65–29.26 m el. MSL). Note: The bedding dip varies from near-vertical (left) to 75 to 80 degrees (middle) to 46 degrees (right), to locally completely deformed. This exposure is morphologically a “twin” of deposits photographically documented during the tsunami that followed the 1994 Mindoro earthquake in the Philippine Islands (Daag et al., 1995).




Fig. 37. Portion of a “Geologic map of the inner-southern California Continental Margin”. Note: The red square (located at the Carlsbad Submarine Canyon) indicates sea floor faulting that cut strata of Holocene age. Also note the adjacent large submarine landslide (adapted from: Clarke et al., 1987).

Grant and Rockwell, 2002). It has also been suggested that the NIRC dips eastward beneath the coast (Kuhn et al., 1994), possibly merging with a deep northeast-dipping Neogene detachment fault system (Legg, 1991; Crouch and Suppe, 1993). Though presently somewhat speculative, an east-dipping seismogenic zone under Carlsbad to Encinitas may trap seismic wave energy in the hanging wall (Legg et al., 1994; Kuhn et al., 2000, 2004).

Approximately 1.5 km offshore lies the Carlsbad Submarine Canyon (Shepard and Emery, 1941), characterized by Holocene sea floor offset, and large subsurface landslides (Kennedy et al., 1985, 1987; Fig. 37). Similarly, Fischer et al. (1992) indicated that a coastal “tectonic dam” had beheaded the former drainage in the Carlsbad area. These tectonic features are directly offshore the Holocene liquefaction and tsunamigenic, chaotic features described in this paper (Kuhn et al., 2000).

North to northeast-trending branching and secondary fault zones also extend onshore from the NIRC (Hannan, 1973; Adams and Frost, 1981). These are exemplified by the Cristianitos fault zone near San Onofre (Western Geophysical, 1972; Moyle, 1973), and by an unnamed fault zone about 10 km north of Carlsbad (Euge et al., 1972). The Carlsbad to Encinitas area is located at a major transition in the NIRC, the “Carlsbad–Encinitas Overstep” of Fischer and Mills (1991), where the offshore Rose Canyon fault zone changes from a more northerly, transtensional trend to a more westerly, transpressional trend along the South Coast Offshore Zone (Kuhn et al., 2000). In sum, the NIRC, other nearby offshore faults, plus possible local onshore faults are all capable of generating relatively high-magnitude earthquakes. These collectively can produce the myriad of paleoseismic features now exposed in the north San Diego County area.

#### 8.2. Urbanization-caused increase in the hazards of seismically-induced liquefaction

The population of coastal North San Diego County is expected to triple over the next decade (SANDAG, 1991). Excavations of new residential tracts have provided excellent exposures of the neotectonic features described herein, and likely will continue largely on the marine and non-marine terraces. Terrace sediments are typically fine to medium sands, which are inherently susceptible to liquefaction when saturated. Although permanent ground water levels are presently low along the undeveloped coast, shallow perched water tables occur locally. Also, surface water also recharges perched and regional water levels via pre-existing fractures and sand dikes (Fig. 38). Almost all water for urban landscaping and recreational use is imported, so that soil moisture and infiltration will likely increase from the present 250 mm/year to an anticipated 2000 mm/year in the coming decade (Shlemon and Kuhn, 1997). Groundwater is typically perched within the Quaternary sediments (terrace deposits) on the underlying, relatively impermeable bedrock surface or “hardpan.” These “high-elevation” perched water levels will rise an expected ten-fold owing to urban-water infiltration (Shlemon, 2000). The active NIRC is within 4 to 6 km of the coast, and if an east-dipping fault plane exists at depth, coastal development will lie within the 5-km, near-




Fig. 38. Horizontal exposure made by grading on a coastal terrace at Carlsbad, California (29.26 m el. MSL). Note: Rainfall-induced surface runoff water flows into the subsurface along pre-existing fractures and sand dikes, thus recharging groundwater levels.

source zone for this active fault. Consequently, enhanced shaking may be expected from moderate to large earthquakes along this portion of the NIRC. Such strong shaking increases potential liquefaction of late Pleistocene sands and in the overlying engineered fills. Accordingly, the potential for coastal area, liquefaction-induced ground failures will likely increase during the coming decades. Structures may then be subject to permanent ground displacements associated with lateral spreading, as well as widespread sand blows and fissuring. Such hazards have heretofore been largely ignored.

## 9. Conclusions

Recent man-made exposures at Carlsbad and Encinitas reveal late Pleistocene to probable Holocene faults, tsunamigenic deposits, and ground-failure hazards. Features interpreted to be of seismic liquefaction origin are widespread and common in the study area. Paleoliquefaction features include sand-filled dikes and sills, lateral spreads, filled craterlets, fissures fills, and other unconsolidated sand deformations derived from underlying Tertiary and Pleistocene

marine and dune sediments. A paleoseismic liquefaction origin, rather than soft-sediment loading is deduced for these features based on internal stratigraphy, morphology, field setting, and proximity to active faults. The paleoliquefied sands are upward-fining, penetrate and locally deform regressive marine and prograding continental deposits and soils that mantle marine-terrace platforms at elevations from about 3 m to 130 m, and such marine-terrace platforms range in age from about 80 ka (lowest) to 400 ka. Some injection dikes and fissure fills are young, for they displace Holocene, Native-American Indian middens, burial grounds, and transitory camps, located on 60-m high terraces, far above modern regional ground levels. The paleoearthquakes were likely at least  $M \sim 7$ , and may have occurred during unusually wet times, when perched groundwater saturated underlying sediments. Many paleoseismic events probably originated on the nearby Newport–Inglewood/Rose Canyon fault system. Eastern dipping segments of the NIRC project under the north coast of San Diego County. Accordingly, the seismic hazard of this area may be substantially higher than heretofore assumed.

## Acknowledgments

Support for the publication of this article was provided by the International Foundation for Applied Research in the Natural Sciences (IFARNS). I am especially grateful to John Franklin of GeoSoils, Inc. for his helpful reviews of the manuscript, assistance in the graphic enhancement of all of the text figures, and use of the GeoSoils Inc. laboratory; to Dennis Gallegos for his review of the archaeological sections; and to Tom Dibblee Jr., Robert Dill, Mark Legg, David McArthur, Rick Riefner, Roy Shlemon, James Slosson, and Richard Terry for their thoughtful comments and constructive reviews of the manuscript, and to Jennifer Bauer for her exceptional field assistance.

## References

- Adams, M.A., Frost, E.G., 1981. The La Costa Avenue fault: an example of a secondary structure developed in a strike-slip zone. In: Abbott, P.L., O'Dunn, S. (Eds.), *Geological Investigations of the San Diego Coastal Plain*. San Diego Assn. of Geol. Guidebook, pp. 21–24.



- Aigner, T., 1985. Storm Depositional Systems: Dynamic Stratigraphy in Modern and Ancient Shallow-Marine Sequences. Springer-Verlag, Berlin (174 pp.).
- Amick, D., Gelinas, R., 1991. The search for evidence of large prehistoric earthquakes along the Atlantic seaboard. *Science* 251, 655–658.
- Anderson, Warren, Associates, 1977. Results of microfossil identification and geologic age correlation, vicinity of San Onofre Generating Station, California. Unpubl. report prepared for Southern California Edison Company (Rosemead), California. Reprinted In: Legg, M.R., Kuhn, G.G., Shlemon R.J. (Eds.), Neotectonics and Coastal Instability: Orange and Northern San Diego Counties, California. Joint field conference, Am. Assn. Of Petroleum Geologists, Pacific Section and SPE, Western Section, held in Long Beach, California on June 19–22, 2000, 2: pp. II-138 to II-147.
- Arkley, R.J., Brown, H.C., 1954. The origin of mima mound (hogwallow) microrelief in the far western states. *Soil Sci. Soc. Proc.*, 195–199.
- Astiz, L., Shearer, P.M., 2000. Earthquake locations in the Inner Continental Borderland, offshore southern California. *Bull. Seismol. Soc. Am.* 90 (2), 425–449.
- Aten, L.E., 1981. Archaeological evidence for pimple (prairie) mound genesis. *Science* 213, 1374–1376.
- Atwater, B.F., Burrell, K.S., Cisternas, M., Higman, V.B., Barnhardt, W.A., Kayen, R.E., Minasian, D., Satake, K., Shimokawa, K., Haraguchi, T., Takada, K., Baker, D., Nakata, T., 2001. Grouted sediment slices show signs of earthquake shaking. *Eos*, 82 (49) 603, 609.
- Aydin, A., Johnson, A.M., Fleming, R.W., 1992. Right-lateral-reverse surface rupture along the San Andreas and Sargent faults associated with the October 17, 1989, Loma Prieta, California, earthquake. *Geology* 20, 1063–1067.
- Baldwin, J.N., Kelson, K.I., Randolph, C.E., 2000. Late Quaternary fold deformation along the Northridge Hills fault, Northridge, California: deformation coincident with past Northridge blind-thrust earthquakes and other nearby structures? *Bull. Seismol. Soc. Am.* 90 (3), 629–642.
- Barnes, G.W., 1879. The hillock or mound formations of San Diego County. In: Packard Jr., A.S., Cope, E.D. (Eds.), *The American Naturalist*, vol. XIII, pp. 565–571.
- Barrows, A.G., 1974. A review of the geology and earthquake history of the Newport–Inglewood structural zone, southern California. *Spec. Rep., Calif. Div. Mines Geol.* 114 (115 pp.).
- Bartholomew, M.J., Stickney, M.C., Wilde, E.M., Dundas, R.G., 2002. Late Quaternary paleoseismites: syndepositional features and section restoration used to indicate paleoseismicity and stress-field orientations during faulting along the main Lima Reservoir fault, southwestern Montana. In: Ettensohn, F.R., Rast, N., Brett, C.E. (Eds.), *Ancient Seismites*, Spec. Pap.-Geol. Soc. Am., vol. 359, pp. 29–48.
- Berg, A.W., 1990. Formation of mima mounds: a seismic hypothesis. *Geology* 18, 281–284.
- Blake, W.P., 1856a. Geological report, explorations and surveys for a railroad route from the Mississippi River to the Pacific Ocean. 33rd Congress, 2nd Session, Session Document, vol. 78, pp. 122–139.
- Blake, W.P., 1856b. Observations on the physical geography and physical geology of the coast of California from Bodega Bay to San Diego. US Coast and Geodetic Survey Rept. to the Superintendent for Year of 1855., pp. 376–398.
- Bohannon, R.G., Geist, E., 1998. Upper crustal structure and Neogene tectonic development of the California Continental Borderland. *Geol. Soc. Amer. Bull.* 110, 779–800.
- Bondevik, S., Svensen, J.I., Mangerud, J., 1995. Identification of tsunami deposits in lake basins, Western Norway (abstr.). *Tsunami Deposits: Geological Warnings of Future Inundation*. U S Geological Survey/NOAA Sponsored Conference, held on May 22–23, 1995 at the University of Washington at Seattle, Abstracts with Programs, pp. 21–22.
- Borrero, J.C., Ortiz, M., Titov, V., Synolakis, C.E., 1997. Field survey of Mexican tsunami produces new data, unusual photos. *Eos* 78 (8), 87.
- Borrero, J.C., Dolan, J.F., Synolakis, C.E., 2001. Tsunamis within the eastern Santa Barbara Channel. *Geophys. Res. Lett.* 28 (4), 643–646.
- Bourgeois, J., 1995. Tsunami deposits (abstr.). *Tsunami Deposits: Geological Warnings of Future Inundation: US Geological Survey/NOAA sponsored conference*, held on May 22–23, 1995 at the University of Washington at Seattle, Abstracts with Programs, p. 2.
- Breschini, G.S., Haversat, T., Erlandson, J., 1992. California Radiocarbon Dates. Coyote Press (147 pp.).
- Brookfield, M., 2004. Deposits of tsunamis and their recognition. Rapid and Catastrophic Environmental Changes in the Holocene and Human Response. 1st Joint Meeting of the International Geological Correlation Programme (IGCP 490) and IGSU Environmental Catastrophes held on January 4–18, 2004 in the Atar Desert/banc d'Arguin wetland.
- Carey, S., Morelli, D., Sigurdsson, H., Bronto, S., 2001. Tsunami deposits from major explosive eruptions: an example from the 1883 eruption of Krakatau. *Geology* 29 (4), 347–350.
- Carter, G.F., 1957. Pleistocene Man at San Diego. Johns Hopkins Press, Baltimore, MA (400 pp.).
- Carver, G.A., Gilpin, L.M., 1995. Observations of deposits resulting from the 1964 tsunami on Kodiak Island, Alaska (abstr.). *Tsunami Deposits: Geological Warnings of Future Inundation*. US Geological Survey/NOAA-Sponsored Conference, held on May 22–23, 1995 at the University of Washington, at Seattle, Abstracts with Programs, p. 6.
- Clague, J.J., Hutchinson, I., Mathewes, R.W., Patterson, R.T., 1999. Evidence for late Holocene tsunamis at Catala Lake. *British Columbia (abstr.)*. *J. Coast. Res.* 15 (1), 45–60.
- Clarke, S.H., Greene, H.G., Kennedy, M.P., 1985. Identifying potentially active faults and unstable slopes offshore. In: Ziony, J.I. (Ed.), *Evaluating Earthquake Hazards in the Los Angeles Region—An Earth Science Perspective*, U. S. Geol. Surv. Prof. Pap., vol. 1360, pp. 347–373.
- Clarke, S.H., Greene, H.G., Kennedy, M.P., Vedder, J.G., 1987. Geologic map of the inner southern California continental margin. Map no. 1A. In: Greene, H.G., Kennedy, M.P. (Eds.), *Geologic Map Series of the California Continental Margin*. *Bull.-Calif., Div. Mines Geol.* 207, Area 1 of 7, (NOS 1206N-16), Scale 1:250,000.

- Coleman, P.J., 1968. Tsunamis as geological agents. *J. Geol. Soc. Aust.* 15 (pt. 2), 267–273.
- Costa, P., 2004. Detecting storm and tsunami deposits in coastal lagoons: Preliminary results from Lagoa de Obido (West Portugal). Rapid and Catastrophic Environmental Changes in the Holocene and Human Response, 1st Joint Meeting of International Geological Correlation Programme (IGCP 490) and ICSU Environmental Catastrophes, held on January 4–18, 2004 at the Atar desert/banc d'Arguin wet-land, Mauritania, Abstracts with Programs (4 pp.).
- Cox, G.W., 1990. Form of mima mounds in relation to occupancy by pocket gophers. *J. Mammal.* 71, 90–94.
- Cox, G.W., 1991. Pocket gophers and mima terrain in North America. *Nat. Areas J.* 11, 193–198.
- Cox, R.T., Van Arsdales, R.B., Harris, J.B., Larson, D., 2001. Neotectonics of the southeastern Reelfoot rift zone margin, central United States, and implications for regional strain accumulation. *Geology* 29 (5), 419–422.
- Crouch, J.K., Suppe, J., 1993. Late Cenozoic tectonic evolution of the Los Angeles basin and Inner California borderland: a model for core complex-like crustal extension. *Geol. Soc. Amer. Bull.* 105, 1415–1434.
- Crowell, J.C., 1974. Origin of late Cenozoic basins in southern California. In: Dickinson, W.R. (Ed.), *Tectonics and Sedimentation, Spec. Publ.-Soc. Econ. Paleontol. Mineral.*, vol. 22, pp. 190–204.
- Daag, A.S., De los Reyes, P.J., Tubianosa, B.S., Javier, D.V., Punongbayan, R.S., 1995. Tsunami deposit of the November, 1994 Mindoro Earthquake, Philippines (abstr.). *Tsunami Deposits: Geological Warnings of Future Inundation*. US Geological Survey/NOAA Sponsored Conference, held on May 22–23, 1995 at the University of Washington at Seattle, Abstracts with Programs, p. 3.
- Dahlquist, W.W., Scheffer, V.B., 1942. The origin of the mima mounds of western Washington. *J. Geol.* 50, 68–84.
- Darigo, N.J., Osborne, R.H., 1986. Quaternary stratigraphy and sedimentation of the Inner Continental Shelf, San Diego County, California. In: Knight, R.J., McLean, J.R. (Eds.), *Shelf Sands and Sandstones, Mem.-Can. Soc. Pet. Geol.*, vol. II, pp. 73–98.
- Davis, T.L., Namson, J.S., Yerkes, R.F., 1989. A cross section of the Los Angeles area: seismically active fold and thrust belt, the 1987 Whittier Narrows earthquake, and earthquake hazard. *J. Geophys. Res.* 94, 9644–9664.
- Dawson, A.G., 1994. Geomorphological effects of tsunami runup and backwash. *Geomorphology* 10, 83–94.
- Dawson, A.G., 1995. Geomorphological processes associated with tsunami runup (abstr.). *Tsunami Deposits: Geologic Warnings of Future Inundation*. US Geological/NOAA Sponsored Conference, held on May 22–23, 1995 at the University of Washington at Seattle, Abstracts with Programs, p. 19.
- Dawson, A.G., 1999. Linking tsunami deposits, submarine slides, and offshore earthquakes. *Quat. Int.* 60, 119–126.
- Dawson, A.G., Shi, S., Dawson, S., Takahashi, T., Shuto, N., 1996. Coastal sedimentation associated with the June 2nd and 3rd, 1994 tsunami in Rajegwesi, Java. *Quat. Sci. Rev.* 15, 901–912.
- Dietz, R.S., 1945. The small mounds of the gulf coastal plain. *Science* 102 (2658), 596–597.
- Dutton, C.E., 1889. The Charleston earthquake of August 31, 1886. US Geological Survey Ninth Annual Report of the Directorate (1887–88), pp. 209–528.
- Ehlig, P.L., 1977. Geologic report on the area adjacent to the San Onofre Nuclear Generating Station, Northwestern San Diego County, California. Unpubl. report prepared for Southern California Edison Company (Rosemead), California. Reprinted In: Legg, M.R., Kuhn, G.G., Shlemon, R.J. (Eds.), *Neotectonics and Coastal Instability: Orange and Northern San Diego Counties, California*. Joint field conference, Am. Assn. Of Petroleum Geologists, Pacific Section and SPE, Western Section, held in Long Beach, California on June 19–22, 2000, 2: pp. II-112 to II-137.
- Eiby, G.A., 1982. Earthquakes and tsunamis in a region of diapiric folding. *Tectonophysics* 85, T1–T8.
- Eisenberg, L.I., 1985. Pleistocene faults and marine terraces, northern San Diego County, California. In: Abbott, P.L. (Ed.), *On the Manner of the Eocene Strata in Northern San Diego County*. San Diego Assn. of Geologists, pp. 87–91.
- Ellis, A.J., Lee, C., 1919. Geology and ground waters of the western part of San Diego County, California. US Geological Survey Water-Supply Paper 446 (321 pp.).
- Emery, K.O., 1950a. Ironstone concretions and beach ridges of San Diego County, California. *California Div. Mines Geol.* 46 (2), 213–221.
- Emery, K.O., 1950b. Contorted pleistocene strata at Newport Beach, California. *J. Sediment. Petrol.* 20 (2), 111–115.
- Emery, K.O., 1960. The Sea off Southern California—A Modern Habitat of Petroleum. John Wiley and Sons, Inc., New York (366 pp.).
- Emery, K.O., Kuhn, G.G., 1982. Sea cliffs: Their processes, profiles, and classification. *Geol. Soc. Amer. Bull.* 93 (7), 644–654.
- Ettensohn, F.R., Kulp, M.R., Rast, N., 2002. Interpreting ancient marine seismites and apparent epicentral areas for paleo-earthquakes, Middle Ordovician Lexington Limestone, central Kentucky. In: Ettensohn, F.R., Rast, N., Brett, C.E. (Eds.), *Ancient Seismites, Spec. Pap.-Geol. Soc. Am.*, vol. 359, pp. 177–190.
- Euge, K.M., Miller, D.A., Palmer, L.A., 1972. Evidence for a possible onshore extension of the Rose Canyon fault in the vicinity of Oceanside, California (abstr.). *Geol. Soc. of Am., Cordilleran Section Meeting*, held in Portland, Oregon, Abstracts with programs, 5: p. 39. Reprinted In: Legg, M.R., Kuhn, M.R., Shlemon R.J. (Eds.), *Neotectonics and Coastal Instability: Orange and Northern San Diego Counties, California*. Joint field conference, Am. Assn. Of Petroleum Geologists, Pacific Section, and SPE, Western Section, held in Long Beach on June 19–22, 2000, 2: pp. II-1 to II-18.
- Fairbanks, H.W., 1893. Geology of San Diego County, also portions of Orange and San Bernardino counties. *Calif. State Min. Bur.*, 11th Rept., pp. 76–120.
- Field, M.E., Richmond, W.C., 1980. Sedimentary and structural patterns on the northern Santa Rosa-Rosa Ridge, southern California. *Mar. Geol.* 24, 79–98.
- Fischer, P.J., Mills, G.I., 1991. The offshore Newport-Inglewood/Rose Canyon Fault Zone, California: Structure, segmentation, and tectonics. In: Abbott, P.L., Elliott, W.J. (Eds.), *Environmental Perils, San Diego Region*. *Geol. Soc. of Am.*

- Field Trip Guidebook Prepared by the San Diego Assn. of Geologists, pp. 17–36.
- Fischer, P.J., Gorsline, D.S., Shlemon, R.J., 1992. Late Quaternary geology of the Dana Point–San Onofre–Carlsbad Margin, California. In: Heath, E.G., Lewis, W.L. (Eds.), *The Regressive Pleistocene Shoreline. Coastal California*, pp. 195–218.
- Foster, I.D.L., Albon, A.J., Bardell, K.M., Fletcher, J.L., Jardine, T.C., Mothers, R.J., Pritchard, M.A., Turner, S.E., 1991. High energy coastal sedimentary deposits; an evaluation of depositional processes in Southwest England. *Earth Surf. Process. Landf.* 16 (4), 341–356.
- Franklin, J.P., Kuhn, G.G., 2000. Paleoseismic features exposed by trenching the lowest coastal terrace at Carlsbad, California. In: Shlemon, R.J., Kuhn, G.G., Legg, M.R. (Eds.), *Neotectonics and Coastal Instability, Orange and Northern San Diego Counties, California. Joint Field Conference*, Am. Assn. of Petroleum Geologists, Pacific Section, and SPE, Western Section, held in Long Beach, California on June 19–22, 1, pp. I-1–I-13.
- Fugro, Inc., 1975a. Summary of geomorphic and age data for the first emergent terrace (Qt1) at the San Onofre Nuclear Generating Station: Unpubl. report prepared for Southern California Edison Company (Rosemead), California. Reprinted In: Legg, M.R., Kuhn, G.G., Shlemon R.J. (Eds.), *Neotectonics and Coastal Instability, Orange and Northern San Diego Counties, California. Joint field conference*, Am. Assn. Of Petroleum Geologists, Pacific Section, and SPE, Western Section, held in Long Beach, California on June 19–22, 2000, 2: pp. II-19 to II-49.
- Fugro, Inc., 1975b. Geomorphic analysis of terraces in San Juan and Bell Canyons, Orange County, California. Unpubl. report prepared for Southern California Edison Company (Rosemead), California. Reprinted In: Legg, M.R., Kuhn, G.G., Shlemon R.J. (Eds.), *Neotectonics and Coastal Instability, Orange and Northern San Diego Counties, California. Joint field conference*, Am. Assn. Of Petroleum Geologists, Pacific Section, and SPE, Western Section, held in Long Beach, California on June 19–22, 2000, 2: pp. II-50 to II-60.
- Fugro, Inc., 1977a. Supplemental report of geological investigations, Trail #6 landslide [San Onofre State Beach] and Horno Canyon [Camp Pendleton], southeast of San Onofre Nuclear Generating Station: Unpubl. report prepared for Southern California Edison Company (Rosemead), California, Project #77-206-01. Reprinted In: Legg, M.R., Kuhn, G.G., Shlemon R.J. (Eds.), *Neotectonics and Coastal Instability: Orange and San Diego Counties, California. Joint field conference*, Am. Assn. of Petroleum Geologists, Pacific Section, and SPE, Western Section, held in Long Beach, California on June 19–22, 2000, 2: pp. II-61 to II-92.
- Fugro, Inc., 1977b. Geological investigations of offsets in Target Canyon, Camp Pendleton, California. Unpubl. report prepared for Southern California Edison Company (Rosemead). Reprinted In: Legg, M.R., Kuhn, G.G., Shlemon R.J. (Eds.), *Neotectonics and Coastal Instability: Orange and Northern San Diego Counties, California. Joint field conference*, Am. Assn. Of Petroleum Geologists, Pacific Section, and SPE, Western Section, held in Long Beach on June 19–22, 2000, 2: pp. II-93 to II-162.
- Fuller, M.L., 1912. The New Madrid earthquake. *U.S. Geol. Surv. Bull.* 494 (119 pp.).
- Gallegos, D.R., 1987. A review and synthesis of environmental and cultural material for the Batiquitos Lagoon region. In: Hector, S.M., Van Wormer, S.R. (Eds.), *San Dieguito–La Jolla: Chronology and Controversy*, San Diego Archaeological Soc. Res. Papers, vol. 1, pp. 23–34.
- Gallegos, D.R., 2002. Southern California in transition: Late Holocene occupation of southern San Diego County. In: Erlandson, J.M., Jones, T.L. (Eds.), *Catalysts to Complexity: Late Holocene Societies of the California Coast. Perspectives in California Archaeology*, J.E. Arnold (Senior Series Editor), Cotsen Institute of Archaeology, Univ. of California, Los Angeles, California, 6 27–40.
- Gallegos, D.R., 1997. Batiquitos Lagoon monitoring program: Archaeological test at Site CA-SDI-#11953/#600, Carlsbad, California. Unpubl. report prepared for Tetra Tech., San Diego, California, 32 pp.
- Galli, P., Ferrelli, L., 1995. A methodological approach for historical liquefaction research. In: Serva, L. (Ed.), *Perspectives in Paleoseismology*, Assn. of Eng. Geologists, Spec. Pub., vol. 6, pp. 35–48 (Chap. 4).
- Goodyear, W.A., 1888. San Diego County. *California State Mining Bureau, 8th Annual Report*, 516–528.
- Graham, S.A., Bachman, S.B., 1983. Structural controls on submarine-fan geometry and internal architecture: upper La Jolla fan system, offshore Southern California. *Bull. Am. Assoc. Pet. Geol.* 67 (1), 83–96.
- Grant, L.B., Rockwell, T.K., 2002. A northward-propagating earthquake sequence in coastal southern California? *Seismol. Res. Lett.* 73 (4), 461–469.
- Grant, L.B., Mueller, K.L., Gath, E.M., Cheng, H., Edwards, R.L., Munro, R., Kennedy, G.L., 1999. Late Quaternary uplift and earthquake potential of the San Joaquin Hills, southern Los Angeles basin, California. *Geology* 27 (11), 1031–1034.
- Grant, L.B., Ballenger, L.J., Runnerstrom, E.E., 2002. Coastal uplift of the San Joaquin Hills, southern Los Angeles Basin, California, by a large earthquake since A.D. 1635. *Bull. Seismol. Soc. Am.* 92 (2), 590–599.
- Greb, S.F., Dever Jr., G.R., 2002. Critical evaluation of possible seismites: examples from the carboniferous of the Appalachian basin. In: Ettensohn, F.R., Rast, N., Brett, C.E. (Eds.), *Ancient Seismites*, Spec. Pap.-Geol. Soc. Am., vol. 359, pp. 109–125.
- Greene, H. G., Kennedy, M. P., (Eds.), 1987. *Geological Map Series of the California Continental Margin*. California Div. Mines and Geol. Bull. No. 207, 7 Areas, Scale 1:250,000.
- Guptill, P.D., Heath, E.W., 1981. Surface faulting along the Newport–Inglewood zone of deformation. *Calif. Geol.*, 142–148.
- Hall, N.T., 1984. Holocene history of the San Andreas fault between Crystal Springs Reservoir and San Andreas Dam, San Mateo County, California. *Bull. Seismol. Soc. Am.* 74, 281–299.
- Hand, B.M., Emery, K.O., 1963. Turbidites and topography of the north end of San Diego Trough, California. *J. Geol.* 72, 526–542.
- Hannan, D.L., 1973. Faulting in the Oceanside, Carlsbad, and Vista areas, northern San Diego County, California. In: Ross, A., Dowlen, R.J. (Eds.), *Studies on the Geology of Camp Pendleton*



- and western San Diego County, California. San Diego Assn. of Geologists, pp. 56–59, 2 maps.
- Harden, J.W., Matti, J.C., 1989. Holocene and late Pleistocene slip rates on the San Andreas fault in Yucaipa, California, using displaced alluvial-fan deposits and soil chronology. *Geol. Soc. Amer. Bull.* 101, 1107–1117.
- Harp, E.L., Jibson, R.W., Kayen, R.E., Keefer, D.K., Sherrod, B.L., Carver, G.A., Collins, B.D., Moss, E.S., Sitar, N., 2003. Landslides and liquefaction triggered by the M 7.9 Denali Fault earthquake of 3 November 2002. *GSA Today* 13 (8), 4–10.
- Hertlein, L.G., Grant IV, U.S., 1944. The geology and paleontology of the marine Pliocene of San Diego, California. San Diego Soc. Nat. Hist. Mem., 2: pt. 1, Geology Section, pp. 18–27, 57.
- Hertlein, L.G., Grant IV, U.S., 1954. Geology of the Oceanside-San Diego coastal area, southern California. *Geol. South. Calif. Bull.* 170 (1), 53–63.
- Hindson, R.A., Andrade, A., 1999. Sedimentation and hydrodynamic processes associated with the tsunami generated by the 1755 Lisbon earthquake. *Quat. Int.* 56, 27–38.
- Holzer, T.L., Clark, M.M., 1993. Sand boils without earthquakes. *Geology* 21, 873–876.
- Holzer, T.L., Youd, T.L., Hanks, T.C., 1989. Dynamics of liquefaction during the 1987 Superstition Hills, California earthquake. *Science* 244, 56–59.
- Holzer, T.L., Noce, T.E., Bennett, M.J., Di Alessandro, C., Boatwright, J., Tinsley III, J.C., Sell, R.W., Rosenberg, L.L., 2004. Liquefaction-induced lateral spreading in Oceano, California, during the 2003 San Simeon earthquake. Open-File Rep. (USGS) 2004-1269 (41 pp.).
- Hough, S.E., 2001. Triggered earthquakes and the 1811–1812 New Madrid, Central United States, earthquake sequence. *Bull. Seismol. Soc. Am.* 91, 1574–1581.
- Housner, G.W., 1958. The mechanism of sandblows. *Bull. Seismol. Soc. Am.* 48, 155–161.
- Imamura, F., Subandono, D., Watson, G., Moore, A., Takahashi, T., Matsutomi, H., Hidayet, R., 1997. Irian Jaya earthquake and tsunami cause serious damage. *Eos*, 78 (19) 198, 201.
- Jachens, R.C., Langenheim, V.E., Matti, J.C., 2002. Relationship of the 1999 Hector Mine and 1992 Landers fault ruptures to offsets on Neogene faults and distribution of late Cenozoic Basins in eastern California shear zone. *Bull. Seismol. Soc. Am.* 92 (4), 1592–1605.
- Kawata, Y., Benson, B.C., Borrero, J.C., Davies, H.L., deLange, W.P., Imamura, F., Letz, H., Nott, J., Synolakis, C., 1999. Tsunami in Papua New Guinea was as intense as first thought. *Eos*, 80 (9): pp.101, 104, 105.
- Kelsey, H.M., Nelson, A.R., Hemphill-Haley, E., 1995. Properties and depositional characteristics of tsunamis in south coastal Oregon from a paired coastal lake and marsh study (abstr.). Tsunami Deposits: Geologic Warnings of Future Inundations. US Geological Survey/NOAA Sponsored Conference, held on May 22–23, 1995 at the University of Washington at Seattle, Abstracts with Programs, p. 20.
- Kennedy, M.P., 1975. Geology of San Diego metropolitan area, California, section A—Western San Diego metropolitan area. California Div. Mines Geol., 200.
- Kennedy, M.P., Tan, S.S., Chapman, R.H., Chase, C.W., 1975. Character and recency of faulting, San Diego Metropolitan area. Spec. Rep., Calif. Div. Mines Geol., 123.
- Kennedy, M.P., Clarke, S.H., Greene, H.G., Legg, M.R., 1980a. Recency and character of faulting offshore Metropolitan San Diego, California. California Div. Mines and Geol., Map. Sheet 41, Scale 1:50,000.
- Kennedy, M.P., Clarke, S.H., Greene, H.G., Legg, M.R., 1980b. Recency and character of faulting offshore San Diego, California. California Div. Mines and Geol., Map Sheet 42, Scale 1:50,000.
- Kennedy, M.P., Clarke, S.H., Greene, H.G., Lonsdale, P.F., 1985. Observations from DSRV ALVIN of Quaternary faulting on the southern California continental margin. Open-File Rep. (U. S. Geol. Surv.) 85-39, 1–24.
- Kennedy, M.P., Greene, H.G., Clarke, S.H., 1987. Geology of the California continental margin: explanation of the continental margin geologic map series. *Bull.-Calif. Div. Mines Geol.* 207 (110 pp.).
- Kern, J.P., Rockwell, T.K., 1992. Chronology and deformation of Quaternary marine shorelines, San Diego County, California. Quaternary Coasts of the US: Marine and Lacustrine Systems, Soc. of Economic Paleontologists and Mineralogists (SEPM), Special Report, vol. 48.
- Krinitzsky, E.L., 1949. Origin of pimple mounds. *Am. J. Sci.* 248, 355.
- Kuenen, P.H., 1947. Two problems of marine geology: atolls and canyons. *Verh. K. Ned. Akad. Wet. Natuurkd.* 43 (3), 1–69.
- Kuenen, P.H., 1957. Sole markings of graded graywacke beds. *J. Geol.* 65, 241–258.
- Kuhn, G.G., 2000. Sea cliff, canyon, and coastal terrace erosion between 1887 and 2000: San Onofre State Beach, Camp Pendleton Marine Corps Base, San Diego County, California. In: Shlemon, R.J., Kuhn, G.G., Legg, M.R. (Eds.), *Neotectonics and Coastal Instability: Orange and Northern San Diego Counties, California*. Joint field conference, Am. Assn. of Petroleum Geologists, Pacific Section, and Soc. of Petroleum Engineers, Western Section, held in Long Beach, California, on June 19–22, 2000, I: pp. I-31 to I-87, 267–284.
- Kuhn, G.G., Shepard, F.P., 1984. Sea Cliffs, Beaches, and Coastal Valleys of San Diego County: Some Amazing Histories and Some Horrifying Implications. Univ. of California Press, Berkeley (193 pp.).
- Kuhn, G.G., Legg, M.R., Frost, E.G., 1994. Large prehistoric earthquake(s) in coastal San Diego County, California. Proceedings of the Workshop on Paleoseismology, held at Marshall, California on 18–22 September, 1994, US Geological Survey Open-File Report, vol. 94-568, pp. 100–102.
- Kuhn, G.G., Johnson, J.A., Shlemon, R.J., 1995a. Paleoliquefaction evidence for prehistoric earthquakes in north coastal San Diego County, California. Am. Geophysical Union (AGU) Fall Meeting, held on 11–15 December, 1995 at San Francisco, California, Abstracts with Programs, Session No. S12C, No.7, p. F362.
- Kuhn, G.G., Legg, M.R., Frost, E.G., 1995b. Late prehistoric earthquake(s) in coastal San Diego County, California. Am. Assn. of Petroleum Geologists, Pacific Section newsletter (feature article), no. 1, 3 pp.

- Kuhn, G.G., Legg, M.R., Johnson, J.A., Frost, E.C., 1995c. Paleoliquefaction evidence for large prehistoric earthquake(s) in north coastal San Diego, County, California. In: Munasinghe, T., Rosenberg, P. (Eds.), *Geology and Natural Resources of Coastal San Diego County, California*. San Diego Assn. of Geologists Guidebook, pp. 16–24.
- Kuhn, G.G., Legg, M.R., Shlemon, R.J., Bauer, J.L., 2000. Neotectonics in the north coastal area, San Diego County, California. In: Shlemon, R.J., Kuhn, G.G., Legg, M.R. (Eds.), *Neotectonics and Coastal Instability, Orange and Northern San Diego Counties, California*. Joint Field Conference, Am. Assn. of Petroleum Geologists, Pacific Section, and Soc. of Petroleum Engineers, Western Section, held in Long Beach, California, on June 19–22, 1. , pp. 1-88–1-103.
- Kuhn, G.G., Legg, M.R., Franklin, J.P., 2004. Paleoseismology of blind and offshore faults, north coast San Diego County, California, USA. In: Doolin, D., Kammerer, A., Nogami, T., Seed, R.B., Towhata, I. (Eds.), *Proceedings of the 11th International Conf. on Soil Dynamics and Earthquake Engineering and 3rd International Conf. on Earthquakes Geotechnical Engineering*, held on 7–9 January, 2004 at Berkeley, California, 2, pp. 597–604.
- Lajoie, K.R., Ponti, D.J., Powell, C.L., Mathieson, S.A., Sarna-Wojcicki, A.M., 1991. Emergent marine strandlines and associated sediments, coastal California: a record of Quaternary sea-level fluctuations, vertical tectonic movements, climatic changes, and coastal processes. In: Morrison, R.B. (Ed.), *Quaternary Nonglacial Geology: Contemporaneous U.S. The Geology of North America; K-2*. Geol. Soc. of Amer, Boulder, Colorado, pp. 190–214.
- Lamont, A., 1936. Palaeozoic Seismicity. *Nature* (London) 138, 243–244.
- Lander, J.F., Lockridge, P.A., Kozuch, M.J., 1993. Tsunamis affecting the West Coast of the United States. NOAA, National Geophysical Data Center, Key to geophysical records, Documentation no. 29, Boulder, Colorado, 242 pp.
- Landuzzi, A., Nichols, R., Vai, G.B., 1995. Sand volcanoes in the Marnoso-Arenacea Formation (Miocene, Northern Apennines, Italy): new evidence of paleoseismicity. In: Serva, L. (Ed.), *Perspectives in Paleoseismology*, Assn. of Eng. Geol. Spec. Pub., vol. 6, pp. 49–61 (Chap. 5).
- Larson, E.S., 1948. Batholith and associated rocks of Corona, Elsinore, and San Luis Rey Quadrangles, southern California. *Mem. Geol. Soc. Amer.* 29 (182 pp.).
- Lawson, A.C., 1908. The California earthquake of April 18, 1906: [Effect of the earthquake upon underground waters, significance of the phenomena]. In *Report of the state earthquake investigation commission*, Pub. No. 87, 1: part 1, pp. 402–409, Carnegie Institution of Washington, Wash. D.C.
- Lee, K.L., Seed, H.B., 1967. Cyclic stress conditions causing liquefaction of sand. *J. Soil Mech. Found. Div. (ASCE)* 93 (SMI), 47–70.
- Legg, M.R., 1985. Geologic structure and tectonics of the inner continental borderland offshore northern Baja California, Mexico. Ph.D. dissertation, University of California, Santa Barbara, Santa Barbara, California, 410 pp.
- Legg, M.R., 1987. Earthquake epicenters and selected fault plane solutions of the inner-southern California continental margin. In: Greene, H.G., Kennedy, M.P. (Eds.), *Geologic Map Series of the California Continental Margin*. California Div. of Mines and Geology, Area 1 of 7, (NOS 1206N-16), Map no. 1B, Scale 1:250,000.
- Legg, M.R., 1991. Developments in understanding the tectonic evolution of the California Continental Borderland. In: Osborne, R.H. (Ed.), *Soc. of Economic Paleontologists and Mineralogists (SEPM), Shepard Commemorative Volume*, pp. 291–312.
- Legg, M.R., 2004. Pers. Com.
- Legg, M.R., Kamberling, M.J., 2003. Large-scale basement-involved landslides, California Continental Borderland. *Pure Appl. Geophys.* 160, 2033–2051.
- Legg, M.R., Kennedy, M.P., 1979. Faulting offshore San Diego and northern Baja California. In: Abbott, P.L., Elliott, W.J. (Eds.), *Earthquakes and other Perils—San Diego Region*. San Diego Assn. of Geol. Guidebook, pp. 29–46.
- Legg, M.R., Kennedy, M.P., 1991. Oblique divergence and convergence in the California Continental Borderland. In: Abbott, P.L., Elliott, W.J. (Eds.), *Environmental Perils of the San Diego Region, California*. San Diego Assn. of Geol. Guidebook, pp. 1–16.
- Legg, M.R., Nicholson, C., Sorlien, C., 1992. Active faulting and tectonics of the inner California continental borderland (abstr.). *Trans.-Am. Geophys. Union* 73, 588.
- Legg, M.R., Kuhn, G.G., Frost, E.G., 2004. Episodic uplift of the southern California coast by large strike-slip earthquakes (abstr.). *Am. Geophysical Union (AGU), Fall Meeting*, held on December 5–9, 1994 at San Francisco, California. Abstracts with programs, session no. T31B-17.
- Legg, M.R., Borrero, J.C., Synolakis, C.E., 2003. Evaluate tsunami risk to southern California coastal cities. Final report prepared for the Earthquake Engineering Research Institute (EERI), Oakland, California, under a 2002 EERI/FEMA NEHERP Professional Fellowship, 32 pp.
- Legg, M.R., EERI, M., Borrero, J.C., Synolakis, C.E., 2004. Tsunami hazards associated with the Catalina Fault in southern California. *Earthq. Spectra* 20 (3), 917–950.
- Li, Y., Craven, J., Schweig, E.S., Obermeier, S.F., 1996. Sand boils induced by the 1993 Mississippi River flood: could they one day be misinterpreted as earthquake-induced liquefaction? *Geology* 24, 171–174.
- Lienkaemper, J.J., 2001. 1857 slip on the San Andreas fault southeast of Cholame, California. *Bull. Seismol. Soc. Am.* 91 (6), 1659–1672.
- Lindvall, S.C., Rockwell, T.K., 1995. Holocene activity of the Rose Canyon fault zone in San Diego, California. *J. Geophys. Res.* 100 (B12), 24,121–24,132.
- Lowe, D.R., 1975. Water escape structures in coarse-grained sediments. *Sedimentology* 22, 157–204.
- Lowe, D.R., 1976. Subaqueous liquefied and fluidized sediment flows and their deposits. *Sedimentology* 23, 285–308.
- Lucci, F.R., 1995. Sedimentological indicators of paleoseismicity. In: Serva, L. (Ed.), *Perspectives in Paleoseismology*, Assn. of Eng. Geologists, Spec. Pub., vol. 6, pp. 7–17 (Chap. 2).

- Mariotti, G., Corda, L., Brandano, M.B., Civitelli, G., 2002. Indicators of paleoseismicity in the middle Miocene Guadagnolo Formation, central Apennines, Italy. In: Ettensohn, F.R., Rast, N., Brett, C.E. (Eds.), *Ancient Seismites*, Spec. Pap.-Geol. Soc. Am., vol. 359, pp. 87–98.
- Marple, R.T., Schweig III, E.S., 1992. Remote sensing of alluvial terrain in a humid, tectonically active setting: the New Madrid Seismic Zone. *Photogramm. Eng. Remote Sensing* 58 (2), 209–219.
- Marple, R.T., Talwani, P., 2000. Evidence for a buried fault system in the coastal plain of the Carolinas and Virginia: Implications for neotectonics in the southeastern United States. *Geol. Soc. Amer. Bull.* 112, 200–220.
- McCarthy, R.J., Bernard, E.N., Legg, M.R., 1993. The Cape Mendocino Earthquake: A local tsunami wakeup call? Coastal Zone 93, Proc. 8th Symposium on Coastal and Ocean Management, Sponsored by the American Shore and Beach Preservation Assn./ASCE, held July 19–23, 1993 in New Orleans, Louisiana, pp. 2812–2828.
- McCulloch, D.S., 1985. Evaluating tsunami potential. *Evaluating Earthquake Hazards in the Los Angeles Region: An Earth Science Perspective*, US Geological Survey Professional Paper, vol. 1360, pp. 374–413.
- Meier, L.S., 1993. The susceptibility of a gravelly soil site to liquefaction. M. S. thesis, Blacksburg, Virginia Polytechnic Institute and State University, 74 pp.
- Meisling, K.E., 1979. Possible emplacement history of a sandblow structure at Pallett Creek, California. In: Abbott, P.L. (Ed.), *Geological Excursions in the Southern California Area*. Dept. of Geological Sciences, San Diego State University, San Diego, CA, pp. 63–66.
- Merriam, D.F., Forster, A., 2002. Stratigraphic and sedimentological evidence for late Paleozoic earthquakes and recurrent structural movement in the U.S. Midcontinent. In: Ettensohn, F.R., Rast, N., Brett, C.E. (Eds.), *Ancient Seismites*, Spec. Pap.-Geol. Soc. Am., vol. 359, pp. 99–108.
- Michael, A.J., Ross, S.L., Stenner, H.D., 2002. Displaced rocks, strong motion, and the mechanics of shallow faulting associated with the 1999 Hector Mine, California, earthquake. *Bull. Seismol. Soc. Am.* 92 (4), 1561–1569.
- Michetti, A.M., Brunamonte, F., Serva, L., 1995. Paleoseismological evidence in the epicentral area of the January 1968 earthquakes, Belice, southwestern Sicily. In: Serva, L. (Ed.), *Perspectives in Paleoseismology*, Assn. of Eng. Geologists, Spec. Pub., vol. 6, pp. 127–139 (Chapter 10).
- Minoura, K., Nakaya, S., Uchida, M., 1994. Tsunami deposits in a lacustrine sequence of the Sanriku coast, northeast Japan. *Sediment. Geol.* 89 (1–2), 25–31.
- Moore, D.G., 1969. Reflection profiling studies of the California Continental Borderland—structure and quaternary turbidite basins. *Spec. Pap.-Geol. Soc. Am.* 107 (142 pp.).
- Moore, G.W., 1972. Offshore extension of the Rose Canyon Fault, San Diego, California. *U. S. Geol. Surv. Prof. Pap.* 800-C, C113–C116.
- Moore, A.L., Lewis, D.R., 1995. Landward fining in a suspected tsunami deposit, Molokai, Hawaii (abstr.). *Tsunami Deposits: Geologic Warnings of Future Inundations*. US Geological Survey/NOAA Sponsored Conference, held on May 22–23, 1995 at the University of Washington at Seattle, Abstracts with Programs, pp. 28–29.
- Moretti, M., Pieri, P., Tropeano, M., 2002. Late Pleistocene soft-sediment deformation structures interpreted as seismites in paralic deposits in the city of Bari (Apulian foreland, southern Italy). In: Ettensohn, F.R., Rast, N., Brett, C.E. (Eds.), *Ancient Seismites*, Spec. Pap.-Geol. Soc. Am., vol. 359, pp. 75–86.
- Morner, Nils-Axel, 1996. Liquefaction and varve deformation as evidence of paleoseismic events and tsunamis, the autumn 10,430 BP case in Sweden. *Quat. Sci. Rev.* 15, 939–948.
- Morse, W.C., 1941. New Madrid earthquake craters. *Bull. Seismol. Soc. Am.* 31, 309–319.
- Moyle Jr., W.R., 1973. Geological map of the western part of Camp Pendleton, Southern California. US Geological Survey Open-File Report. In: Ross, A., Dowlen, R.J., (Eds.), *Studies on the geology of Camp Pendleton, and western San Diego County, California*. San Diego Assn. Of Geol. Guidebook for 1975.
- Munson, P.J., Munson, C.A., Pond, E.C., 1995. Paleoliquefaction evidence for a strong Holocene earthquake in south-central Indiana. *Geology* 23, 325–328.
- Namson, J.S., Davis, T.L., 1988. Seismically active fold and thrust belt in the San Joaquin Valley, central California. *Geol. Soc. Amer. Bull.* 100, 257–273.
- Natland, M.L., Kuenen, P.H., 1951. Sedimentary history of the Ventura Basin, California, and the action of turbidity currents. In: Hough, J.L. (Ed.), *Turbidity Currents and the Transportation of Coarse Sediments to Deep Water*, SEPM Spec. Pub., vol. 2, pp. 76–107.
- National Research Council, 1985. *Liquefaction of soils during earthquakes*. National Academy Press, Washington, DC (226 pp.).
- Newcomb, R.C., 1952. Origin of the mima mounds, Thurston County region, Washington. *J. Geol.* 60, 461–472.
- Nichols, R.J., 1995. The liquefaction and re-mobilization of sandy sediments. In: Hartley, A.J., Prosser, D.J. (Eds.), *Characterization of Deep Marine Clastic Systems*, Geol. Soc. Spec. Pub., vol. 94, pp. 63–76.
- Nishimura, Y., Miyaji, N., 1995. Spatial distribution and lithofacies of recent tsunami and storm surge deposits (abstr.). *Tsunami Deposits: Geologic Warnings of Future Inundation*. US Geological Survey/NOAA Sponsored Conference, held on May 22–23, 1995 at the University of Washington at Seattle, Abstracts with Programs, p. 4.
- Oakshott, G.B. (Ed.), 1975. The San Fernando, California earthquake of 9 February 1971. *Bull.-Div. Mines Geol.*, vol. 196. California Dept. of Conservation (463 pp.).
- Obermeier, S.F., 1987. Identification and geologic characteristics of earthquake-induced liquefaction features. In: Crone, A.J., Omdahl, E.M. (Eds.), *Directions in Paleoseismology*, Open-File Rep. (U. S. Geol. Surv.), vol. 87-673, pp. 173–177.
- Obermeier, S.F., 1989. The New Madrid earthquakes: an engineering-geologic interpretation of relict liquefaction features. *U. S. Geol. Surv. Prof. Pap.* 1336-B (114 pp.).
- Obermeier, S.F., 1996a. Using liquefaction-induced features for paleoseismic analysis. In: McCalpin, J.P. (Ed.), *Paleoseismology*. Academic Press Inc., pp. 331–396 (Chap. 7).



- Obermeier, S.F., 1996b. Use of liquefaction-induced features for paleoseismic analysis—an overview of how seismic liquefaction features can be distinguished from other features and how their regional distribution and properties of source sediment can be used to infer the location and strength of Holocene paleo-earthquakes. *Eng. Geol.* 44, 1–76.
- Obermeier, S.F., Dickenson, S.E., 2000. Liquefaction evidence for the strength of ground motions resulting from Late Holocene cascadian subduction earthquakes, with emphasis on the event of 1700 AD. *Bull. Seismol. Soc. Am.* 90, 876–896.
- Obermeier, S.F., Pond, E.C., 1999. Issues in using liquefaction features for paleoseismic analysis. *Seismol. Soc. Res. Lett.* 70 (1), 34–58.
- Obermeier, S.F., Gohn, G.S., Weems, R.E., Gelinas, R.L., Rubin, M., 1985. Geologic evidence for recurrent moderate to large earthquakes near Charleston, South Carolina. *Science* 227, 408–411.
- Obermeier, S.F., Jacobson, R.B., Smoot, J.P., Weems, R.E., Gohn, G.S., Monroe, J.E., Powers, D.S., 1990. Earthquake-induced liquefaction features in the coastal setting of South Carolina and in the fluvial setting of the New Madrid seismic zone. *U. S. Geol. Surv. Prof. Pap.* 1504 (44 pp.).
- Obermeier, S.F., Bleuer, N.R., Munson, C.A., Munson, P.J., Martin, W.S., McWilliams, K., Tabaczynski, D.A., Odum, J.K., Rubin, M., Eggert, D.L., 1991. Evidence of strong earthquake shaking in the lower Wabash Valley from prehistoric liquefaction features. *Science* 251, 1061–1063.
- Obermeier, S.F., Martin, J.R., Frankel, A.D., Youd, T.L., Munson, P.J., Munson, C.A., Pond, E.C., 1993. Liquefaction evidence for one or more strong Holocene earthquakes in the Wabash Valley of southern Indiana and Illinois, with a preliminary estimate of magnitude. *U. S. Geol. Surv. Prof. Pap.* 1536 (27 pp.).
- Obermeier, S.F., Pond, E.C., Olson, S.M., Green, R.A., 2002. Paleoliquefaction studies in continental settings. In: Ettensohn, F.R., Rast, N., Brett, C.E. (Eds.), *Ancient Seismites, Spec. Pap.-Geol. Soc. Am.*, vol. 359, pp. 13–27.
- Obermeier, S.F., Durbin, J.M., Pouch, G., 2002. Ground fissuring as a paleoseismic indicator: (abstr.). *Geol. Soc. of Am., Northeastern Section (39th Annual Meeting) and Southeastern Section (53rd Annual Meeting)*, Wash. D.C. March.
- Okal, E.A., Fryer, G.J., Borrero, J.C., Ruscher, C., 2002a. The landslide and local tsunami of 13 September 1999 on Fatu Hiva (Marquesas Islands; French Polynesia). *Bull. Soc. Geol. Fr.* 173 (4), 359–367.
- Okal, E.A., Synolakis, C.E., Fryer, G.J., Heinrich, P., Borrero, J.C., Ruscher, C., Arcas, D., Guille, G., Rousseau, D., 2002b. A field survey of the 1946 Aleutian tsunami in the far field. *Seismol. Res. Lett.* 73 (4), 490–503.
- Orcutt, C.R., 1887. Aquatic plants of the vicinity of San Diego. *The West American Scientist* III (6), 123–126.
- Ortiz, M., Singh, S.K., Pacheco, J., Kostogodov, V., 1998. Rupture length of the October 9, 1995 Colima–Jalisco earthquake (Mw 8) estimated from tsunami data. *Geophys. Res. Lett.* 25 (15), 2857–2880.
- Owen, G., 1987. Deformation processes in unconsolidated sands. In: Jones, M.E., Preston, R.M.F. (Eds.), *Deformation of Sediments and Sedimentary Rocks*, *Geol. Soc. Spec. Publ.*, vol. 29. Blackwell Scientific Publications, Oxford [Oxfordshire], pp. 11–24.
- Peters, K.E., Hermann, R.B., 1986. First-hand observations of the Charleston earthquake of August 31, 1886, and other earthquake materials. *Bull.-C. Geol. Surv.* 41, 116.
- Peterson, C., Priest, G.R., 1995. Paleotsunami barrier-overtopping, one piece of the puzzle (abstr.). *Tsunami Deposits: Geologic Warnings of Future Inundations*. US Geological Survey/NOAA Sponsored Conference, held on May 22–23, 1995 at the University of Washington at Seattle, Abstracts with Programs, p. 35.
- Pratt, B.R., 2002. Storms versus tsunamis: dynamic interplay of sedimentary, diagenetic, and tectonic processes in the Cambrian of Montana. *Geology* 30 (5), 423–426.
- Prentice, C.S., Schwartz, D.P., 1991. Re-evaluation of 1906 surface faulting, geomorphic expression, and seismic hazard along the San Andreas Fault in the southern Santa Cruz Mountains. *Bull. Seismol. Soc. Am.* 81, 1424–1479.
- Prentice, C.S., Keefer, D.K., Sims, J.D., 1992. Surface effects of the earthquakes. *The Cape Mendocino Earthquakes, Earthquake Volcanoes*, vol. 23, pp. 127–134.
- Rajendran, C.P., Rajendran, K., 2003. Studying earthquake recurrence in the Kachchh region, India. *Eos*, 84 (48): pp. 529, 533, 536.
- Reimnitz, E., Marshall, N.F., 1965. Effects of the Alaska earthquake and tsunami on recent deltaic sediments. *J. Geophys. Res.* 70 (10), 2363–2376.
- Reiter, L., 1995. Paleoseismology—a user's perspective. In: Serva, L., Slemmons, D.B. (Eds.), *Perspectives in Paleoseismology*, *Assn. of Eng. Geol. Spec. Pub.*, vol. 6, pp. 3–17.
- Riefner Jr., R.E., Pryor, D.R., 1996. New locations and interpretations of vernal pools in southern California. *Phytologia* 80 (4), 296–327.
- Ritchie, A.M., 1952. The erosional origin of the mima mounds of southwest Washington. *J. Geol.* 61, 41–50.
- Rivero, C.J., Shaw, J.H., Mueller, K., 2000. Oceanside and thirty-mile bank blind thrusts: implications for earthquakes hazards in coastal southern California. *Geology* 28 (10), 891–894.
- Rymer, M.J., Seitz, G.G., Weaver, K.D., Orgil, A., Faneros, G., Hamilton, J.C., Goetz, C., 2002a. Geologic and paleoseismic study of the Lavic Lake Playa, Mojave Desert, California. *Bull.-Seismol. Soc. Am.* 92, 1577–1591.
- Rymer, M.J., Boatwright, J., Seekins, L.C., Yule, J.D., Liu, J., 2002b. Triggered surface slips in the Salton Trough associated with the 1999 Hector Mine, California earthquake. *Bull. Seismol. Soc. Am.* 92 (4), 1300–1317.
- San Diego Association of Governments (SANDAG), 1991. *Shoreline Preservation Strategy for the San Diego Region (Draft)*. San Diego Association of Governments (SANDAG), California (63 pp.).
- Saucier, R.T., 1987. Geomorphological interpretations of late Quaternary terraces in Western Tennessee and their regional tectonic implications. In: Russ, D.P., Crone, A.J. (Eds.), *The New Madrid, Missouri, Earthquake Region—Geological, Seismological, and Geotechnical Studies*, *U. S. Geol. Surv. Prof. Pap.*, vol. 1336-A (19 pp.).

- Saucier, R.T., 1991a. Comment on formation of mima mounds: a seismic hypothesis. *Geology* 19 (3), 284.
- Saucier, R.T., 1991b. Geoarchaeological evidence of strong prehistoric earthquakes in the New Madrid (Missouri) seismic zone. *Geology* 19, 296–298.
- Saul, R.B., 1975. Geology of the southeast slope of the Santa Susana Mountains and geological effects of the San Fernando earthquake. In: Oakshott, G.B. (Ed.), *San Fernando Earthquake of 9 February, 1971*, Bull. - Calif., Div. Mines Geol., vol. 196, pp. 53–70 (Chap. 5).
- Scheffers, A., Kelletat, D., 2003. Sedimentologic and geomorphologic tsunami imprints; a worldwide review. *Earth-Sci. Rev.* 63, 83–92.
- Scott, R.F., Zuckerman, K.A., 1973. Sandblows and liquefaction. The Great Alaska Earthquake of 1964. National Research Council, 6: Engineering volume. National Academy of Science, Washington, DC.
- Seed, H.B., 1968. Landslides during earthquakes due to soil liquefaction. *Am. Soc. Civil Eng., J. Soil Mech. Found. Div.* 94, 1055–1122.
- Seed, H.B., Idriss, I.M., 1983. Evaluation of liquefaction potential using field performance data. *J. Geotech. Eng.* 109 (3), 458–482.
- Seed, H.B., Lee, K.L., 1966. Liquefaction of saturated sands during cyclic loading. *Am. Soc. Civil Eng., J. Soil Mech.* 92 (SM6), 105–134 (paper 4972).
- Seilacker, A., 1969. Fault-graded beds interpreted as seismites. *Sedimentology* 13, 155–159.
- Seilacker, A., 1984. Sedimentary structures tentatively attributed to seismic events. *Mar. Geol.* 55, 1–12.
- Seitz, G.G., 1983. Normal faulting associated with strike-slip faulting in the Leucadia area of San Diego County. Unpubl. *Geology Senior thesis*, San Diego State University, San Diego, California, 32 pp.
- Shackleton, N.J., Opdike, N.D., 1976. Oxygen-isotope and paleomagnetic stratigraphy of pacific core V28–239 late Pliocene to latest Pleistocene. *Mem. Geol. Soc. Amer.* 145, 449–464.
- Shaw, J., Shearer, P., 2000. An elusive blind-thrust beneath metropolitan Los Angeles. *Science* 283, 1516–1518.
- Shepard, F.P., 1948. *Submarine Geology*, first edition Harper and Brothers Publishers (348 pp.).
- Shepard, F.P., Emery, K.O., 1941. Submarine topography off the southern California coast: canyons and tectonic interpretation. *Spec. Pap.-Geol. Soc. Am.* 31 (171 pp.).
- Shepard, F.P., Kuhn, G.G., 1977. Geological and historical analysis of coastal zone environmental hazards and liability for losses caused by them, Carlsbad to Oceanside, California. US Department of Commerce, NOAA/SEA GRANT, Project. No. R/CZ-43.
- Shepard, F.P., Macdonald, G.A., Cox, D.C., 1950. The tsunami of April 1, 1946. *Bull. of the Scripps Institution of Oceanography of the University of California*, Univ. of California Press, Berkeley and Los Angeles, California, 5 (6), 391–528.
- Shi, S., Dawson, A.G., Smith, D.E., 1993. Geomorphological impact of the Flores tsunami of 12th December, 1992. *Tsunami* 93, Proc. IUGG/IOC Int. Tsunami Symposium, held in Wakayama, Japan on August 23–27, pp. 689–696.
- Shlemon, R.J., 1977. Geomorphic analysis of Fault “E”, Camp Pendleton, California. Unpubl. report prepared for Southern California Edison Company (Rosemead), California. Reprinted In: Legg, M.R., Kuhn, G.G., Shlemon R.J. (Eds.), *Neotectonics and Coastal Instability: Orange and Northern San Diego Counties, California*. Joint field conference Am. Assn. of Petroleum Geologists, Pacific Section, and SPE, Western Section held on June 19–22, 2000 in Long Beach, California, 2: pp. II-163 to II-182.
- Shlemon, R.J., 1978a. Late Quaternary evolution of the Camp Pendleton -San Onofre State Beach area, California. Unpubl. report prepared for Southern California Edison Company (Rosemead), California. Reprinted In: Legg, M.R., Kuhn, G.G., Shlemon, R.J. (Eds.), *Neotectonics and Coastal Instability: Orange and Northern San Diego Counties, California*. Joint field conference, Am. Assn. of Petroleum Geologists, Pacific Section, and SPE, Western Section, held in Long Beach, California on June 19–22, 2000, 2: pp. II-183 to II-251.
- Shlemon, R.J., 1978b. Late Quaternary rates of deformation, Laguna Beach to San Onofre State Beach, Orange and San Diego Counties, California. Unpubl. report prepared for Southern California Edison Company (Rosemead), California. Reprinted In: Legg, M.R., Kuhn, G.G., Shlemon, R.J. (Eds.), *Neotectonics and Coastal Instability: Orange and Northern San Diego Counties, California*. Joint field conference, Am. Assn. of Petroleum Geologists, Pacific Section, and SPE, Western Section, held in Long Beach, California on June 19–22, 2000, 2: pp. II-252 to II-282.
- Shlemon, R.J., 1979a. Late Quaternary rates of sedimentation and soil formation, Camp Pendleton and San Onofre State Beach coastal area, southern California. Unpubl. report prepared for Southern California Edison Company (Rosemead), California. Reprinted In: Legg, M.R., Kuhn, G.G., Shlemon, R.J. (Eds.), *Neotectonics and Coastal Instability: Orange and Northern San Diego Counties, California*. Joint field conference, Am. Assn. of Petroleum Geologists, Pacific Section, and SPE, Western Section, held in Long Beach, California, on June 19–22, 2000, 2: p. II-283.
- Shlemon, R.J., 1979b. Late Cenozoic stratigraphy: Capistrano embayment coastal area, Orange County, California. Unpubl. report prepared for Southern California Edison Company (Rosemead), California. Reprinted In: Legg, M.R., Kuhn, G.G., Shlemon, R.J. (Eds.), *Neotectonics and Coastal Instability: Orange and Northern San Diego Counties, California*. Joint field conference, Am. Assn. of Petroleum Geologists, Pacific Section, and SPE, Western Section, held in Long Beach, California, on June 19–22, 2000, 2: pp. II-284 to II-303.
- Shlemon, R.J., 1979c. Age of “Dana Poin”, “Vaciadero”, and “Carr” faults, Capistrano Embayment coastal area, Orange County, California. Unpubl. report prepared for Southern California Edison Company (Rosemead), California. Reprinted In: Legg, M.R., Kuhn, G.G., Shlemon, R.J. (Eds.), *Neotectonics and Coastal Instability: Orange and Northern San Diego Counties, California*. Joint field conference, Am. Assn. of Petroleum Geologists, Pacific Section, and SPE, Western Sec-

- tion, held in Long Beach, California on June 19–22, 2000, 2: pp. II-304 to II-322.
- Shlemon, R.J., 1979d. Late Pleistocene channel of the lower Santa Margarita River, San Diego County, California. Reprinted In: Legg, M.R., Kuhn, G.G., Shlemon, R.J. (Eds.), *Neotectonics and Coastal Instability: Orange and Northern San Diego Counties, California*. Joint field conference, Amer. Assn. of Geologists, Pacific Section, and SPE, Western Section, held in Long Beach California on June 19–22, 2000, 2: pp. II-323 to II-329.
- Shlemon, 1999. Pers. com.
- Shlemon, R.J., 2000. State-of-the-art to standard-of-practice: Active faults, paleoliquefaction, and tsunamis in the Carlsbad area, San Diego County, California. *Geol. Soc. of Am. Ann. Meeting*, held in Reno, Nevada on November 9–18, 2000, Abstracts with Programs, Session no. 54, T69, p. A121.
- Shlemon, R.J., Kuhn, G.G., 1997. Seismically-induced liquefaction increased by urbanization, northern San Diego County, California (abstr.). *Geol. Soc. of Amer. Meeting, Cordilleran Section*, held on May 21–23, 1997 at Kona, Hawaii, Abstracts with Programs, 29 (5): poster session no. 23, (#16943), pp. 24/48.
- Shlemon, R.J., Begg, E.L., Huntington, G.L., 1973. Fracture traces. *Pac. Discovery*, (California Academy of Sciences San Francisco, California) XXVI (1), 31–32.
- Shlemon, R.J., Kuhn, G.G., Boka, B., Riefner, R.E., 1997. Origin of a mima-mound field, San Clemente State Park, Orange County, California: A test for the seismic hypothesis. *Assn. of Engineering Geologists, 40th Annual Meeting*, held on September 30 to October 4, 1997 at Portland, Oregon, Abstracts with Programs, pp. 148–149.
- Sieh, K.E., 1978. Prehistoric large earthquakes produced by slip on the San Andreas fault at Palmett Creek, California. *J. Geophys. Res.* 83, 3907–3939.
- Sims, J.D., Garvin, C.D., 1995. Recurrent liquefaction induced by the 1989 Loma Prieta earthquake and 1990 and 1991 aftershocks: implications for paleoseismicity studies. *Bull. Seismol. Soc. Am.* 85 (1), 51–65.
- Sliter, R.W., Ryan, H.F., Normark, W.R., 2001. Does recent deformation at the base of slope provide evidence of a connection between the Newport–Inglewood and the Rose Canyon fault zones offshore southern California? *Am. Geophys. Union (AGU) Fall Meeting* 82 (47), F802.
- Slosson, J.E., Kuhn, G.G., McCarthy, R.J., Legg, M.R., 2001. Use of space photography for fault and geomorphic studies. *Geol. Soc. of Am., Cordilleran Section, 96th annual meeting* held on April 27–29, 2000, in Vancouver, British Columbia, Session no. 80051.
- Smith, B.F., 1996. Results of a data recovery program at sites SDI-6,132,SDI-10,671, and SDI-12,814; The Carlsbad Ranch Project, Carlsbad, California. Unpubl. report prepared for the Carls Development Company, Carlsbad, California, 42 pp., 80 tables, 6 appendices.
- Society of Economic Paleontologists and Mineralogists (SEPM), 1951. Turbidity currents and the transportation of coarse sediments to deep water. In: Hough, J.L. (Ed.), *A Symposium, SEPM Spec. Pub.*, vol. 2 (106 pp.).
- Stewart, K.G., Dennison, J.M., Bartholomew, M.J., 2002. Late Mississippian paleoseismites from southeastern West Virginia and southwestern Virginia. In: Ettensohn, F.R., Rast, N., Brett, C.E. (Eds.), *Ancient Seismites*, pp. 127–144.
- Stone, E.M., Grant, L.B., Arrowsmith, J.R., 2002. Recent rupture history of the San Andreas fault southeast of Cholame in the northern Carrizo Plain, California. *Seismol. Soc. Am.* 92 (3), 983–997.
- Su, Wen-June, Follmer, L.R., 1992. Soft-sediment structures attributed to paleo-earthquakes in southeastern Illinois —A preliminary interpretation. *Proc. of the Assn. of Engineering Geologists, 35th annual meeting* held in Los Angeles, California on October 2–4, pp. 459–468.
- Sylvester, A.G., 1988. Strike-slip faults. *Geol. Soc. Amer. Bull.* 100, 1666–1703.
- Sylvester, A.G., Burmeister, K.C., Wise, W.S., 2002. Faulting and effects of associated shaking at Pisgah Crater Volcano caused by the 16 October 1999 Hector Mine earthquake (Mw 7.1), central Mojave Desert, California. *Bull. Seismol. Soc. Am.* 92 (4), 1333–1340.
- Synolakis, C.E., Carrier, P., Liu, G., Yeh, H., 1997. Tsunamigenic sea-floor deformations. *Science* 278, 598–600.
- Takashimizu, Y., Masuda, F., 2000. Depositional facies and sedimentary successions of earthquake-induced tsunami deposits in upper Pleistocene incised valley fills, central Japan. *Sediment. Geol.* 135, 295–320.
- Talwani, P., Cox, J., 1985. Paleoseismic evidence for recurrence of earthquakes near Charleston, South Carolina. *Science* 229, 379–380.
- Tan, S.S., 1986. Landslide hazards in the Encinitas Quadrangle, San Diego County, California. *California Div. Mines and Geology Open-File Report 86-8 LA*, 3 plates, scale 1:24,000.
- Tan, S.S., Kennedy, M.P., 1996. Geological maps of the northwestern part of San Diego County, California. *California Div. Mines and Geology Open-File Report 96-02*, 2 plates, scale 1:24,000.
- Tuttle, M.P., Ruffman, A., Anderson, T., Jeter, H., 1995. Comparison of tsunami and storm deposits along the Atlantic Seaboard (abstr.). *Tsunami Deposits: Geologic Warnings of Future Inundation*. US Geological Survey/NOAA sponsored conference, held on May 22–23, 1995 at the University of Washington at Seattle, Abstracts with Programs, pp. 9–10.
- Tuttle, M.P., Lafferty III, R.H., Guccione, M.J., Schweig III, E.S., Lopinot, N., Cande, R.F., Dyer-Williams, K., Hayes, M., 1996. Use of archaeology to date liquefaction features and seismic events in the New Madrid seismic zone, Central United States. *Geoarchaeology* 11, 451–480.
- Twidale, C.R., 1976. *Analysis of Landforms*. Wiley and Sons, Australasia Pty. Ltd., pp. 197–201.
- US Department of Agriculture (USDA), 1918. Reconnaissance soil survey of the San Diego Region, California. In: Holmes, L.C., Pendleton, R.L. (Authors, Eds.), *US Government Printing Office*, Wash. D.C., 77 pp.
- US Department of Agriculture (USDA), 1953. Aerial photograph No. AXM-8M-100.
- US Department of Agriculture (USDA), 1973. *Soil Survey: San Diego Area, California*. US Government Printing Office, Wash. D.C., 2 parts, I: (113p.), II: (127p.), 76 soil maps.



- US Geological Survey (USGS), 1947. Aerial photograph GS-CP, No. 11–167.
- US Geological Survey (USGS), 1990a. The Loma Prieta, California, earthquake: an anticipated event. *Science* 247, 286–293.
- US Geological Survey (USGS), 1990b. Tecumseh's Prophecy: Preparing for the next New Madrid earthquake —A plan for an intensified study of the New Madrid Seismic Zone. In: Hamilton, R.M., Johnson, A.C. (Eds.), *US Geological Survey Circular*, vol. 1066 (29 pp.).
- US Geological Survey (USGS), 1992. The Loma Prieta, California earthquake of October 17, 1989 —Marina District. In: O'Rourke, T.D. (Ed.), *U. S. Geol. Surv. Prof. Pap.*, 1551-F, pp. F1–F215.
- US Geological Survey (USGS), 1998. The Loma Prieta, California earthquake of October 17, 1989 —Liquefaction. In: Holzer, T.J. (Ed.), *U. S. Geol. Surv. Prof. Pap.*, 1551-B, pp. B1–B314.
- US Geological Survey (USGS), Southern California Earthquake Center (SCEC), 1994. The magnitude 6.7 Northridge, California, earthquake of 17 January, 1994. *Science* 266, 389–397.
- Van Dorn, W.G., 1965. Tsunamis. *Advances in Hydrosience*, vol. 2. Academic Press, New York, pp. 1–48.
- Vaughan, P.R., Thorup, K.M., Rockwell, T.K., 1999. Paleoseismology of the Elsinore fault at Agua Tibia Mountain, southern California. *Bull. Seismol. Soc. Am.* 89 (6), 1447–1457.
- Washburn, A.L., 1997. Plugs and plug circles: a basic form of patterned ground, Cornwallis Island, Arctic Canada—origin and implications. *Mem. Geol. Soc. Amer.* 190 (97 pp.).
- Weber Jr., F.H., 1982. Recent slope failures, ancient landslides, and related geology of the north-central coastal area, San Diego County, California. *California Div. Mines and Geol. Open-File Report* 82-12 LA, 77 pp.
- Weldon II, R., Sieh, J., 1985. Holocene rate of slip and tentative interval for large earthquakes on the San Andreas fault, Cajon Pass, southern California. *Geol. Soc. Amer. Bull.* 96, 793–812.
- Weldon II, R.J., McCalpin, J.P., Rockwell, T.K., 1996. Paleoseismology in strike-slip tectonic environments. In: McCalpin, J.P. (Ed.), *Paleoseismology*. Academic Press, pp. 271–325 (Chap. 6).
- Western Geophysical, 1972. Final report, southern California Edison and San Diego Gas and Electric Company, San Onofre offshore investigations. Unpubl. report relating to the San Onofre Nuclear Generating Station, Units 2 and 3, Amendment No. 11 to Preliminary Safety Analysis Report, 34 pp.
- Wheeler, R.L., 2002. Distinguishing seismic from nonseismic soft-sediment structures: Criteria from seismic-hazard analysis. In: Ettensohn, F.R., Rast, N., Brett, C.E. (Eds.), *Ancient Seismites*, *Spec. Pap.-Geol. Soc. Am.*, vol. 359, pp. 1–11.
- Wilson, K.L., 1972. Eocene and related geology of a portion of the San Luis Rey and Encinitas Quadrangles, San Diego County, California. [master's thesis], Univ. of California at Riverside, 135 pp.
- Woods, M.C., Seiple, W.R. (Eds), 1995. The Northridge, California Earthquake, California Dept. of Conservation. *Spec. Publ., Div. Mines Geol.* 116 (302 pp.).
- Yegian, M.K., Ghaghrman, V.G., Harutinunyan, R.N., 1994. Liquefaction and embankment failure case histories. *J. Geotech. Geoenviron. Eng. Am. Soc. Civil Eng. (ASCE)* 120 (3), 581–596.
- Youd, T.L., 1973. Liquefaction, flow, and associated ground failure. *U.S. Geol. Surv. Circ.* 688 (12 pp.).
- Youd, T.L., 1984. Ground effects—Liquefaction and associated ground failure. *Proc. in the Geological and Hydrologic Hazards Training Program, US Geological Survey Open-File Report*, vol. 84-760, pp. 210–232.
- Youd, T.L., 1985. Landslides caused by earthquakes: discussion and reply. *Geol. Soc. Amer. Bull.* 96, 1091–1094.
- Youd, T.L., Asce, M., Perkins, D.M., 1978. Mapping liquefaction-induced ground failure potential. *Proc. of the Am. Soc. of Civil Engineers*, 104 (GT4) presented at the October 17–21, 1977 ASCE Annual Convention, Exposition and Continuing Education Program, held at San Francisco, California, *J. Geotech. Eng.*, pp. 433–446.
- Zitellini, N., Memdes, L.A., Cordoba, D., Danobeitia, J., Nicolich, R.N., Pellis, G., Ribeiro, A., R. Sartori, Toelli, L., Bartolome, R., Bortoluzzi, G., Calafato, A., Carrillho, F., Casoni, L., Chierici, F., Corela, C. Correggiari, A., Della Vedova, B., Gracia, E., Jornet, P., Landuzzi, M., Ligi, M., Magagnoli, A., Marozzi, G., Matias, L., Penitenti, D., Rodriguez, P., Rovere, M., Terrinha, P., Vigliotti, L., Ruiz, A.Z., 2001. Source of 1755 Lisbon earthquake and tsunami investigated. *Eos* 82 (26) 285, 290, 291.